

THE ABEL-JACOBI MAP FOR HIGHER CHOW GROUPS, II

MATT KERR AND JAMES D. LEWIS

CONTENTS

1. Introduction	2
1.1. Notation	4
1.2. Acknowledgements	4
2. Absolute Hodge cohomology	4
3. The weight-filtered Abel-Jacobi map	13
3.1. Gysin spectral sequences and higher residue maps	13
3.2. Generalization of the Griffiths prescription	16
3.3. Higher Hodge conjecture	19
3.4. Coniveau filtration	22
4. Higher normal functions	23
4.1. Spreads and the Leray filtration	23
4.2. A geometric construction: the Λ -filtration	25
4.3. Comparing filtrations	28
5. Leray filtration in the product case	32
5.1. Higher Abel-Jacobi maps	32
5.2. Weight filtration redux	35
6. Detecting indecomposables in the regulator kernel	37
6.1. An invariant for cycles with complete spread	37
6.2. First application: exterior products	40
6.3. Relation to a criterion of Lewis	41
6.4. Second application: Pontryagin cycles	43
7. Reduced higher Abel-Jacobi maps	47
8. Abel-Jacobi for motivic cohomology of relative and singular varieties	49
8.1. Currents with pullback	50
8.2. Higher Chow chains with pullback	53
8.3. Abel-Jacobi maps	59
8.4. Regulator period computations on singular varieties	61
References	66

1. INTRODUCTION

This paper has its origin in two distinct, well-established problems: that of detecting $K_n^{\text{alg}}(X)$ classes — especially indecomposable ones — in the regulator kernel for X smooth projective; and that of computing the limit of the regulator for a family of such classes as X degenerates. It has become a sequel to [KLM] because these problems benefit from two “dual” extensions of the regulator formulas introduced there.

To explain this, we recall the relation between Bloch’s higher Chow groups, K -theory and motivic cohomology

$$(1.1) \quad CH^p(X, n)_{\mathbb{Q}} \cong K_n^{\text{alg}}(X)_{\mathbb{Q}}^{(p)} \cong H_{\mathcal{M}}^{2p-n}(X, \mathbb{Q}(p))$$

for X smooth projective. For our purposes, the regulator is the absolute Hodge cycle-class map

$$(1.2) \quad c_{\mathcal{H}}^X : CH^p(X, n)_{\mathbb{Q}} \longrightarrow H_{\mathcal{H}}^{2p-n}(X, \mathbb{Q}(p)),$$

and [KLM] introduced an explicit morphism of complexes, from cycles to “triples” of currents, computing this. Nontrivial cycles $\mathcal{Z} \in \ker(c_{\mathcal{H}}^X)$ can be studied by *spreading out* $(X, \mathcal{Z} \mapsto \mathcal{X}_U, \mathfrak{Z}_U)$ as in §4.1) and computing $c_{\mathcal{H}}^{\mathcal{X}_U}(\mathfrak{Z}_U)$. Here $\mathcal{X}_U$ is quasi-projective but still smooth, and so (1.1), (1.2) still make sense; however, [KLM] does not give a map of complexes computing $c_{\mathcal{H}}$ for quasi-projective varieties.

This is remedied in §3, which as a byproduct also yields a formula for

$$c_{\mathcal{H}, Y} : CH^p(Y, n) \longrightarrow H_{\mathcal{H}, Y}^{2p-n-2}(X, \mathbb{Q}(p+1))$$

when $Y \subset X$ is a complete normal crossing divisor (NCD). But this is the “wrong” regulator for singular varieties, as far as many arithmetic applications or (finite) limits under degeneration are concerned. Indeed, for *general* Y

$$CH^p(Y, n) \cong G_n(Y)_{\mathbb{Q}}^{(p)} \cong H_{2d_Y-2p+n}^{\mathcal{M}, B.M.}(Y, \mathbb{Q}(d_Y - p))$$

is motivic Borel-Moore *homology*; whereas

$$(1.3) \quad c_{\mathcal{H}}^Y : H_{\mathcal{M}}^{2p-n}(Y, \mathbb{Q}(p)) \longrightarrow H_{\mathcal{H}}^{2p-n}(Y, \mathbb{Q}(p))$$

is the interesting map. Again, a morphism of complexes computing this — for many types of (complete) singular and relative varieties — is the subject of §§8.1–3. The construction involves partially dualizing that of §3, as well as considerably greater technical difficulties.

Applications of the second construction get relatively short shrift in this paper, but §8.4 does compute (1.3) for higher cycles on a degenerate elliptic curve and $K3$ surface — examples related to irrationality of $\zeta(2)$ and $\zeta(3)$, respectively, see [Ke5]. (The cycles actually do deform

in each case as the variety “smooths”.) The formulas of §8 are also used in [DK].

On the other hand, almost half the paper (§§4–7) concerns applications of $c_{\mathcal{H}}$ for smooth quasi-projective varieties, to define and compute *higher Abel-Jacobi invariants* for higher Chow cycles¹ (on projective X) in the kernel of (1.2). These are, roughly speaking, graded pieces of $c_{\mathcal{H}}^{X_U}(\mathfrak{Z}_U)$ under the Leray filtration for a smooth morphism $\mathcal{X}_U \rightarrow U$ (in the limit as we shrink U). These higher invariants appear in various guises: in §4, as a generalization of the classical notion of normal functions; in §5, as Künneth components of the [KLM] map; and in §7, as generalized membrane integrals on X in the spirit of [GG] and [Ke1, sec. 11].

The actual double-complex formalism used in the formula for $c_{\mathcal{H}}$ in §3, only shows up once in the higher AJ discussion — in §5.2 to prove a very precise result on nontrivial *decomposable* cycles in the regulator kernel (Theorem 5.1). These are detected by composing the higher AJ ’s with *higher residue maps*, a feature not present when spreading out usual algebraic cycles. But the most beautiful results on (higher) cycles are to be found in §6, where we prove that certain regulator-trivial exterior products (Theorem 6.2) and Pontryagin products (Theorem 6.3) of higher Chow cycles are strongly indecomposable, i.e not in the image of

$$CH^{p-1}(X_K, n-1) \otimes K^* \longrightarrow CH^p(X_K, n).$$

The “indecomposable higher AJ ” invariant used to detect these (Theorem 6.1) is an improvement of an earlier invariant of the second author [Le2], and Theorem 6.3 is likewise a strengthening of a result of Collino and Fakhruddin [CF] which used [Le2].

Section 2 is partly expository: we show how to compute $H_{\mathcal{H}}^q(\mathcal{W}, \mathbb{Q}(p))$ ($\mathcal{W}$ smooth quasi-projective) as the cohomology of a double complex, then place a “weight” filtration on it and compute the graded pieces (Prop. 2.7). This is all just to clear the runway for §3.

We want this paper (in conjunction with [KLM]) to be useful as a handbook of sorts on various cycle-class maps, so we have attempted a “modular” format as far as possible. With the following exceptions, a well-informed reader can probably start at the beginning of any section: §3 depends on §2, §5.2 on §3.1, and §6 and §7 each on §5.1. Our definition of “well-informed” includes familiarity with (or at least reference to) §§5.1–5.7 of [KLM], which is used almost everywhere.

¹to be clear: [KLM] studied usual AJ on higher cycles; [GG, Le1, Ke5] studied higher AJ on usual (algebraic) cycles; here we study higher AJ on higher cycles, which includes both.

1.1. Notation. Here we forewarn the reader of some potentially confusing idiosyncracies. Most importantly: **cycle groups throughout this paper are automatically** $\otimes \mathbb{Q}$, in particular the higher precycles $Z^p(X, m)$, $Z_{\mathbb{R}}^p(X, m)$ (see §8.2 for a definition of the latter). These are not to be confused with the “topological” C^∞ chains resp. cycles with coefficients in $(2\pi i)^q \mathbb{Q}$, written $C_{top}^r(X_{\mathbb{C}}^{an}, \mathbb{Q}(q))$ resp. $Z_{top}^r(X_{\mathbb{C}}^{an}, \mathbb{Q}(q))$. Here r refers to *real* codimension (in contrast to p). The sheaf of C^∞ forms of degree q is denoted $\Omega_{X^\infty}^q$; currents $\mathcal{D}_X^{2\dim(X)-q}$ are the distributions that act on them.

A higher Chow precycle $\mathcal{Z} \in Z^p(X, m)_{\partial_{\mathcal{B}}\text{-closed}}$ has rational equivalence class $\langle \mathcal{Z} \rangle \in CH^p(X, m)$ and homological equivalence class $[\mathcal{Z}] \in Hom_{\text{MHS}}(\mathbb{Q}(0), H^{2p-n}(X, \mathbb{Q}(p)))$. When it is not crucial, we are sometimes lax about the $\mathcal{Z} \leftrightarrow \langle \mathcal{Z} \rangle$ distinction.

Given a Hodge structure $\mathcal{H}$, $F_h^q \mathcal{H}$ is defined to be the largest sub-Hodge structure of $\mathcal{H}$ contained in $\mathcal{H} \cap F^q \mathcal{H}_{\mathbb{C}}$. If $\mathcal{W}$ is smooth quasi-projective with good compactification $\overline{\mathcal{W}}$, then for our various cycle/cohomology groups, an underline denotes the image of the restriction from $\overline{\mathcal{W}}$ to $\mathcal{W}$: e.g., $\underline{CH}^p(\mathcal{W}, m) := \text{im}\{CH^p(\overline{\mathcal{W}}, m) \rightarrow CH^p(\mathcal{W}, m)\}$.

The following is more standard: for a complex $(\mathcal{K}^\bullet, d)$, the truncation $\tau_{\bullet \leq q} \mathcal{K}^\bullet$ is $\mathcal{K}^\bullet$ for $\bullet < q$, 0 for $\bullet > q$, and $\{\ker(d) \subset \mathcal{K}^q\}$ for $\bullet = q$. $D^b(\text{---})$ is the bounded derived category, and $\mathbf{s}^\bullet(\text{---})$ is the simple complex associated to a double complex.

1.2. Acknowledgements. This paper was conceived at the May 2005 Antalya Algebra Days in Antalya, Turkey; both authors would like to thank the organizers for a stimulating atmosphere. We would also like to thank Donu Arapura for sharing [Ar2], and Shuji Saito for helpful conversations.

2. ABSOLUTE HODGE COHOMOLOGY

Let X be a smooth projective analytic variety $/\mathbb{C}$, $Y = Y_1 \cup \dots \cup Y_N \subset X$ a normal-crossing divisor (NCD) with smooth components (and smooth intersections of all orders), and $\mathbb{A} \subseteq \mathbb{R}$ a subring such that $\mathbb{A} \otimes_{\mathbb{Z}} \mathbb{Q}$ is a field. The principal aim of this section is to produce explicit complexes whose cohomologies compute the absolute $\mathbb{A}$ -Hodge cohomology groups

$$H_{\mathcal{H}}^q(X, \mathbb{A}(p)), \quad H_{\mathcal{H}}^q(X \setminus Y, \mathbb{A}(p)), \quad H_{\mathcal{H}, Y}^q(X, \mathbb{A}(p))$$

as defined in [Ja1, sec. 2]. At least half of the section is expository, albeit brief and sketchy.

Definition 2.1. [Ja1],[BZ] A mixed $\mathbb{A}$ -Hodge complex consists of the following data:

- A bounded below complex $K_{\mathbb{A}}^{\bullet}$ of $\mathbb{A}$ -modules (in the derived category), such that $H^p(K_{\mathbb{A}}^{\bullet})$ is an $\mathbb{A}$ -module of finite type for all p .
- A bounded below filtered complex $(K_{\mathbb{A} \otimes \mathbb{Q}}^{\bullet}, W)$ of $\mathbb{A} \otimes_{\mathbb{Z}} \mathbb{Q}$ -vector spaces, and an isomorphism $K_{\mathbb{A} \otimes \mathbb{Q}}^{\bullet} \xrightarrow{\sim} K_{\mathbb{A}}^{\bullet} \otimes \mathbb{Q}$ in the derived category.
- A bifiltered complex $(K_{\mathbb{C}}^{\bullet}, W, F)$ of $\mathbb{C}$ -vector spaces, and a filtered isomorphism $\alpha : (K_{\mathbb{C}}^{\bullet}, W) \xrightarrow{\sim} (K_{\mathbb{A} \otimes \mathbb{Q}}^{\bullet}, W) \otimes \mathbb{C}$ in the filtered derived category.

Further,

- For every $m \in \mathbb{Z}$,

$$Gr_m^W K_{\mathbb{A} \otimes \mathbb{Q}}^{\bullet} \rightarrow (Gr_m^W K_{\mathbb{C}}^{\bullet}, F)$$

is a (polarizable) $\mathbb{A} \otimes \mathbb{Q}$ -Hodge complex of weight m , i.e. the differentials of $Gr_m^W K_{\mathbb{C}}^{\bullet}$ are strictly compatible with the induced filtration F , and F induces a pure (polarizable) $\mathbb{A} \otimes \mathbb{Q}$ -Hodge structure of weight $m + r$ on $H^r(Gr_m^W K_{\mathbb{A} \otimes \mathbb{Q}}^{\bullet})$ for $r \in \mathbb{Z}$.

A mixed $\mathbb{A}$ -Hodge complex gives rise to a diagram:

(2.1)

$$\begin{array}{ccccc} & & 'K_{\mathbb{A} \otimes \mathbb{Q}}^{\bullet} & & ('K_{\mathbb{C}}^{\bullet}, W) \\ & \nearrow \alpha_1 & & \nwarrow \alpha_2 & \nearrow \beta_1 \\ K_{\mathbb{A}}^{\bullet} & & (K_{\mathbb{A} \otimes \mathbb{Q}}^{\bullet}, W) & & (K_{\mathbb{C}}^{\bullet}, W, F), \\ & & \nwarrow \beta_2 & & \end{array}$$

where $\alpha_j, \beta_j, j = 1, 2$, are morphisms of complexes, α_2 being a quasi-isomorphism, β_1 a filtered morphism, and β_2 a filtered quasi-isomorphism. By the work of Deligne and Beilinson, the construction of mixed $\mathbb{A}$ -Hodge complexes is equivalent to the construction of mixed $\mathbb{A}$ -Hodge structures.

Let $\mu : (M^{\bullet}, d_M) \rightarrow (N^{\bullet}, d_N)$ be a morphism of complexes. The cone of μ is the complex:

$$Cone\{M^{\bullet} \xrightarrow{\mu} N^{\bullet}\} = C_{\mu}^{\bullet} := M[1]^{\bullet} \oplus N^{\bullet},$$

with differential $D(a, b) = (-d_M(a), \mu(a) + d_N(b))$. The absolute Hodge cohomology $H_{\mathcal{H}}^{\ell}(K^{\bullet})$ (of a mixed $\mathbb{A}$ -Hodge complex $K^{\bullet}$) is given by H^{ℓ} of

$$(2.2) \quad Cone\{K_{\mathbb{A}}^{\bullet} \oplus \hat{W}_0 K_{\mathbb{A} \otimes \mathbb{Q}}^{\bullet} \oplus (\hat{W}_0 \cap F^0) K_{\mathbb{C}}^{\bullet} \xrightarrow{(\alpha, \beta)} 'K_{\mathbb{A} \otimes \mathbb{Q}}^{\bullet} \oplus \hat{W}_0 'K_{\mathbb{C}}^{\bullet}\}[-1],$$

where

$$(\alpha, \beta) (\xi_{\mathbb{A}}, \xi_{\mathbb{Q}}, \xi_{\mathbb{C}}) = (\alpha_1 \xi_{\mathbb{A}} - \alpha_2 \xi_{\mathbb{Q}}, \beta_1 \xi_{\mathbb{Q}} - \beta_2 \xi_{\mathbb{C}}),$$

and $\hat{W}_{\bullet}$ is the filtration decalée [De2, sec. 2]. More precisely, set $'W^{\bullet} := W_{-\bullet}$, $\hat{W}_{\bullet} := (\text{Dec } 'W)^{-\bullet}$, where

$$(\text{Dec } 'W)^{\ell} K^m := \ker \left\{ 'W^{\ell+m} K^m \xrightarrow{d_K} \frac{K^{m+1}}{'W^{\ell+m+1} K^{m+1}} \right\}.$$

We henceforth assume $K_{\mathbb{A}}^{\bullet} = 'K_{\mathbb{A}}^{\bullet}$, $K_{\mathbb{C}}^{\bullet} = 'K_{\mathbb{C}}^{\bullet}$ with α_2, β_2 identity maps, and $K_{\mathbb{A} \otimes \mathbb{Q}}^{\bullet} = K_{\mathbb{A}}^{\bullet} \otimes_{\mathbb{Z}} \mathbb{Q}$ with α_1 the inclusion.

Remark 2.2. Suppose also $\mathbb{A} = \mathbb{Q}$, so $\alpha_1 = \text{identity}$. In this case, there is a quasi-isomorphism

$$(*) \quad \text{Cone} \{ \hat{W}_0 K_{\mathbb{Q}}^{\bullet} \oplus F^0 \hat{W}_0 K_{\mathbb{C}}^{\bullet} \xrightarrow{\beta_1 - \beta_2} \hat{W}_0 K_{\mathbb{C}}^{\bullet} \} [-1]$$

$$\downarrow \simeq$$

$$\text{Cone} \{ K_{\mathbb{Q}}^{\bullet} \oplus \hat{W}_0 K_{\mathbb{Q}}^{\bullet} \oplus F^0 \hat{W}_0 K_{\mathbb{C}}^{\bullet} \xrightarrow{(\alpha, \beta)} K_{\mathbb{Q}}^{\bullet} \oplus \hat{W}_0 K_{\mathbb{C}}^{\bullet} \} [-1]$$

given by $(\xi_1, \xi_2, \xi_3) \mapsto (\xi_1, \xi_1, \xi_2, 0, \xi_3)$. (The differential in $(*)$ is $D(\xi_1, \xi_2, \xi_3) = (-d\xi_1, -d\xi_2, d\xi_3 + \beta_1 \xi_1 - \beta_2 \xi_2)$.) We refer to $(*)$ as the normalized AH complex. It has an obvious (non-quasi-isomorphic) embedding in the related “Deligne” (or weak AH) complex

$$\text{Cone} \left\{ K_{\mathbb{Q}}^{\bullet} \oplus F^0 K_{\mathbb{C}}^{\bullet} \xrightarrow{\beta_1 - \beta_2} K_{\mathbb{C}}^{\bullet} \right\} [-1],$$

which simply forgets the weights.

We do the basic example for general $\mathbb{A}$, and then restrict to the case of Remark 2.2; though we do explain (Remark 2.6) how to extend the result of the Proposition to arbitrary $\mathbb{A}$.

Example 2.3. (Absolute $\mathbb{A}$ -Hodge cohomology of X .) Let

$$('K_{\mathbb{A}}^{\bullet} =) K_{\mathbb{A}}^{\bullet} := C(X, \mathbb{A}(p)) [2p]^{\bullet} = C^{2p+\bullet}(X, \mathbb{A}(p)),$$

$$K_{\mathbb{A} \otimes \mathbb{Q}}^{\bullet} := C^{2p+\bullet}(X, \mathbb{A}(p)) \otimes_{\mathbb{Z}} \mathbb{Q},$$

$$('K_{\mathbb{C}}^{\bullet} =) K_{\mathbb{C}}^{\bullet} := \mathcal{D}(X)(p) [2p]^{\bullet},$$

with $K_{\mathbb{A} \otimes \mathbb{Q}}^{\bullet} \xrightarrow{\beta_1} K_{\mathbb{C}}^{\bullet}$ given by integration: $(2\pi i)^p \gamma \mapsto (2\pi i)^p \delta_{\gamma}$. $F^{\bullet}$ is the Hodge filtration (on $\mathcal{D}_X^{\bullet}$, twisted by p); and $W_r K_{\bullet}^{\bullet} := \begin{cases} K_{\bullet}^{\bullet} & r \geq 0 \\ 0 & r < 0 \end{cases}$ (the stupid weight filtration) yields e.g.

$$\hat{W}_0 K_{\mathbb{A} \otimes \mathbb{Q}}^m = \ker \left\{ W_{-m} K_{\mathbb{A} \otimes \mathbb{Q}}^m \rightarrow \frac{K_{\mathbb{A} \otimes \mathbb{Q}}^{m+1}}{W_{-m-1} K_{\mathbb{A} \otimes \mathbb{Q}}^{m+1}} \right\} = \begin{cases} K_{\mathbb{A} \otimes \mathbb{Q}}^m, & m < 0 \\ \ker(\partial) \subset K_{\mathbb{A} \otimes \mathbb{Q}}^0, & m = 0 \\ 0, & m > 0 \end{cases}.$$

Writing out (2.2) explicitly gives a complex which is

$$C^{2p+\bullet}(X, \mathbb{A}(p)) \oplus C^{2p+\bullet-1}(X, \mathbb{A}(p)) \otimes_{\mathbb{Z}} \mathbb{Q}$$

for $\bullet > 1$,

$$C^{2p+1}(X, \mathbb{A}(p)) \oplus C^{2p}(X, \mathbb{A}(p)) \otimes_{\mathbb{Z}} \mathbb{Q} \oplus \{\ker(d) \subset \mathcal{D}^{2p}(X)\}$$

for $\bullet = 1$,

$$C^{2p}(X, \mathbb{A}(p)) \oplus \{\ker(\partial) \subset C^{2p}(X, \mathbb{A}(p)) \otimes_{\mathbb{Z}} \mathbb{Q}\} \oplus \{\ker(d) \subset F^p \mathcal{D}^{2p}(X)\} \\ \oplus C^{2p-1}(X, \mathbb{A}(p)) \otimes_{\mathbb{Z}} \mathbb{Q} \oplus \mathcal{D}^{2p-1}(X)$$

for $\bullet = 0$, and

$$C^{2p+\bullet}(X, \mathbb{A}(p)) \oplus C^{2p+\bullet}(X, \mathbb{A}(p)) \otimes_{\mathbb{Z}} \mathbb{Q} \oplus F^p \mathcal{D}^{2p+\bullet}(X) \\ \oplus C^{2p+\bullet-1}(X, \mathbb{A}(p)) \otimes_{\mathbb{Z}} \mathbb{Q} \oplus \mathcal{D}^{2p+\bullet-1}(X)$$

for $\bullet < 0$. For $q \in \mathbb{Z}$, its q^{th} cohomology is $H_{\mathcal{H}}^{2p+q}(X, \mathbb{A}(p))$. This maps to $H_{\mathcal{D}}^{2p+q}(X, \mathbb{A}(p))$ by forgetting all $\hat{W}_0$'s in the definition of the *Cone* complex; the map is an isomorphism (only) in degrees $q \leq 0$.

Now for each $\ell \in \mathbb{N}$, let Y^ℓ denote the union of the ℓ -fold intersections of various components of Y , so that $\widetilde{Y}^\ell$ is their disjoint union. Set $Y^0 = X$ and $Y^\ell = \emptyset$ for $\ell < 0$. The proper hypercovering $\widetilde{Y}^\bullet \rightarrow Y$ (together with the inclusion $Y \hookrightarrow X$) leads to double complexes

$$\mathcal{D}^{2p+2i+j}(\widetilde{Y}^{-i})(p+i) =: DR(p)^{i,j}, \quad C^{2p+2i+j}(\widetilde{Y}^{-i}, \mathbb{Q}(p+i)) =: \mathcal{B}(p)^{i,j}$$

with differentials Gy ($i \mapsto i+1$, see formula in §3) and d resp. ∂ ($j \mapsto j+1$). (Here $\mathcal{B}$, DR stand for “Betti”, “de Rham”.) We have associated simple complexes with total differential $\mathbb{D}$, “weight” filtrations ($\mathcal{K} := \mathcal{B}$ or DR)

$$\mathbf{s}^k \mathcal{K}(p) := \oplus_{i(\leq 0)} \mathcal{K}(p)^{i, k-i} \supset \oplus_{\ell \leq i(\leq 0)} \mathcal{K}(p)^{i, k-i} =: ({}'W)^\ell \mathbf{s}^k \mathcal{K}(p),$$

and Hodge filtration

$$F^q \mathbf{s}^k DR(p) := \oplus_i F^q \mathcal{D}^{2p+2i+j}(\widetilde{Y}^{-i})(p+i) \cong \oplus_i F^{q+p+i} \mathcal{D}^{2p+2i+j}(\widetilde{Y}^{-i}).$$

The function of these complexes is to compute cohomology of $X \setminus Y$:

$$H^*(F^q \mathbf{s}^\bullet DR(p)) \cong F^{q+p} H^{*+2p}(X \setminus Y, \mathbb{C}), \quad H^*(\mathbf{s}^\bullet \mathcal{B}(p)) \cong H^{*+2p}(X \setminus Y, \mathbb{Q}(p)).$$

The only nontrivial observation involved in proving this is that

$$(2.3) \quad \frac{\mathcal{D}^\bullet(X)}{\sum_j \iota_*^{Y_j} \mathcal{D}^\bullet(Y_j)[-2]} \rightarrow \frac{\mathcal{D}^\bullet(X)}{\mathcal{D}^\bullet(\text{on } Y)(X)} \cong \mathcal{D}^\bullet(\log Y)(X)$$

is an $F^\bullet$ -filtered quasi-isomorphism [Ja1], so that H^* of $[F^p \text{ of}]$ the l.h.s. computes $[F^p] H^*(X \setminus Y, \mathbb{C})$. Then one looks at the spectral sequence

$\{E_0^{i,j} = DR(p)^{i,j}, d_0 = Gy\}$ computing $H^*(\mathbf{s}^\bullet DR(p))$; its $E_1^{i,j} = 0$ for $i \neq 0$, and $E_1^{0,\bullet}$ is the l.h.s. of (2.3) shifted by $[2p]$.

Now we explain briefly the function of the decalée filtration

$$\begin{aligned} (\text{Dec}('W))^\ell \mathbf{s}^k DR(p) &= \ker \left\{ 'W^{\ell+k} \mathbf{s}^k DR(p) \xrightarrow{\mathbb{D}} \frac{\mathbf{s}^{k+1} DR(p)}{'W^{\ell+k+1} \mathbf{s}^{k+1} DR(p)} \right\} \\ &= \left\{ \ker(d) \subset \mathcal{D}^{2p+2k+\ell}(\widetilde{Y^{-\ell-k}})(p+k+\ell) \right\} \bigoplus \bigoplus_{\ell+k < i(\leq 0)} \mathcal{D}^{2p+i+k}(\widetilde{Y^{-i}})(p+i) \\ &= \mathbf{s}^k \left\{ \tau_{j \leq -\ell} DR(p)^{i,j} \right\}. \end{aligned}$$

The usual weight filtration on cohomology is defined by

$$\begin{aligned} (2.4) \quad W_{2p+k+\ell} H^{2p+k}(X \setminus Y, \mathbb{C}) &:= 'W^{-\ell} H^k(\mathbf{s}^\bullet DR(p)) \\ &:= \text{im}\{H^k('W^{-\ell} \mathbf{s}^\bullet DR(p)) \rightarrow H^k(\mathbf{s}^\bullet DR(p))\}. \end{aligned}$$

To relate $'W$ and $\text{Dec}('W)$, consider their associated spectral sequences (both $\implies H^*(\mathbf{s}^\bullet DR(p))$), with 0^{th} pages

$$\mathcal{E}_0^{i,j} := Gr_W^i \mathbf{s}^{i+j} DR(p) \cong DR(p)^{i,j}, \quad \hat{\mathcal{E}}_0^{i',j'} := Gr_{\text{Dec}('W)}^{i'} \mathbf{s}^{i'+j'} DR(p),$$

and $d_0 = Gr \mathbb{D}$. The first is just the Gysin spectral sequence, which by [De1] degenerates at $\mathcal{E}_2$; hence the second spectral sequence degenerates at $\hat{\mathcal{E}}_1$, since by [De2, sec. 2] $\hat{\mathcal{E}}_r^{-j, i+2j} \xrightarrow{\cong} \mathcal{E}_{r+1}^{i,j} \forall r \geq 1$. So $\mathbb{D}$ on $\mathbf{s}^\bullet DR(p)$ is strictly compatible with $\text{Dec}('W)^\bullet$, and

$$\begin{aligned} (2.5) \quad H^k(\text{Dec}('W)^{-\ell-k} \mathbf{s}^\bullet DR(p)) &\cong \\ \text{im}\{H^k(\text{Dec}('W)^{-\ell-k} \mathbf{s}^\bullet DR(p)) &\rightarrow H^k(\mathbf{s}^\bullet DR(p))\}. \end{aligned}$$

The above compatibility of spectral sequences also implies agreement of the last lines of (2.4) and (2.5). Writing $\hat{W}_0 = \text{Dec}('W)^0$ and specializing to $k = -\ell$, we have shown that

$$(2.6) \quad H^k(\hat{W}_0 \mathbf{s}^\bullet DR(p)) \cong W_{2p} H^{2p+k}(X \setminus Y, \mathbb{C}).$$

This argument can be applied also to $\mathbf{s}^\bullet \mathcal{B}(p)$ and $F^* \mathbf{s}^\bullet DR(p)$.

To define absolute Hodge cohomology, we take our mixed $\mathbb{Q}$ -Hodge complex $K^\bullet$ to be given by

$$K_{\mathbb{Q}}^\bullet := \mathbf{s}^\bullet \mathcal{B}(p), \quad K_{\mathbb{C}}^\bullet := \mathbf{s}^\bullet DR(p),$$

with $W_\bullet (= 'W^{-\bullet})$ and $F^\bullet$ as above, and β_1 given by integration (on each $\widetilde{Y^\ell}$).

Definition 2.4. $H_{\mathcal{H}}^{2p+k}(X \setminus Y, \mathbb{Q}(p)) := H_{\mathcal{H}}^k(K^\bullet)$.

This is computed by the k^{th} cohomology of $(*)$ in Remark 2.2, or
(2.7)

$$\begin{aligned} \text{Cone} \left\{ [(F^0 \mathbf{s}^\bullet DR(p)) \cap \mathbf{s}^\bullet(\tau_{j \leq 0} DR(p))] \xrightarrow{\beta_1 - \beta_2} \mathbf{s}^\bullet(\tau_{j \leq 0} DR(p)) \right\} [-1] \\ = \mathbf{s}^\bullet \mathcal{H}(p)^{\bullet, \bullet}, \end{aligned}$$

where

$$(2.8) \quad \mathcal{H}(p)^{i,j} := \begin{cases} 0 & j > 1 \\ \{\ker(d) \subset \mathcal{D}^{2(p+i)}(\widetilde{Y^{-i}})\} & j = 1 \\ \left(\begin{array}{l} \{\ker(\partial) \subset C^{2(p+i)}(\widetilde{Y^{-i}}, \mathbb{Q}(p+i))\} \\ \oplus \{\ker(d) \subset F^{p+i} \mathcal{D}^{2(p+i)}(\widetilde{Y^{-i}})\} \\ \oplus \mathcal{D}^{2(p+i)-1}(\widetilde{Y^{-i}}) \end{array} \right) & j = 0 \\ \left(\begin{array}{l} C^{2p+2i+j}(\widetilde{Y^{-i}}, \mathbb{Q}(p+i)) \oplus F^{p+i} \mathcal{D}^{2p+2i+j}(\widetilde{Y^{-i}}) \\ \oplus \mathcal{D}^{2p+2i+j-1}(\widetilde{Y^{-i}}) \end{array} \right) & j < 0 \end{cases}$$

has differentials D (vertical $Cone$ differential) and Gy (horizontal). Note that this is 0 for $i > 0$. Elements of $\mathcal{H}(p)^{i,j}$ are always written (a, b, c) , with a, b both zero if $j = 1$.

Proposition 2.5. *The (Gysin) spectral sequence $\mathcal{H}(p)^{\bullet, \bullet}$ associated to $\mathcal{H}(p)^{\bullet, \bullet}$ (with $\mathcal{H}(p)_0^{i,j} := \mathcal{H}(p)^{i,j}$, $d_0 := D$) converges to $H_{\mathcal{H}}^{2p+*}(X \setminus Y, \mathbb{Q}(p))$.*

Remark 2.6. This does not in general degenerate at $\mathcal{H}_2$; see the proof of Prop. 2.7 below. It is easy to modify $\mathcal{H}(p)^{\bullet, \bullet}$ to compute $H_{\mathcal{H}}^{2p+*}(X \setminus Y, \mathbb{A}(p))$ for more general $\mathbb{A}$. Take the 0^{th} column to be the complex from Example 2.3, computing $H_{\mathcal{H}}^{2p+*}(X, \mathbb{A}(p))$, and similarly replace the other columns by complexes computing AH cohomologies of the $\widetilde{Y^{-i}}$.

The presentation (2.7) of $\mathbf{s}^\bullet \mathcal{H}(p)$ as a $Cone$, together with (2.6) (and variants), leads to a short-exact sequence (s.e.s.)

$$(2.9) \quad \begin{aligned} 0 \rightarrow \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), H^{2p-m-1}(X \setminus Y, \mathbb{Q}(p))) &\xrightarrow{\alpha} H_{\mathcal{H}}^{2p-m}(X \setminus Y, \mathbb{Q}(p)) \\ &\xrightarrow{\beta} \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), H^{2p-m}(X \setminus Y, \mathbb{Q}(p))) \rightarrow 0. \end{aligned}$$

There is also a localization long-exact sequence (l.e.s.)

(2.10)

$$\rightarrow H_{\mathcal{H}}^{2p-m}(X, \mathbb{Q}(p)) \rightarrow H_{\mathcal{H}}^{2p-m}(X \setminus Y, \mathbb{Q}(p)) \rightarrow H_{\mathcal{H}, Y}^{2p-m+1}(X, \mathbb{Q}(p)) \rightarrow$$

produced as follows. If we define double complexes

$$\mathcal{H}_Y(p)^{i,j} := \begin{cases} \mathcal{H}(p)^{i,j} & i < 0 \\ 0 & i \geq 0 \end{cases}, \quad \mathcal{H}_X(p)^{i,j} := \begin{cases} \mathcal{H}(p)^{0,j} & i = 0 \\ 0 & i \neq 0 \end{cases},$$

then $\mathbf{s}\mathcal{H}_Y(p)[-1]^{\bullet}$, $\mathbf{s}^{\bullet}\mathcal{H}_X(p) = \mathcal{H}(p)^{0,\bullet}$ have respective k^{th} cohomologies $H_{\mathcal{H}, Y}^{2p+k}(X, \mathbb{Q}(p))$, $H_{\mathcal{H}}^{2p+k}(X, \mathbb{Q}(p))$. The s.e.s. $\mathcal{H}_X(p)^{\bullet,\bullet} \hookrightarrow \mathcal{H}(p)^{\bullet,\bullet} \twoheadrightarrow \mathcal{H}_Y(p)^{\bullet,\bullet}$ of double-complexes translates to a s.e.s. of complexes

$$\mathbf{s}^{\bullet}\mathcal{H}_X(p) \hookrightarrow \mathbf{s}^{\bullet}\mathcal{H}(p) \twoheadrightarrow (\mathbf{s}\mathcal{H}_Y(p)[-1])[+1]^{\bullet},$$

which produces (2.10).

The spectral sequence of Proposition 2.5 induces a “weight” filtration with graded pieces $Gr_j^W H_{\mathcal{H}}^{2p+i+j}(X \setminus Y, \mathbb{Q}(p)) \cong \mathcal{H}(p)_{\infty}^{i,j}$, which will concern us (in a broader context) in §3. For now, we show how to compute these graded pieces. To grease the skids for this endeavor (and for describing the AJ map in §3), define double complexes

$$\mathcal{W}_{\mathbb{Q}}(p)^{i,j} := \tau_{j \leq 0} \mathcal{B}(p)^{i,j} = \begin{cases} 0 & j > 0 \\ \{\ker(\partial) \subset C^{2(p+i)}(\widetilde{Y^{-i}}, \mathbb{Q}(p+i))\} & j = 0 \\ C^{2p+2i+j}(\widetilde{Y^{-i}}, \mathbb{Q}(p+i)) & j < 0 \end{cases},$$

$\mathcal{W}(p)^{i,j} := \tau_{j \leq 0} DR(p)^{i,j}$, with upward shifts $\widetilde{\mathcal{W}}_{[\mathbb{Q}]}(p)^{i,j} := \mathcal{W}_{[\mathbb{Q}]}(p)^{i,j-1}$ (and $\widetilde{\mathcal{B}}(p)^{i,j} := \mathcal{B}(p)^{i,j-1}$). These are each the 0^{th} page of a spectral sequence with d_0 = vertical differential. By [De1], $DR(p)^{\bullet,\bullet}$ and $\mathcal{B}(p)^{\bullet,\bullet}$ degenerate at the 2^{nd} page; so therefore do the $\mathcal{W}$ -spectral sequences, with e.g.

$$\widetilde{\mathcal{W}}_{\mathbb{Q}}(p)_{\infty}^{i,j} = \widetilde{\mathcal{W}}_{\mathbb{Q}}(p)_2^{i,j} \cong Gr_{j-1}^W H^{2p+i+j-1}(X \setminus Y, \mathbb{Q}(p)),$$

and

$$\widetilde{\mathcal{W}}_{\mathbb{Q}}(p)_1^{i,j} \cong H^{2p+2i+j-1}(\widetilde{Y^{-i}}, \mathbb{Q}(p+i)).$$

Also note that besides being a sub-double complex of $DR(p)^{\bullet,\bullet-1}$, $\widetilde{\mathcal{W}}(p)^{\bullet,\bullet}$ injects into $\mathcal{H}(p)^{\bullet,\bullet}$ by sending $\xi \in \mathcal{W}(p)^{i,j-1}$ to $(0, 0, \xi) \in \mathcal{H}(p)^{i,j}$. This produces a map of spectral sequences.

Now write $H^{[k]} := H^{2p+k}(X \setminus Y, \mathbb{Q}(p))$ and consider the s.e.s. of MHS with $j \leq 0$:

$$0 \rightarrow Gr_{j-1}^W H^{[k]} \rightarrow \frac{W_0}{W_{j-2}} H^{[k]} \rightarrow \frac{W_0}{W_{j-1}} H^{[k]} \rightarrow 0.$$

This leads to a l.e.s.:

(2.11)

$$0 \rightarrow \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), \frac{W_0}{W_{j-2}} H^{[k]}) \hookrightarrow \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), \frac{W_0}{W_{j-1}} H^{[k]}) \xrightarrow{\delta_j^{[k]}} \\ \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), Gr_{j-1}^W H^{[k]}) \rightarrow \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), \frac{W_0}{W_{j-2}} H^{[k]}) \longrightarrow \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), \frac{W_0}{W_{j-1}} H^{[k]}) \rightarrow 0,$$

and we set $\text{im}(\delta_j^{[k]}) =: \Xi_j^{[k]}$. Note that $k = i + j - 1 \implies Gr_{j-1}^W H^{[k]} \cong \widetilde{\mathcal{W}}(p)_2^{i,j}$.

Proposition 2.7. *For $j = 0$ we have a s.e.s.*

$$0 \rightarrow \frac{\text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), Gr_{-1}^W H^{[i-1]})}{\Xi_0^{[i-1]}} \rightarrow \mathcal{H}(p)_\infty^{i,0} \rightarrow \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), H^{[i]}) \rightarrow 0.$$

Otherwise,

$$\mathcal{H}(p)_\infty^{i,j} \cong \begin{cases} 0, & j > 1 \text{ (or } i > 0) \\ \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), Gr_0^W H^{[i]}), & j = 1 \\ \frac{\text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), Gr_{j-1}^W H^{[i+j-1]})}{\Xi_j^{[i+j-1]}}, & j < 0 \end{cases}.$$

Proof. We first determine $\mathcal{H}(p)_1^{\bullet,\bullet}$ and $\mathcal{H}(p)_2^{\bullet,\bullet}$. Recall that H^* of $\mathcal{H}(p)^{0,\bullet}$ computes $H_{\mathcal{H}}^{2p+*}(X, \mathbb{Q}(p))$; it follows easily from this that the other columns compute AH cohomology of the $\widetilde{Y}^{-i}$:

$$0 \rightarrow \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), \widetilde{\mathcal{W}}_{\mathbb{Q}}(p)_1^{i,j}) \rightarrow H_{\mathcal{H}}^{2p+2i+j}(\widetilde{Y}^{-i}, \mathbb{Q}(p+i)) \rightarrow \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), \mathcal{W}_{\mathbb{Q}}(p)_1^{i,j}) \rightarrow 0 \\ \parallel \\ \mathcal{H}(p)_1^{i,j}$$

Now $\mathcal{W}_{\mathbb{Q}}(p)_1^{i,j}$ is of pure weight j . Suppose $j \neq 0$, so that the r.h. term vanishes and

$$\mathcal{H}(p)_1^{i,j} \cong \frac{\widetilde{\mathcal{W}}(p)_1^{i,j}}{F^0 \widetilde{\mathcal{W}}(p)_1^{i,j} + \widetilde{\mathcal{W}}_{\mathbb{Q}}(p)_1^{i,j}}.$$

The $\widetilde{\mathcal{W}}(p)_1^{i,j}$ are polarized HS's; by semisimplicity, d_1 -cohomology passes under the Ext and

$$(2.12) \quad \mathcal{H}(p)_2^{i,j} \cong \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), \widetilde{\mathcal{W}}_{\mathbb{Q}}(p)_2^{i,j}) \longleftarrow \widetilde{\mathcal{W}}(p)_2^{i,j}.$$

On the other hand, for $j = 0$, the s.e.s. of complexes

$$0 \rightarrow \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), \widetilde{\mathcal{W}}_{\mathbb{Q}}(p)_1^{\bullet,0}) \rightarrow \mathcal{H}(p)_1^{\bullet,0} \rightarrow \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), \mathcal{W}_{\mathbb{Q}}(p)_1^{\bullet,0}) \rightarrow 0$$

yields a d_1 -cohomology l.e.s.

$$\rightarrow \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), \mathcal{W}_{\mathbb{Q}}(p)_2^{i-1,0}) \xrightarrow{\delta_0^{[i-1]}} \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), \widetilde{\mathcal{W}}_{\mathbb{Q}}(p)_2^{i,0}) \rightarrow \mathcal{H}_2^{i,0}$$

$$\rightarrow \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), \mathcal{W}_{\mathbb{Q}}(p)_2^{i,0}) \xrightarrow{\delta_0^{[i]}} \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), \widetilde{\mathcal{W}}_{\mathbb{Q}}(p)_2^{i+1,0}) \rightarrow$$

where we have again used semisimplicity (to pass d_1 under Ext and Hom). Using (2.11), we get a s.e.s.

(2.13)

$$0 \rightarrow \frac{\text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), \widetilde{\mathcal{W}}_{\mathbb{Q}}(p)_2^{i,0})}{\Xi_0^{[i-1]}} \xrightarrow{\alpha} \mathcal{H}_2^{i,0} \xrightarrow{\beta} \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), \frac{W_0}{W_{-2}} H^{[i]}) \rightarrow 0.$$

Now the higher differentials d_ℓ ($\ell \geq 2$) are all 0 on $\widetilde{\mathcal{W}}(p)_{\bullet}^{\bullet}$, which means that the surjectivity (2.12) extends to the entire map of spectral sequences $\widetilde{\mathcal{W}}(p)_r^{\bullet} \rightarrow \mathcal{H}(p)_r^{\bullet}$ ($r \geq 2$) off the 0^{th} row. Hence the only nontrivial higher differentials on $\mathcal{H}(p)_{\bullet}^{\bullet}$ are the

$$d_{-j+1} : \mathcal{H}_{-j+1}^{i+j-1,0} \rightarrow \mathcal{H}_{-j+1}^{i,j}$$

for $j < 0$. (For $j = 1$, the proof is now finished.) Moreover, since $\widetilde{\mathcal{W}}(p)_2^{i,0} \rightarrow \mathcal{H}(p)_2^{i,0}$ factors through, and surjects onto, the l.h. term of (2.13), we have $d_2|_{\text{im}(\alpha)} = 0$ and so d_2 factors through β :

$$\begin{array}{ccc} \mathcal{H}_2^{i,0} & \xrightarrow{d_2} \mathcal{H}_2^{i+2,-1} \cong \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), \text{Gr}_{-2}^W H^{[i]}) & \\ & \searrow \beta & \uparrow \delta_{-1}^{[i]} \\ & \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), \frac{W_0}{W_{-2}} H^{[i]}) & \end{array}$$

Using again (2.11) we get a s.e.s.

$$\begin{array}{ccc} 0 \rightarrow \frac{\text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), \widetilde{\mathcal{W}}_{\mathbb{Q}}(p)_3^{i,0})}{\Xi_0^{[i-1]}} \xrightarrow{\alpha} \mathcal{H}_3^{i,0} \xrightarrow{\beta_3} \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), \frac{W_0}{W_{-3}} H^{[i]}) \rightarrow 0, & & \\ & \parallel & \parallel \\ & \ker(d_2) \longrightarrow \ker(\delta_{-1}^{[i]}) & \end{array}$$

and iterating the argument completes the proof for $j = 0$. (At ∞ , we have r.h. term $\text{Hom}_{\text{MHS}}(\mathbb{Q}(0), W_0 H^{[i]}) = \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), H^{[i]})$.) This argument also shows that each d_{-j+1} factors

$$\begin{array}{ccc} \mathcal{H}_{-j+1}^{i+j-1,0} & \xrightarrow{d_{-j+1}} \mathcal{H}_{-j+1}^{i,j} & \\ & \searrow \beta_{-j+1} & \uparrow \delta_j^{[i+j-1]} \\ & \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), \frac{W_0}{W_{j-1}} H^{[i+j-1]}) & \end{array}$$

so that $\text{im}(d_{-j+1}) = \text{im}(\delta_j^{[i+j-1]}) = \Xi_j^{[i+j-1]}$ and we are done. $\square$

3. THE WEIGHT-FILTERED ABEL-JACOBI MAP

Having dispensed with the preliminaries of §2, we are ready to explicitly describe the absolute Hodge cycle-class map

$$c_{\mathcal{H}}^{p,m} : CH^p(U, m) \rightarrow H_{\mathcal{H}}^{2p-m}(U, \mathbb{Q}(p))$$

for smooth quasi-projective U . We do this first in terms of a map of double-complexes (§3.1), then in §3.2 via generalized membrane integrals. The double-complex construction is then applied in a discussion of the Beilinson-Hodge conjecture (in §3.3).

3.1. Gysin spectral sequences and higher residue maps. We start by laying out a number of complexes: given a smooth projective algebraic $W/k \subseteq \mathbb{C}$, set

$$C_{\mathcal{M}}^{\bullet}(W, \mathbb{Q}(q)) := Z_{\mathbb{R}}^q(W, 2q - \bullet).$$

Also writing W for $W_{\mathbb{C}}^{\text{an}}$, define

$$C_{\mathcal{B}}^{\bullet}(W, \mathbb{Q}(q)) := C_{\text{top}}^{\bullet}(W, \mathbb{Q}(q)), \quad C_{[F]DR}^{\bullet}(W, \mathbb{C}) := [F^q]\mathcal{D}^{\bullet}(W),$$

$$\begin{aligned} C_{\mathcal{D}}^{\bullet}(W, \mathbb{Q}(q)) &:= \text{Cone} \{ C_{\text{top}}^{\bullet}(W, \mathbb{Q}(q)) \oplus F^q \mathcal{D}^{\bullet}(W) \rightarrow \mathcal{D}^{\bullet}(W) \} [-1]^{\bullet} \\ &= C_{\text{top}}^{\bullet}(W, \mathbb{Q}(q)) \oplus F^q \mathcal{D}^{\bullet}(W) \oplus \mathcal{D}^{\bullet-1}(W), \end{aligned}$$

$$C_{\mathcal{H}}^{\bullet}(W, \mathbb{Q}(q)) := \begin{cases} 0, & \bullet > 2q + 1 \\ \mathcal{D}_{d\text{-cl}}^{2q}(W), & \bullet = 2q + 1 \\ Z_{\text{top}}^{2q}(W, \mathbb{Q}(q)) \oplus F^q \mathcal{D}_{d\text{-cl}}^{2q}(W) \oplus \mathcal{D}^{2q-1}(W), & \bullet = 2q \\ C_{\mathcal{D}}^{\bullet}(W, \mathbb{Q}(q)), & \bullet < 2q \end{cases}$$

and let $\mathcal{K}$ stand for $\mathcal{B}$, DR , FDR , $\mathcal{D}$, $\mathcal{H}$, or $\mathcal{M}$. (The respective differentials ∂_{top} , d , d , D , D , and $\partial_{\mathcal{B}}$ are then uniformly denoted $d_{\mathcal{K}}$.) We have $H^*(C_{[F]DR}^{\bullet}(W, \mathbb{C})) = [F^q]H_{DR}^*(W, \mathbb{C})$, and for all other $\mathcal{K}$

$$H^*(C_{\mathcal{K}}^{\bullet}(W, \mathbb{Q}(q))) \cong H_{\mathcal{K}}^*(W, \mathbb{Q}(q)).$$

Now let $\mathcal{Y} = \bigcup \mathcal{Y}_i \subset \mathcal{X}$ be a NCD (in $\mathcal{X}$ smooth projective/ k), $\mathcal{Y}_I := \bigcap_{i \in I} \mathcal{Y}_i$, and $\mathcal{Y}^{\ell} := \bigcup_{|I|=\ell} \mathcal{Y}_I$, so that $\widetilde{\mathcal{Y}^{\ell}} \cong \Pi_{|I|=\ell} \mathcal{Y}_I$. (By convention, $\mathcal{Y}^0 := \mathcal{X}$.) In what follows, we exclude $\mathcal{K} = [F]DR$ for uniformity of notation, though similar results hold. (In particular, the spectral sequences $DR(p)$, $FDR(p)$ would degenerate at the 2nd page.)

Define (mostly 3rd quadrant) double complexes

$$\mathcal{K}_{\mathcal{X} \setminus \mathcal{Y}}(p)_0^{a,b} := C_{\mathcal{K}}^{2p+2a+b}(\widetilde{\mathcal{Y}^{-a}}, \mathbb{Q}(p+a))$$

with vertical differential $d_{\mathcal{K}}$, horizontal differential²

$$Gy := 2\pi i \sum_{|I|=-a} \sum_{i \in I} (-1)^{\langle i \rangle_{I \setminus \{i\}}} (\iota_{\mathcal{Y}_I \subset \mathcal{Y}_{I \setminus \{i\}}})_*$$

($\langle i \rangle_J$:= position in which i occurs in J), and total differential $\mathbb{D} := d_{\mathcal{K}} + (-1)^b Gy$ (on the associated simple complex $s^\bullet \mathcal{K}(p)$).

Example 3.1. $\mathcal{M}(p)_0^{\bullet, \bullet}$ sits in the 3rd quadrant, with upper-right-hand 3×3

$$\begin{array}{ccccccc} & \longrightarrow & Z^{p-2}(\widetilde{\mathcal{Y}}^2) & \xrightarrow{Gy} & Z^{p-1}(\widetilde{\mathcal{Y}}^1) & \xrightarrow{Gy} & Z^p(\mathcal{X}) \\ & & \uparrow d & & \uparrow d & & \uparrow d \\ & \longrightarrow & Z^{p-2}(\widetilde{\mathcal{Y}}^2, 1) & \xrightarrow{Gy} & Z^{p-1}(\widetilde{\mathcal{Y}}^1, 1) & \xrightarrow{Gy} & Z^p(\mathcal{X}, 1) \\ & & \uparrow d & & \uparrow d & & \uparrow d \\ & \longrightarrow & Z^{p-2}(\widetilde{\mathcal{Y}}^2, 2) & \xrightarrow{Gy} & Z^{p-1}(\widetilde{\mathcal{Y}}^1, 2) & \xrightarrow{Gy} & Z^p(\mathcal{X}, 2) . \\ & & \uparrow & & \uparrow & & \uparrow \end{array}$$

Taking $d_0 := d_{\mathcal{K}}$ gives Gysin spectral sequences $\mathcal{K}(p)_r^{a,b} \xRightarrow{(*=a+b)}$

$H^*(s^\bullet \mathcal{K}(p)) \cong H_{\mathcal{K}}^{2p+*}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p))$. (Omitting the 0th column would compute $H_{\mathcal{K}, \mathcal{Y}}^{2p+*+1}(\mathcal{X}, \mathbb{Q}(p))$.) One finds in each case $\mathcal{K}(p)_1^{a,b} \cong H_{\mathcal{K}}^{2p+2a+b}(\widetilde{\mathcal{Y}}^{-a}, \mathbb{Q}(p+a))$; and for $\mathcal{K} = \mathcal{B}$ we have degeneration at $\mathcal{B}(p)_{2(=\infty)}$ with $\mathcal{B}(p)_\infty^{a,b} = Gr_b^W H_{(\mathcal{B})}^{2p+a+b}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p))$, where $W_\bullet$ is the weight filtration of Deligne [De1]. To generalize the weight filtration to arbitrary $\mathcal{K}$, set $\tilde{W}_j s^\bullet \mathcal{K}(p) := s^\bullet \mathcal{K}(p)^{\bullet \geq -j, \bullet}$ and define

$$W_{*+j} H_{\mathcal{K}}^{2p+*}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)) := \text{im} \left\{ H^*(\tilde{W}_j s^\bullet \mathcal{K}(p)) \rightarrow H^*(s^\bullet \mathcal{K}(p)) \right\},$$

so that $Gr_b^W H_{\mathcal{K}}^{2p+a+b}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)) \cong \mathcal{K}_{\mathcal{X} \setminus \mathcal{Y}}(p)_\infty^{a,b}$; in particular,

$$\mathcal{K}_{\mathcal{X} \setminus \mathcal{Y}}(p)_\infty^{0,b} \cong \underline{H_{\mathcal{K}}^{2p+b}}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)) := \text{im} \left\{ H_{\mathcal{K}}^{2p+b}(\mathcal{X}, \mathbb{Q}(p)) \rightarrow H_{\mathcal{K}}^{2p+b}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)) \right\}$$

(lowest weight). While the Gysin spectral sequences for $\mathcal{K} = \mathcal{M}, \mathcal{H}, \mathcal{D}$ do not degenerate at $\mathcal{K}(p)_2$, for $\mathcal{H}, \mathcal{D}$ one can show all $d_{r \geq 2} : \mathcal{K}(p)_r^{a,b < 0} \rightarrow \mathcal{K}(p)_r^{a+r, b-r+1}$ are zero; precise formulas for the $\mathcal{H}(p)_\infty^{a,b}$ were worked out in Prop. 2.7.

More generally and perhaps more usefully, one can map all $\mathcal{K}_\infty$ terms to easily describable targets. First, to compute $H_{\mathcal{K}}$ of $\mathcal{Y}^k \setminus \mathcal{Y}^{k+1} \cong$

²ignore the $2\pi i$ (in Gy) for $\mathcal{K} = \mathcal{M}$.

$\widetilde{\mathcal{Y}}^k \setminus (\mathcal{Y}^{k+1} \times_{\mathcal{Y}^k} \widetilde{\mathcal{Y}}^k)$ (with good compactification $\widetilde{\mathcal{Y}}^k$), we can use

$$(3.1) \quad \mathcal{K}_{\mathcal{Y}^k \setminus \mathcal{Y}^{k+1}}(q)^{\alpha, \beta} = C_{\mathcal{K}}^{2q+2\alpha+\beta} \left(\widetilde{\mathcal{Y}^{k-\alpha} \times_{\mathcal{Y}^k} \mathcal{Y}^k}, \mathbb{Q}(q+\alpha) \right).$$

Now while the quotient double complex $\mathcal{K}^{\leq -k}(p)^{a,b} := \begin{cases} 0 & a > -k \\ \mathcal{K}(p)^{a,b} & a \leq -k \end{cases}$ (of $\mathcal{K}(p)_0^{a,b}$) does not compute $H_{\mathcal{K}, \mathcal{Y}^k}^{2p+*}(\mathcal{X}, \mathbb{Q}(p))$ for $k > 1$, it does map to (3.1). More precisely, there are maps of double complexes

$$\mathcal{K}_{\mathcal{X} \setminus \mathcal{Y}}(p)_0^{a,b} \rightarrow \mathcal{K}_{\mathcal{X} \setminus \mathcal{Y}}^{\leq -k}(p)_0^{a,b} \rightarrow \mathcal{K}_{\mathcal{Y}^k \setminus \mathcal{Y}^{k+1}}(p-k)_0^{a+k,b}$$

inducing maps of spectral sequences, hence of ∞ terms

$$\mathcal{K}_{\mathcal{X} \setminus \mathcal{Y}}(p)_{\infty}^{-k,b} \rightarrow \mathcal{K}_{\mathcal{Y}^k \setminus \mathcal{Y}^{k+1}}(p-k)_{\infty}^{0,b} \cong \underline{H_{\mathcal{K}}^{2p-2k+b}(\mathcal{Y}^k \setminus \mathcal{Y}^{k+1}, \mathbb{Q}(p-k))}.$$

This defines *higher residue maps*

$$Res^{-a} : Gr_b^W H_{\mathcal{K}}^{2p+a+b}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)) \rightarrow \underline{H_{\mathcal{K}}^{2p+2a+b}(\mathcal{Y}^{-a} \setminus \mathcal{Y}^{-a+1}, \mathbb{Q}(p+a))},$$

which are in general neither injective nor surjective. More concretely, if we let $U^{-a} :=$ connected component of $\mathcal{Y}^{-a} \setminus \mathcal{Y}^{-a+1}$, the Res^{-a} into $\underline{H_{\mathcal{K}}^{2p+2a+b}(U^{-a}, \mathbb{Q}(p+a))}$ is computed by taking $(-a)$ successive residues along a “flag” of connected components $U^i \subseteq \mathcal{Y}^i \setminus \mathcal{Y}^{i+1}$ with $\overline{U^{-a}} \subsetneq \overline{U^{-a-1}} \subsetneq \dots \subsetneq \overline{U^1} \subsetneq \overline{U^0} = \mathcal{X}$.

Consider the “realization” maps (of double complexes)

$$\mathcal{M}(p)_0^{a,b} \rightarrow \mathcal{H}(p)_0^{a,b} \rightarrow \mathcal{D}(p)_0^{a,b} \rightarrow \mathcal{B}(p)_0^{a,b},$$

with the first given by sending $\mathfrak{Z} \in Z_{\mathbb{R}}^{p+a}(\widetilde{\mathcal{Y}^{-a}}, -b)$ to $(2\pi i)^{p+a+b}((2\pi i)^{-b}T_3, \Omega_3, R_3)$, and the latter two obvious. (Here $T_3 \in C_{top}^{2p+2a+b}(\widetilde{\mathcal{Y}^{-a}}, \mathbb{Q}(p+a))$, $\Omega_3 \in F^{p+a}\mathcal{D}^{2p+2a+b}(\widetilde{\mathcal{Y}^{-a}})$, $R_3 \in \mathcal{D}^{2p+2a+b-1}(\widetilde{\mathcal{Y}^{-a}})$ are as in [KLM, sec. 5.3-4].) By considering the induced maps on the cohomologies of the associated simple complexes, we have *$W_{\bullet}$ -filtered cycle-class maps*

$$c_{\mathcal{K}}^{p,m} : CH^p(\mathcal{X} \setminus \mathcal{Y}, m) \cong H_{\mathcal{M}}^{2p-m}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)) \rightarrow H_{\mathcal{K}}^{2p-m}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)),$$

with graded pieces

$$Gr_b^W c_{\mathcal{K}}^{p,m} : \mathcal{M}(p)_{\infty}^{-b-m,b} \rightarrow \mathcal{K}(p)_{\infty}^{-b-m,b}.$$

Remark 3.2. If we omit the 0th column from the entire construction, we get cycle-class maps

$$CH^{p-1}(\mathcal{Y}, m-1) \cong H_{\mathcal{M}, \mathcal{Y}}^{2p-m+1}(\mathcal{X}, \mathbb{Q}(p)) \rightarrow H_{\mathcal{K}, \mathcal{Y}}^{2p-m+1}(\mathcal{X}, \mathbb{Q}(p))$$

for cohomology with support on $\mathcal{Y}$.

Since $\mathcal{M}(p)_0^{a,b} = 0$ for $b > 0$, $H_{\mathcal{M}}^{2p+*}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)) = W_0 H_{\mathcal{M}}^{2p+*}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p))$. Therefore one consequence of the above is the existence of a commutative diagram ($\mathcal{K} = \mathcal{H}, \mathcal{D}$, or $\mathcal{B}$)

$$\begin{array}{ccc} CH^p(\mathcal{X} \setminus \mathcal{Y}, m) & \xrightarrow{c_{\mathcal{K}}} & W_0 H_{\mathcal{K}}^{2p-m}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)) \\ \downarrow \text{Res}^m & & \downarrow \text{Res}^m \\ CH^{p-m}(\mathcal{Y}^m \setminus \mathcal{Y}^{m+1}) & \xrightarrow{c_{\mathcal{K}}} & \underline{H_{\mathcal{K}}^{2p-2m}(\mathcal{Y}^m \setminus \mathcal{Y}^{m+1}, \mathbb{Q}(p-m))}. \end{array}$$

(Note that $CH^{p-m}(\mathcal{Y}^m \setminus \mathcal{Y}^{m+1}) = \underline{CH^{p-m}(\mathcal{Y}^m \setminus \mathcal{Y}^{m+1})}$ since we are in the case of algebraic cycles.)

Example 3.3. We give a simple application of the corresponding diagram for $\mathcal{K} = DR$, which has upper-r.h. composition

$$CH^p(\mathcal{X} \setminus \mathcal{Y}, m) \xrightarrow{c_{FDR}} F^p W_{2p} H_{DR}^{2p-m}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{C}) \xrightarrow{\text{Res}^m} F^{p-m} \underline{H^{2p-2m}(\mathcal{Y}^m \setminus \mathcal{Y}^{m+1}, \mathbb{C})},$$

in the case $p = m := n$.

The construction of [DK] produces families of hypersurfaces

$$\begin{array}{ccccc} X_t & \hookrightarrow & \mathcal{X} & \hookrightarrow & \mathbb{P}_{\tilde{\Delta}} \times \mathbb{P}^1 \\ \downarrow & & \downarrow \pi & \swarrow & \\ \{t\} & \hookrightarrow & \mathbb{P}^1 & & \end{array}$$

in a Toric Fano n -fold $\mathbb{P}_{\tilde{\Delta}}$ ($n = 2, 3, 4$) with the generic X_t nonsingular, and X_0 a NCD satisfying $\mathbb{P}_{\tilde{\Delta}} \setminus X_0 \cong (\mathbb{C}^*)^n$ (with coordinates $x_1, \dots, x_n$). We set $\mathcal{Y} := X_0 \subset \mathcal{X}$. The hypotheses of [DK] Thm. 1 guarantee the existence of $\Xi \in CH^n(\mathcal{X} \setminus \mathcal{Y}, m)$ restricting to $\langle \{x_1, \dots, x_n\} \rangle \in CH^n(\mathcal{X} \cap (\mathbb{C}^*)^n, n)$. From this we deduce $\text{dlog} x_1 \wedge \dots \wedge \text{dlog} x_n (= \Omega_{\Xi})$ is a holomorphic form³ on $\mathcal{X} \setminus \mathcal{Y}$ representing $c_{DR}(\Xi)$, or equivalently the infinitesimal invariant of $\{c_{\mathcal{H}}(\Xi|_{X_t})\} \in \Gamma(U \subset \mathbb{P}^1, R^{n-1} \pi_* \mathbb{C}/\mathbb{Q}(n))$ as

$$F^n H^n(\mathcal{X} \setminus \mathcal{Y}, \mathbb{C}) \hookrightarrow \Gamma(U, \Omega_U^1 \otimes F^{n-1} R^{n-1} \pi_* \mathbb{C}).$$

Since $\text{Res}^n(\bigwedge^n \text{dlog} x_i) \in H^0(\mathcal{Y}^n = \text{Ipts.}, \mathbb{C})$ is obviously nonzero, we get generic nontriviality of $\langle \Xi|_{X_t} \rangle \in CH^n(X_t, n)$.

3.2. Generalization of the Griffiths prescription. The composition of $c_{\mathcal{H}}$ with β from (2.9) yields the fundamental class map

$$cl : CH^p(\mathcal{X} \setminus \mathcal{Y}, m) \rightarrow \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), H^{2p-m}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)));$$

³The assertion is that the pullback of $\bigwedge_{i=1}^n \text{dlog} x_i$ from $(\mathbb{C}^*)^n$ to $\mathcal{X} \cap \{(\mathbb{C}^*)^n \times \mathbb{P}^1\}$ extends to a holomorphic form on $\mathcal{X} \setminus X_0 \times \{0\}$. In fact $\mathcal{X} \setminus \{\mathcal{X} \cap ((\mathbb{C}^*)^n \times \mathbb{P}^1)\} = (D \times \mathbb{P}^1) \cup (\mathcal{Y} = X_0 \times \{0\})$, but Ω_{Ξ} can have no residues along $D \times \mathbb{P}^1$.

we write $cl(\xi) =: [\xi]$. The term “Abel-Jacobi map” refers either to $c_{\mathcal{H}}$ or its restriction

$$AJ : CH_{hom}^p(\mathcal{X} \setminus \mathcal{Y}, m) \rightarrow Ext_{MHS}^1(\mathbb{Q}(0), H^{2p-m-1}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p))) =: J^{p,m}(\mathcal{X} \setminus \mathcal{Y})$$

to $\ker(cl)$. An interesting consequence of Prop. 2.7 is that cl factors through $\mathcal{H}(p)_{\infty}^{-m,0}$ (via $Gr_0^W c_{\mathcal{H}}$), so that its kernel $CH_{hom}^p(\mathcal{X} \setminus \mathcal{Y}, m) \supseteq W_{-1}CH^p(\mathcal{X} \setminus \mathcal{Y}, m)$.

We shall be interested in these maps when $\mathcal{X} \xrightarrow{\pi} \mathcal{S}$ is the k -spread of a projective variety X (defined over a field extension of k), and $\mathcal{Y} = \pi^{-1}(D)$ for D a divisor in $\mathcal{S}$. Relevant for the application is an extension of the arithmetic Bloch-Beilinson conjecture to higher Chow groups of smooth projective varieties.

Conjecture 3.4. [BBC] *For all smooth projective $\mathcal{X}/\bar{\mathbb{Q}}$, $\mathcal{Y} = \emptyset$, the $c_{\mathcal{H}}^{p,m}$ are injective ($\forall p, m$).*

For *nonempty* $\mathcal{Y}$ (NCD/ $\bar{\mathbb{Q}}$), BBC together with the Hodge conjecture implies injectivity of the $c_{\mathcal{H}}^{p,m}$ (“quasi-projective BBC”), and even of the $Gr_b^W c_{\mathcal{H}}^{p,m}$. While injectivity generally fails for $(\mathcal{X}, \mathcal{Y})$ defined over $k \subseteq \mathbb{C}$ of higher transcendence degree, one might ask whether $c_{\mathcal{H}}^{p,m}$ is still strictly compatible with $W_{\bullet}$; this seems to be a difficult problem.

Now we describe an approach to AJ to be used in §4. First, define $'C_{\mathcal{H}}^{\bullet}(W, \mathbb{Q}(q)) \subset C_{\mathcal{H}}^{\bullet}(W, \mathbb{Q}(q))$ (W as above) to be the subcomplex consisting of triples of the form $(0, *, *)$ (no C_{top} ’s). It maps quasi-isomorphically to

$$C_{\mathcal{I}}^{\bullet}(W, \mathbb{Q}(q)) := \begin{cases} 0, & \bullet > 0 \\ \mathcal{D}^{2q}(W)/F^q \mathcal{D}^{2q}(W), & \bullet = 0 \\ \widetilde{\mathcal{D}}^{\bullet}(W)/F^q \widetilde{\mathcal{D}}^{\bullet}(W), & \bullet < 0 \end{cases}.$$

Writing $\widetilde{\mathcal{I}}_{\mathcal{X} \setminus \mathcal{Y}}(p)_0^{a,b} := C_{\mathcal{I}}^{2p+2a+b-1}(\widetilde{Y^{-a}}, \mathbb{Q}(p+a))$, we have

$$H^*(s^{\bullet} \widetilde{\mathcal{I}}(p)) \cong \frac{W_{2p} H_{DR}^{2p+*-1}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{C})}{F^p W_{2p} H_{DR}^{2p+*-1}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{C})}$$

and

$$(3.2) \quad \frac{H^{-m}(s^{\bullet} \widetilde{\mathcal{I}}(p))}{H^{-m}(sW_{\mathbb{Q}}(p)[-1]^{\bullet})} \cong J^{p,m}(\mathcal{X} \setminus \mathcal{Y}).$$

A class $\xi \in CH_{hom}^p(\mathcal{X} \setminus \mathcal{Y}, m)$ has a $\mathbb{D}$ -closed representative $\mathfrak{Z}^{\bullet} \in s^{-m}\mathcal{M}(p)$; this is sent to⁴ $(2\pi i)^{p-m}((2\pi i)^{m+\bullet} T_{\mathfrak{Z}^{\bullet}}, \Omega_{\mathfrak{Z}^{\bullet}}, R_{\mathfrak{Z}^{\bullet}}) \in s^{-m}\mathcal{H}(p)$ by [KLM], where $(2\pi i)^{p+\bullet} T_{\mathfrak{Z}^{\bullet}} = \mathbb{D}\{(2\pi i)^{p+\bullet} \Gamma^{\bullet}\}$ for $\{(2\pi i)^{p+\bullet} \Gamma^{\bullet}\} \in s^{-m-1}\mathcal{W}_{\mathbb{Q}}(p)$. The triple is changed by $\mathbb{D}$ -coboundary to

⁴here $\bullet = i$ indexes entries along the $(-m)^{\text{th}}$ diagonal.

$(2\pi i)^{p-m}(0, \Omega_{\mathcal{Z}\bullet}, R_{\mathcal{Z}\bullet} + (2\pi i)^{m+\bullet}\delta_{\Gamma\bullet})$, then “projected” to $\mathbb{D}$ -closed $(2\pi i)^{p-m}\{\overline{R_{\mathcal{Z}\bullet} + (2\pi i)^{m+\bullet}\delta_{\Gamma\bullet}}\} \in \mathbf{s}^{-m}\tilde{\mathcal{I}}(p)$, which yields the class $AJ(\xi)$ in (3.2) directly. This is used in §4.

Given an algebraic cycle $\mathcal{Z} \stackrel{\text{hom}}{\equiv} 0$ on a smooth projective variety X , the classical *Griffiths prescription* ([Gr], or see [Le3, 12.12]) computes $AJ(\mathcal{Z})$ by integrating suitable test forms over a topological membrane bounding on $\mathcal{Z}$. With the last description of $AJ(\xi)$, it is easy to generalize this. Define a new double complex $\Omega(p)_{i,j} := \Omega^{2n-2p-j+1}_{(\mathcal{Y}^{-i})^\infty}(\widetilde{\mathcal{Y}^{-i}})$ ($n = \dim \mathcal{X}$) with differentials $\mathfrak{I} : \Omega(p)_{i,j} \rightarrow \Omega(p)_{i-1,j}$ (dual to Gy , see §8.1), and $d : \Omega(p)_{i,j} \rightarrow \Omega(p)_{i,j-1}$. Letting currents act on C^∞ forms (and multiplying the result by $(2\pi\sqrt{-1})^{-i}$) induces pairings

$$\int : F^{n-p+1}\Omega(p)_{i,j} \otimes \tilde{\mathcal{I}}(p)^{i,j} \rightarrow \mathbb{C},$$

well-defined since $\dim(\widetilde{\mathcal{Y}^{-i}}) = n + i$. These extend by *summing along the diagonal* to

$$(3.3) \quad \not\int : \mathbf{s}_{-m}F^{n-p+1}\Omega(p)_{\bullet,\bullet} \otimes \mathbf{s}^{-m}\tilde{\mathcal{I}}(p)^{\bullet,\bullet} \longrightarrow \mathbb{C},$$

then by taking cohomology to the perfect pairing

$$F^{n-p+1}H^{2n-2p+m+1}((\mathcal{X}, \mathcal{Y}), \mathbb{C}) \otimes \frac{H^{2p-m-1}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{C})}{F^p} \longrightarrow \mathbb{C}.$$

By forgetting weights, $J^{p,m}(\mathcal{X} \setminus \mathcal{Y})$ maps to the Deligne Jacobian

$$J_D^{p,m}(\mathcal{X} \setminus \mathcal{Y}) := \frac{H^{2p-m-1}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{C})}{F^p H^{2p-m-1}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{C}) + H^{2p-m-1}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p))} \cong \frac{\{H_{-m}(\mathbf{s}_\bullet F^{n-p+1}\Omega(p))\}^\vee}{\text{im} H^{-m}(\mathbf{s}^\bullet \tilde{\mathcal{B}}(p))},$$

and the image of $AJ(\xi)$ is just the functional on “test forms” $\omega_\bullet \in \mathbf{s}_{-m}F^{n-p+1}\Omega(p)$ given by $\not\int$ against the “generalized membrane” $(2\pi i)^{p-m}\{R_{\mathcal{Z}\bullet} + (2\pi i)^{m+\bullet}\delta_{\Gamma\bullet}\}$.

Example 3.5. The case of algebraic cycles ($m = 0$) on $\mathcal{X} \setminus \mathcal{Y}$, where $\mathcal{Y} \xhookrightarrow{\iota} \mathcal{X}$ is a smooth divisor, is already entertaining. Fix $n (= \dim \mathcal{X}) = 5$, and let $\langle \mathcal{Z} \rangle \in CH_{\text{hom}}^3(\mathcal{X} \setminus \mathcal{Y})$. Clearly $\overline{\mathcal{Z}} \in Z_{(\mathbb{R})}^3(\mathcal{X})$ gives an element of $\mathbf{s}^0\mathcal{M}_{\mathcal{X} \setminus \mathcal{Y}}(3)$ representing $\langle \mathcal{Z} \rangle$. Since $[\mathcal{Z}] = 0$ in $\text{Hom}_{\text{MHS}}(\mathbb{Q}(0), H^6(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(3)))$, there exist

$$\Gamma \in C_{\text{top}}^5(\mathcal{X}, \mathbb{Q}(3)), \gamma \in C_{\text{top}}^4(\mathcal{Y}, \mathbb{Q}(2)); \Xi \in F^3\mathcal{D}^5(\mathcal{X}), \xi \in F^2\mathcal{D}^4(\mathcal{Y})$$

s.t. $\partial\Gamma + (2\pi i)\iota_*\gamma = (2\pi i)^3\overline{\mathcal{Z}}$, $d\Xi + (2\pi i)\iota_*\xi = (2\pi i)^3\delta_{\overline{\mathcal{Z}}}$. (In fact, Γ, Ξ determine γ, ξ .) This yields an element $\{(\gamma, \xi, 0), (\Gamma, \Xi, 0)\} \in \mathbf{s}^{-1}\mathcal{H}_{\mathcal{X} \setminus \mathcal{Y}}(3)$, while the double-complex AJ applied to $\overline{\mathcal{Z}}$ gives $(2\pi i)^3(\overline{\mathcal{Z}}, \delta_{\overline{\mathcal{Z}}}, 0) \in \mathbf{s}^0\mathcal{H}_{\mathcal{X} \setminus \mathcal{Y}}(3)$. Adding $\mathbb{D}$ of the former to the latter gives

$\{(0, 0, \delta_\gamma - \xi), (0, 0, \delta_\Gamma - \Xi)\}$, so we have $\{\delta_\gamma - \xi, \delta_\Gamma - \Xi\} \mapsto \overline{\{\delta_\gamma, \delta_\Gamma\}} \in \mathbf{s}^0 \widetilde{\mathcal{I}}_{\mathcal{X} \setminus \mathcal{Y}}(3)$.

Now set

$$\begin{aligned} \mathbb{V} &:= \frac{\{(\omega, \eta) \mid \omega \in F^3 \Omega_{\mathcal{X}^\infty}^5(\mathcal{X})_{d\text{-cl}}, \eta \in F^3 \Omega_{\mathcal{Y}^\infty}^4(\mathcal{Y}), \iota^* \omega = d\eta\}}{\{(-d\alpha, \iota^* \alpha - d\beta) \mid \alpha \in F^3 \Omega_{\mathcal{X}^\infty}^4(\mathcal{X}), \beta \in F^3 \Omega_{\mathcal{Y}^\infty}^3(\mathcal{Y})\}} \\ &\cong H_0(\mathbf{s}_\bullet F^3 \Omega(3)) \cong F^3 H^5((\mathcal{X}, \mathcal{Y}), \mathbb{C}), \\ \mathbb{W}_{\mathbb{Q}} &:= \frac{\{(\tau, \sigma) \mid \tau \in C_{top}^5(\mathcal{X}, \mathbb{Q}(3)), \sigma \in Z_{top}^4(\mathcal{Y}, \mathbb{Q}(2)), (2\pi i) \iota_* \beta = \partial \tau\}}{\{(-\partial \varepsilon + 2\pi i \iota_* \varphi, -\partial \varphi) \mid \varepsilon \in C_{top}^4(\mathcal{X}, \mathbb{Q}(3)), \varphi \in C_{top}^3(\mathcal{Y}, \mathbb{Q}(2))\}} \\ &\cong H^0(\mathbf{s}_\bullet \widetilde{\mathcal{B}}(3)) \cong H_5((\mathcal{X}, \mathcal{Y}), \mathbb{Q}(3)). \end{aligned}$$

Using (3.3), $\overline{\{\delta_\gamma, \delta_\Gamma\}}$ gives a functional on $\mathbb{V}$ by

$$(\omega, \eta) \longmapsto \int_{\Gamma} \omega + (2\pi i) \int_{\gamma} \eta.$$

(Note that if γ is algebraic, then $\int_{\gamma} \eta = 0$; but to arrange this in general requires the Hodge conjecture.) This functional yields an element of $J_{\mathcal{D}}^{3,0}(\mathcal{X} \setminus \mathcal{Y}) \cong \frac{\mathbb{V}}{\text{im } \mathbb{W}_{\mathbb{Q}}}$ which depends only on $\langle \mathcal{Z} \rangle$ and not on the choices of Γ, Ξ , etc.

3.3. Higher Hodge conjecture. For X smooth projective and defined over a subfield k of $\mathbb{C}$, the HC (Hodge Conjecture) states that $cl^{p,0} : CH^p(X_{(k)}) \rightarrow Hom_{\text{MHS}}(\mathbb{Q}(0), H^{2p}(X_{\mathbb{C}}^{an}, \mathbb{Q}(p)))$ should be surjective. Beilinson [Be] and Jannsen [Ja2] made a prediction extending this to all p, m and beyond the projective setting, provided the variety is arithmetic ($k \subseteq \bar{\mathbb{Q}}$).

Conjecture 3.6. *For $U/\bar{\mathbb{Q}}$ smooth quasiprojective,*

$$cl_U^{p,m} : CH^p(U, m) \rightarrow Hom_{\text{MHS}}(\mathbb{Q}(0), H^{2p-m}(U_{\mathbb{C}}^{an}, \mathbb{Q}(p)))$$

is surjective.

To show that this is “reasonable”, we relate it to the extended Bloch-Beilinson conjecture BBC (Conj. 3.4). The next result, due to M. Saito [SaM4] (by a somewhat less direct inductive proof), pops right out of the machinery of §§2, 3.1.

Proposition 3.7. *Conjecture 3.6 follows from HC and BBC.*

Proof. Let $\mathcal{X}/\bar{\mathbb{Q}}$ be a good compactification of U , i.e. $\mathcal{Y} := \mathcal{X} \setminus U$ is a NCD (also defined / $\bar{\mathbb{Q}}$). Let

$$\widetilde{Res}^k : W_b H_{\mathcal{K}}^{2p-k+b}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)) \longrightarrow \mathcal{K}(p)_{\infty}^{-k,b}$$

be the natural map. Consider the diagram

$$\begin{array}{ccc}
CH^p(\mathcal{X} \setminus \mathcal{Y}, m) & \xrightarrow{c_{\mathcal{H}}} & W_0 H_{\mathcal{H}}^{2p-m}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)) \\
\downarrow \widetilde{Res}^m & & \downarrow \widetilde{Res}^m \\
\left\{ \bigcap_{i \geq 2} \ker(d_i) \subseteq \mathcal{M}_{\mathcal{X} \setminus \mathcal{Y}}(p)_2^{-m,0} \right\} & \xrightarrow[\text{formula}]{\text{KLM}} & \left\{ \bigcap_{i \geq 2} \ker(d_i) \subseteq \mathcal{H}_{\mathcal{X} \setminus \mathcal{Y}}(p)_2^{-m,0} \right\} \\
\downarrow & & \downarrow \\
Hom_{\text{MHS}}(\mathbb{Q}(0), H^{2p-m}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p))) & \equiv & \left\{ \bigcap_{i \geq 1} \ker(\delta_{1-i}^{[-m]}) \subseteq (*) \right\}
\end{array}$$

cl (curved arrow from $CH^p(\mathcal{X} \setminus \mathcal{Y}, m)$ to $Hom_{\text{MHS}}(\mathbb{Q}(0), H^{2p-m}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)))$)

where $\delta_j^{[-m]}$ is defined in §2 and $(*) =$

$$Hom_{\text{MHS}}(\mathbb{Q}(0), \mathcal{B}_{\mathcal{X} \setminus \mathcal{Y}}(p)_2^{-m,0}) \cong \frac{\ker\{Gy : Hg^{p-m}(\widetilde{\mathcal{Y}}^m) \rightarrow Hg^{p-m+1}(\widetilde{\mathcal{Y}}^{m-1})\}}{\text{im}\{Gy : Hg^{p-m-1}(\widetilde{\mathcal{Y}}^{m+1}) \rightarrow Hg^{p-m}(\widetilde{\mathcal{Y}}^m)\}}.$$

(We have made use of the proof of Prop. 2.7. Note that $Hg^q(Y) := Hom_{\text{MHS}}(\mathbb{Q}(0), H^{2q}(Y, \mathbb{Q}(q)))$.) Also consider

$$\begin{array}{ccc}
CH^{p-m}(\widetilde{\mathcal{Y}}^m) & \xrightarrow{Gy=d_1} & CH^{p-m+1}(\widetilde{\mathcal{Y}}^{m-1}) \\
\downarrow cl^{p-m} & & \downarrow cl^{p-m+1} \\
Hg^{p-m}(\widetilde{\mathcal{Y}}^m) & \xrightarrow{Gy} & Hg^{p-m+1}(\widetilde{\mathcal{Y}}^{m-1}) \\
\uparrow & & \uparrow \\
H_{\mathcal{H}}^{2(p-m)}(\widetilde{\mathcal{Y}}^m, \mathbb{Q}(p-m)) & \xrightarrow{Gy=d_1} & H_{\mathcal{H}}^{2(p-m+1)}(\widetilde{\mathcal{Y}}^{m-1}, \mathbb{Q}(p-m+1)) \\
\uparrow & & \uparrow \\
Ext_{\text{MHS}}^1(\mathbb{Q}(0), H^{2p-2m-1}(\widetilde{\mathcal{Y}}^m, \mathbb{Q}(p-m))) & \xrightarrow{Gy} & Ext_{\text{MHS}}^1(\mathbb{Q}(0), H^{2p-2m+1}(\widetilde{\mathcal{Y}}^{m-1}, \mathbb{Q}(p-m+1)))
\end{array}$$

$c_{\mathcal{H}}^{p-m}$ (curved arrow from $CH^{p-m}(\widetilde{\mathcal{Y}}^m)$ to $Hg^{p-m}(\widetilde{\mathcal{Y}}^m)$)

$c_{\mathcal{H}}^{p-m+1}$ (curved arrow from $CH^{p-m+1}(\widetilde{\mathcal{Y}}^{m-1})$ to $Hg^{p-m+1}(\widetilde{\mathcal{Y}}^{m-1})$)

By the HC, there exists (with $n = \dim(X)$) $\Gamma \in CH^{n-m}(\widetilde{\mathcal{Y}}^{m-1} \times \widetilde{\mathcal{Y}}^m)$ inducing an inverse of $Gy : H^{2p-2m-1}(\widetilde{\mathcal{Y}}^m) \rightarrow H^{2p-2m+1}(\widetilde{\mathcal{Y}}^{m-1})$ on its image, and the 0-map in other degrees.

Given $\xi \in Hom_{\text{MHS}}(\mathbb{Q}(0), H^{2p-m}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)))$, let $\tilde{\xi}$ be a “lift” to $\ker(Gy) \subseteq Hg^{p-m}(\widetilde{\mathcal{Y}}^m)$. By HC for cl^{p-m} , we have $\mathfrak{W} \in CH^{p-m}(\widetilde{\mathcal{Y}}^m)$ with $cl^{p-m}(\mathfrak{W}) = \tilde{\xi}$. One easily checks that $c_{\mathcal{H}}^{p-m+1}$ of $\tilde{\mathfrak{W}} := \mathfrak{W} - \Gamma_* Gy(\mathfrak{W})$ lies in $\ker(Gy)$. By BBC for $c_{\mathcal{H}}^{p-m+1}$, so does $\tilde{\mathfrak{W}}$ itself. (Of course, still $cl^{p-m}(\tilde{\mathfrak{W}}) = \tilde{\xi}$.)

Now $\tilde{\xi} \in \bigcap_{i \geq 1} \ker(\delta_{1-i}^{[-m]}) \subset (*) \implies c_{\mathcal{H}}^{p-m}(\tilde{\mathfrak{W}}) \in \bigcap_{i \geq 1} \ker(d_i)$, again using the proof of Prop. 2.7. So for $i \geq 2$, we have

$$c_{\mathcal{H}}^{p-m+i, i-1}(d_i(\tilde{\mathfrak{W}})) \in \operatorname{im} \left\{ \begin{array}{c} \operatorname{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), H^{2p-2m+i-2}(\widetilde{\mathcal{Y}^{m-i+1}}, \mathbb{Q}(p-m+i-1))) \\ \xrightarrow{Gy} \operatorname{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), H^{2p-2m+i}(\widetilde{\mathcal{Y}^{m-i}}, \mathbb{Q}(p-m+i))) \end{array} \right\}.$$

Let Γ^i be an algebraic cycle inverse to Gy as above; then writing $\widetilde{d_i(\tilde{\mathfrak{W}})}$ for a representative (of $d_i(\tilde{\mathfrak{W}})$) in $CH^{p-m+i}(\widetilde{\mathcal{Y}^{m-i}}, i-1)$, $c_{\mathcal{H}}\{\widetilde{d_i(\tilde{\mathfrak{W}})} - Gy(\Gamma^i \widetilde{d_i(\tilde{\mathfrak{W}})})\} = 0$. Using BBC (for $c_{\mathcal{H}, \widetilde{\mathcal{Y}^{m-i}}}^{p-m+i, i-1}$) gives $d_i(\tilde{\mathfrak{W}}) \in \operatorname{im}(Gy = d_1) \implies d_i(\tilde{\mathfrak{W}}) = 0$. Hence $\tilde{\mathfrak{W}}$ survives to ∞ , and is Gr_0^W of a class in $CH^p(\mathcal{X} \setminus \mathcal{Y}, m)$. $\square$

Remark 3.8. One case where HC and BBC are actually true in all places they are used (in the above proof), is where $\mathcal{X}$ and all irreducible components of each $\mathcal{Y}^j$ are rational.

If (say) $\mathcal{Y}$ is not defined over $\bar{\mathbb{Q}}$, it is easy to construct counterexamples to the Conjecture.

Example 3.9. Let $V_{/\bar{\mathbb{Q}}} \subset \mathbb{P}^3$ be a smooth hypersurface of degree at least 4, with $P \in V(\mathbb{Q})$, and $Q \in V(\mathbb{C})$ very general. Let L_P and L_Q be disjoint lines through P and Q (resp.) intersecting V transversely. (Note that L_Q cannot be defined $/\bar{\mathbb{Q}}$.) We claim the cycle-class maps

$$(3.4) \quad CH^2(V \setminus \{P, Q\}, 1) \rightarrow \operatorname{Hom}_{\text{MHS}}(\mathbb{Q}(0), H^3(V \setminus \{P, Q\}, \mathbb{Q}(2)))$$

$$(3.5) \quad CH^3(\mathbb{P}^3 \setminus V \cup L_P \cup L_Q, 2) \rightarrow \operatorname{Hom}_{\text{MHS}}(\mathbb{Q}(0), H^4(\mathbb{P}^3 \setminus V \cup L_P \cup L_Q, \mathbb{Q}(3)))$$

are not surjective. Consider the blow-ups

$$\beta_V : B_{P \amalg Q}(V) =: \tilde{V} \longrightarrow V, \quad \beta_{\mathbb{P}} : B_{L_Q \amalg L_P}(\mathbb{P}^3) =: \tilde{\mathbb{P}}^3 \longrightarrow \mathbb{P}^3,$$

with (resp.) exceptional divisors $\ell_P \amalg \ell_Q$ and $S_P \amalg S_Q$; set $'\tilde{V} := \beta_{\mathbb{P}}^{-1}(V)$.

For (3.4), let $\mathcal{X} = \tilde{V}$, $\mathcal{Y} = \ell_P \amalg \ell_Q (= \widetilde{\mathcal{Y}^1})$; take $\tilde{\xi} \in Hg^1(\mathcal{Y})$ to be the class of a difference of points (-pt. on ℓ_P , +pt. on ℓ_Q). Then in fact $\tilde{\xi} \in \ker\{Gy = d_1 : H_{\mathcal{H}}^2(\mathcal{Y}, \mathbb{Q}(1)) \rightarrow H_{\mathcal{H}}^4(\mathcal{X}, \mathbb{Q}(2))\}$, so $\tilde{\xi}$ determines a class in the r.h.s. of (3.4). But any $\mathcal{Z} \in CH^1(\mathcal{Y})$ with $cl^1(\mathcal{Z}) = \tilde{\xi}$ has $(\beta_V)_* Gy(\mathcal{Z}) = Q - P \in CH^2(V_{\mathbb{C}})$ which is $\stackrel{\text{rat}}{\neq} 0$; so $d_1(\mathcal{Z}) \neq 0$ and $\mathcal{Z}$ cannot be Gr_0^W of a class in the l.h.s. of (3.4).

For (3.5), take $'\mathcal{X} = \tilde{\mathbb{P}}^3$, $'\mathcal{Y} = '\tilde{V} \cup S_P \cup S_Q \implies '\widetilde{\mathcal{Y}^1} = '\tilde{V} \amalg S_P \amalg S_Q$, $'\widetilde{\mathcal{Y}^2} =$ disjoint union of lines on $'\tilde{V}$ (including ℓ_P, ℓ_Q). Our $\tilde{\xi}$ from above maps to a $'\tilde{\xi} \in Hg^1(' \widetilde{\mathcal{Y}^2}) \cong H_{\mathcal{H}}^2(' \widetilde{\mathcal{Y}^2}, \mathbb{Q}(1))$, $Gy (= d_1)$ of which is 0 in

$H_{\mathcal{H}}^4(' \widetilde{\mathcal{Y}}^1, \mathbb{Q}(2)) \cong Hg^2(' \widetilde{\mathcal{Y}}^1)$. Moreover, d_2 maps $'\tilde{\xi}$ to

$$\frac{H_{\mathcal{H}}^5(' \mathcal{X}, \mathbb{Q}(3))}{Gy\{H_{\mathcal{H}}^3(' \widetilde{\mathcal{Y}}^1, \mathbb{Q}(2))\}} \cong \frac{H^4(' \mathcal{X}, \mathbb{C}/\mathbb{Q}(3))}{Gy\{H^2(' \widetilde{\mathcal{Y}}^1, \mathbb{C})\}} = 0;$$

so we get a class in the r.h.s. of (3.5). But $d_1 = Gy$ of any $'\mathcal{Z} \in CH^1(' \widetilde{\mathcal{Y}}^2)$ with $cl^1(' \mathcal{Z}) = '\tilde{\xi}$, maps to $Q - P \not\equiv 0$ under

$$CH^2(' \widetilde{\mathcal{Y}}^1) \xrightarrow{\text{restrict}} CH^2(' \tilde{V}) \xrightarrow{(\beta_{\mathbb{P}}|_{' \tilde{V}})_*} CH^2(V_{\mathbb{C}});$$

so once again we are done.

3.4. Coniveau filtration. To conclude this section we mention an alternative approach to sequential residue maps on $H_{\mathcal{K}}^{2p+*}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p))$. While the Gysin/weight setup above has the highest-codimension residue defined first, the approach via coniveau reverses this (so that codim.-1 residues are defined first). The drawback is that what follows is valid only for $\mathcal{K} = \mathcal{M}, \mathcal{D}$, and $\mathcal{B}$ (not $\mathcal{H}$).

Define

$$N^k C_{\mathcal{K}}^{2p+*}(\mathcal{X}, \mathbb{Q}(p)) := \text{im} \left\{ (2\pi i)^k \iota_*^{\widetilde{\mathcal{Y}}^k} : C_{\mathcal{K}}^{2p-2i+*}(\widetilde{\mathcal{Y}}^i, \mathbb{Q}(p-i)) \rightarrow C_{\mathcal{K}}^{2p+*}(\mathcal{X}, \mathbb{Q}(p)) \right\};$$

taking

$$' \mathcal{K}(p)_0^{a,b} := Gr_N^a C_{\mathcal{K}}^{2p+a+b}(\mathcal{X}, \mathbb{Q}(p)), \quad d_0 := (Gr_N^a) d_{\mathcal{K}}$$

then yields (4th quadrant) spectral sequences with

$$' \mathcal{K}(p)_1^{a,b} \cong H_{\mathcal{K}}^{2p-a+b}(\mathcal{Y}^a \setminus \mathcal{Y}^{a+1}, \mathbb{Q}(p-a)),$$

$$' \mathcal{K}(p)_{\infty}^{a,b} \cong Gr_N^a H_{\mathcal{K}}^{2p+a+b}(\mathcal{X}, \mathbb{Q}(p)).$$

The higher differentials d_r on $' \mathcal{K}(p)_r^{a,b}$ then yield the desired residue maps

$$'Res^r : \left\{ \ker('Res^{r-1}) \subseteq H_{\mathcal{K}}^{2p+b}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p)) \right\} \\ \rightarrow \left\{ \begin{array}{c} \text{subquotient of} \\ H_{\mathcal{K}}^{2p+b-2k+1}(\mathcal{Y}^r \setminus \mathcal{Y}^{r+1}, \mathbb{Q}(p-r)) \end{array} \right\}$$

with $\bigcap \ker('Res^r) = \underline{H_{\mathcal{K}}^{2p+b}}(\mathcal{X} \setminus \mathcal{Y}, \mathbb{Q}(p))$. The $'Res^r$ have been studied for $\mathcal{K} = \mathcal{M}, \mathcal{D}$ (in a special case) in [Ke4].

It is intriguing to compare the length of the filtration by their kernels to that of the “weight” filtration of §3.1. To determine that an element of $CH^p(\mathcal{X} \setminus \mathcal{Y}, m)$ comes from $CH^p(\mathcal{X}, m)$, one must check the vanishing of m higher $\widetilde{Res}^k$ maps (see §3.3) or $[p - \frac{m}{2}]$ (where $[\cdot]$ = greatest-integer function) higher $'Res^r$ maps.

4. HIGHER NORMAL FUNCTIONS

The next two sections describe our take on higher Abel-Jacobi maps for higher Chow groups — invariants that detect cycles in the regulator (or AJ) kernel. Here we will adopt the formalism of M. Saito’s mixed Hodge modules [SaM2], while §5 (which is more “computational”) reverts to absolute Hodge cohomology in a special (but broadly applicable) case.

4.1. Spreads and the Leray filtration. We first recall the relationship between these, for W (defined over $k \subset \mathbb{C}$) a smooth quasi-projective variety; for us $\mathrm{MHM}(W)$ will always mean $\mathrm{MHM}(W_{\mathbb{C}})$. There is an equivalence of categories between $\mathrm{MHM}(pt.)$ and graded-polarizable mixed Hodge structures (PMHS), so that push-forward along the structure morphism $a_W : W_{\mathbb{C}} \rightarrow \mathrm{Spec} \mathbb{C}$ yields $(a_W)_* : \mathrm{MHM}(W) \rightarrow D^b\mathrm{PMHS}$ satisfying $H^i(a_W)_* \mathbb{Q}_W(p) \cong H^i(W, \mathbb{Q}(p))$. Writing $H^0 := H^{2p-m}(W, \mathbb{Q}(p))$, $H^{-1} := H^{2p-m-1}(W, \mathbb{Q}(p))$, we have a diagram with short-exact rows: (4.1)

$$\begin{array}{ccccc}
 \mathrm{Ext}_{\mathrm{MHS}}^1(\mathbb{Q}(0), H^{-1}) & & H_{\mathcal{H}}^{2p-m}(W, \mathbb{Q}(p)) & & \\
 \parallel & & \parallel & & \\
 \frac{W_0 H_{\mathbb{C}}^{-1}}{F^0 W_0 H_{\mathbb{C}}^{-1} + W_0 H^{-1}} \longrightarrow & \mathrm{Ext}_{D^b\mathrm{MHS}}^{2p-m}(\mathbb{Q}(0), (a_W)_* \mathbb{Q}_W(p)) \longrightarrow & \mathrm{Hom}_{\mathrm{MHS}}(\mathbb{Q}(0), H^0) & & \\
 \uparrow & \uparrow & \parallel & & \\
 \frac{W_{-1} H_{\mathbb{C}}^{-1}}{\{F^0 W_0 H_{\mathbb{C}}^{-1} + W_0 H^{-1}\} \cap W_{-1} H_{\mathbb{C}}^{-1}} \longrightarrow & \mathrm{Ext}_{D^b\mathrm{PMHS}}^{2p-m}(\mathbb{Q}(0), (a_W)_* \mathbb{Q}_W(p)) \longrightarrow & \mathrm{Hom}_{\mathrm{PMHS}}(\mathbb{Q}(0), H^0) & & \\
 \parallel & (*) \parallel & & & \\
 \mathrm{Ext}_{\mathrm{PMHS}}^1(\mathbb{Q}(0), H^{-1}) & & \mathrm{Ext}_{\mathrm{MHM}(W)}^{2p-m}(\mathbb{Q}_W(0), \mathbb{Q}_W(p)) & &
 \end{array}$$

where $(*)$ follows from $\mathbb{Q}_W(0) = (a_W)^* \mathbb{Q}(0)$ and adjointness of $(a_W)^*$, $(a_W)_*$. There is then a cycle-class map (described in [As])

$$c_{\mathrm{MHM}}^W : CH^p(W, m) \rightarrow \mathrm{Ext}_{\mathrm{MHM}(W)}^{2p-m}(\mathbb{Q}_W(0), \mathbb{Q}_W(p))$$

factoring $c_{\mathcal{H}}$ (as defined in §3, thinking of W as having a good compactification).

Next, we briefly revisit the approach to spreads discussed in [Ke2]. Begin with the following data:

- fields $k \subset K \subset \mathbb{C}$, k algebraically closed and K/k finitely generated of transcendence degree t ;
- a smooth projective variety X defined $/K$, of dimension d ; and
- a higher Chow (pre)cycle $\mathcal{Z} \in Z^p(X_{(K)}, m)$ with $\partial_{\mathcal{B}} \mathcal{Z} = 0$.

The theory of spreads then provides:

- a smooth projective variety $\mathcal{S}/k$, with an isomorphism $k(\mathcal{S}) \rightarrow K$; and $p_g : \text{Spec} K \rightarrow \mathcal{S}$ the corresponding generic point. (Clearly $\dim_k \mathcal{S} = t$.) Denote by $U \subset \mathcal{S}$ any affine Zariski-open subvariety defined $/k$.
- $\mathcal{X}$ smooth projective and $\pi : \mathcal{X} \rightarrow \mathcal{S}$ projective, both defined $/k$, s.t. $X = \mathcal{X} \times_{p_g} \text{Spec} K$. Write $\pi^{-1}(U) =: \mathcal{X}_U$.
- $\bar{\mathfrak{Z}} \in Z^p(\mathcal{X}, m)$ with $\mathcal{Z} = \bar{\mathfrak{Z}} \times_{p_g} \text{Spec} K$, and some $U_0 \subset \mathcal{S}$ (as above) s.t. $\bar{\mathfrak{Z}}_{U_0} := \bar{\mathfrak{Z}}|_{\mathcal{X}_{U_0}}$ is $\partial_{\mathcal{B}}$ -closed. [Note: by $\bar{\mathfrak{Z}}$ we do not necessarily mean the Zariski closure, though that would be one possible choice.]

Remark 4.1. It is possible that none of the many choices of $\bar{\mathfrak{Z}}$ is $\partial_{\mathcal{B}}$ -closed; on the other hand, if such a choice (with $\partial_{\mathcal{B}} \bar{\mathfrak{Z}} = 0$) does exist, we call it a *complete k -spread* of $\mathcal{Z}$. Existence or nonexistence of a complete spread depends only on the class of $\mathcal{Z}$ in $CH^p(X_{(K)}, m)$.

To get rid of ambiguities (i.e., the nonuniqueness of $\mathcal{X}$, $\mathcal{S}$, $\bar{\mathfrak{Z}}$, etc.), we take the limit over $U \subset \mathcal{S}$ (U as above): $\eta_{\mathcal{S}} := \varprojlim U$, $\mathcal{X}_{\eta} := \varprojlim \mathcal{X}_U$. These have precise meaning only under a contravariant functor to abelian groups,⁵ e.g.

$$Ext_{\text{MHS}}^1(\mathbb{Q}(0), H^a(\eta_{\mathcal{S}}, R^b \pi_* \mathbb{Q}(p))) := \varinjlim_U Ext_{\text{MHS}}^1(\mathbb{Q}(0), H^a(U, R^b \pi_* \mathbb{Q}(p))),$$

$$H_{DR}^*(\mathcal{X}_{\eta}, \mathbb{C}) := \varinjlim_U H^*(\mathcal{D}^{\bullet}((\mathcal{X}_U)_{\mathbb{C}}^{an})),$$

$$CH^p(\mathcal{X}_{\eta}, m) := \varinjlim_U CH^p(\mathcal{X}_U, m) \xrightarrow[p_g]{\cong} CH^p(X, m).$$

The k -spread $\bar{\mathfrak{Z}}$ of $\mathcal{Z}$ is just the restriction of $\bar{\mathfrak{Z}}_{U_0}$ to $Z^p(\mathcal{X}_{\eta}, m)$; writing $\mathfrak{s} := (p_g^*)^{-1}$, the class $\langle \bar{\mathfrak{Z}} \rangle \in CH^p(\mathcal{X}_{\eta}, m)$ (frequently denoted just $\bar{\mathfrak{Z}}$) is then $\mathfrak{s} \langle \mathcal{Z} \rangle$. Note that the complex points of $\mathcal{S}$ that survive in the limit over U are the very general points (of maximal transcendence degree), corresponding to the various embeddings of $k(\mathcal{S})$ into $\mathbb{C}$.

The spectral sequence

$$(4.2) \quad E_2^{i,j} := Ext_{\text{MHM}(\eta_{\mathcal{S}})}^i \left(\mathbb{Q}_{\eta}(0), H^j \pi_* \mathbb{Q}_{\mathcal{X}_{\eta}}(p) \right)$$

computing

$$(4.3) \quad Ext_{\text{MHM}(\mathcal{X}_{\eta})}^{2p-m} \left(\mathbb{Q}_{\mathcal{X}_{\eta}}(0), \mathbb{Q}_{\mathcal{X}_{\eta}}(p) \right) \cong Ext_{D^b \text{MHM}(\eta_{\mathcal{S}})}^{2p-m} \left(\mathbb{Q}_{\eta}(0), \pi_* \mathbb{Q}_{\mathcal{X}_{\eta}}(p) \right)$$

⁵the resulting direct limits are exact in Ab.

degenerates at E_2 since π is proper; so the resulting Leray filtration $\mathcal{L}^\bullet$ on (4.3) has $Gr_{\mathcal{L}}^i(4.3) \cong E_{2(=\infty)}^{i, 2p-m-i}$. Set $\mathcal{L}^i CH^p(X, m) := (c_{\text{MHM}}^{\mathcal{X}_\eta} \circ \mathfrak{s})^{-1} \mathcal{L}^i$, so that $c_{\text{MHM}}^{\mathcal{X}_\eta}$ induces

$$c_{\text{MHM}}^{X, i} : Gr_{\mathcal{L}}^i CH^p(X, m) \hookrightarrow Ext_{\text{MHM}(\eta_S)}^i(\mathbb{Q}_\eta(0), H^{2p-m-i} \pi_* \mathbb{Q}_{\mathcal{X}_\eta}(p)).$$

The filtration on $CH^p(X, m)$ stabilizes $\mathcal{L}^i = \mathcal{L}^{i+1} \forall i \geq \min\{t+2, p+1\}$ by a standard argument (using relative hard Lefschetz for $\mathcal{X}_\eta \rightarrow \eta_S$, see [Ke1] or [Le1]). If $k = \mathbb{Q}$ then the “quasi-projective BBC” of §3.2 implies that $c_{\mathcal{H}}^{\mathcal{X}_\eta}$, hence $c_{\text{MHM}}^{\mathcal{X}_\eta}$, is injective; so $\mathcal{L}^\bullet$ (conjecturally) stabilizes at $\{0\}$, hence should be viewed as a candidate Bloch-Beilinson filtration.

Remark 4.2. Note that this is essentially the filtration described in [As], except without the limit over all $K \subseteq \mathbb{C}$ finitely generated over k (and containing the minimal field of definition of $X, \mathcal{Z}$).

A second spectral sequence (for the structure morphism $a_\eta : \eta_S \rightarrow \text{Spec } k$) yields short-exact sequences $\underline{E}_\infty^{i,j} \hookrightarrow E_\infty^{i,j} \xrightarrow{p^{i,j}} \underline{\underline{E}}_\infty^{i,j}$ where

$$\underline{E}_\infty^{i,j} = Ext_{\text{PMHS}}^1(\mathbb{Q}(0), H^{i-1}(\eta_S, R^j \pi_* \mathbb{Q}(p))),$$

$$\underline{\underline{E}}_\infty^{i,j} = Hom_{(\text{P})\text{MHS}}(\mathbb{Q}(0), H^i(\eta_S, R^j \pi_* \mathbb{Q}(p))).$$

(Arapura [Ar2] has proved that the MHS on $H^i(U, R^j \pi_* \mathbb{Q}(p))$ arising from Saito’s theory agrees with the “natural” MHS coming from his geometric definition [Ar1] of the Leray filtration on $H^*(\mathcal{X}_U, \mathbb{Q}(p))$.) In particular, the $[\mathfrak{Z}]_i := p^{i, 2p-m-i}(c_{\text{MHM}}^{X, i}(\mathcal{Z}))$ are just Leray graded pieces of the fundamental class $[\mathfrak{Z}] \in Hom_{\text{MHS}}(\mathbb{Q}(0), H^{2p-m}(\mathcal{X}_\eta, \mathbb{Q}(p)))$. Both are obviously (well-)defined (given $\langle \mathcal{Z} \rangle$) without recourse to any of the theory just discussed (except spreads).

4.2. A geometric construction: the Λ -filtration. We now propose a notion of *higher normal functions* having the $[\mathfrak{Z}]_i$ as “topological invariants”, which arose from an attempt to extend Arapura’s geometric approach to Leray to absolute Hodge cohomology. The definition does not use MHM, and leads to a more concrete description of $\mathcal{L}^\bullet$.

Recall that a complex subvariety $\mathcal{T}_{\mathbb{C}} \subset \mathcal{S}_{\mathbb{C}}$ is called *very general* if no rational function $f \in \bar{\mathbb{Q}}(\mathcal{S})^*$ has $f|_{\mathcal{T}_{\mathbb{C}}} \equiv 0$. Equivalently, the minimal field of definition $L(\subset \mathbb{C})$ of $\mathcal{T}$ has $\text{trdeg}(L/k) = \text{codim}_{\mathcal{S}}(\mathcal{T})$; $\mathcal{T}$ always means $\mathcal{T}_L$, and $\eta_{\mathcal{T}} := \varprojlim \mathcal{V}$ over $\mathcal{V} \subset \mathcal{T}$ affine Zariski open $/L$. Write π (by abuse of notation) for the proper morphism $\mathcal{X}_{\eta_{\mathcal{T}}} \rightarrow \eta_{\mathcal{T}}$, and $\mathcal{S}[i]$ for the set of $(i-1)$ -dimensional very general subvarieties of $\mathcal{S}_{(\mathbb{C})}$.

Definition 4.3. If $[\mathfrak{Z}]_0 = \cdots = [\mathfrak{Z}]_{i-1} = 0$, the i^{th} higher normal function

$$\nu_3^i : \mathcal{S}[i] \rightarrow \coprod_{\mathcal{T} \in \mathcal{S}[i]} \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), H^{2p-m-1}(\mathcal{X}_{\eta_{\mathcal{T}}}, \mathbb{Q}(p)))$$

associated to $\mathfrak{Z}$, is given by $\nu_3^i(\mathcal{T}) := AJ(\mathfrak{Z}|_{\mathcal{X}_{\eta_{\mathcal{T}}}})$.

Remark 4.4. This can be “higher” in two senses ($i > 1$ and $m > 0$). If $i = 1$ and $m = 0$, ν_3^1 is essentially a classical normal function, restricted to $\eta_{\mathcal{S}}(\mathbb{C}) = \varprojlim U(\mathbb{C}) = \mathcal{S}[1]$.

To see that this is well-defined, note that $\pi : \mathcal{X}_{\eta} \rightarrow \eta (= \eta_{\mathcal{T}} \text{ or } \eta_{\mathcal{S}})$ is proper and smooth, so that $E_2^{a,b} = H^a(\eta, R^b \pi_* \mathbb{Q}(p))$ converges at E_2 and the (canonical) Leray filtration has $\text{Gr}_{\mathcal{L}}^a H^{a+b}(\mathcal{X}_{\eta}, \mathbb{Q}(p)) = E_{2,\eta}^{a,b}$. If $[\mathfrak{Z}]_a \in E_{2,\eta_{\mathcal{S}}}^{a,2p-m-a}$ vanishes for $0 \leq a \leq i-1$, then so do $[\mathfrak{Z}|_{\mathcal{X}_{\eta_{\mathcal{T}}}}] \in E_{2,\eta_{\mathcal{T}}}^{a,2p-m-a}$, while $E_{2,\eta_{\mathcal{T}}}^{a,2p-m-a} = 0$ for $a \geq i$ (as $\dim(\eta_{\mathcal{T}}) = i-1$ and $\eta_{\mathcal{T}}$ is a limit of affines); hence $[\mathfrak{Z}|_{\mathcal{X}_{\eta_{\mathcal{T}}}}] = 0$ and AJ is defined.

Moreover, we have (noncanonically) $H^*(\mathcal{X}_{\eta}, \mathbb{Q}(p)) \cong \oplus_{a+b=*} E_{2,\eta}^{a,b}$ as MHs (with $\mathcal{L}^q = \oplus_{a \geq q}$), so that all $\mathcal{L}^{q+1} \rightarrow \mathcal{L}^q \rightarrow \text{Gr}_{\mathcal{L}}^q$ split and $L^a \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), H^{a+b}(\mathcal{X}_{\eta}, \mathbb{Q}(p))) := \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), \mathcal{L}^a H^{a+b})$ satisfies $\text{Gr}_{\mathcal{L}}^a \cong \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), E_{2,\eta}^{a,b}) =: \mathcal{E}_{2,\eta}^{a,b}$. If $\mathcal{T}' \subset \mathcal{T}$ is a (very general) hyperplane section, consider $\theta^n : H^n(\mathcal{X}_{\eta_{\mathcal{T}}}, \mathbb{Q}(p)) \rightarrow H^n(\mathcal{X}_{\eta_{\mathcal{T}'}} , \mathbb{Q}(p))$, with $\theta^{a,b} := \text{Gr}_{\mathcal{L}}^a \theta^{a+b} : E_{2,\eta_{\mathcal{T}}}^{a,b} \rightarrow E_{2,\eta_{\mathcal{T}'}}^{a,b}$ injective for $a \leq i-2 (= \dim(\mathcal{T}'))$. If also $2a+b < 2p$, then weights of $E_{2,\eta_{\mathcal{T}}}^{a,b}$ are < 0 , so that $\text{Hom}_{\text{MHS}}(\mathbb{Q}(0), \text{coker}(\theta)) = 0$ and $\mathcal{E}_{2,\eta_{\mathcal{T}}}^{a,b} \xrightarrow[\text{Ext}^1(\theta^{a,b})]{} \mathcal{E}_{2,\eta_{\mathcal{T}'}}^{a,b}$. In particular, if $i \leq m+2$, then $\text{Ext}^1(\theta^{a,2p-m-a-1})$ is injective for $0 \leq a \leq i-2$, while $\mathcal{E}_{2,\eta_{\mathcal{T}}}^{a,2p-m-a-1}$ (resp. $\mathcal{E}_{2,\eta_{\mathcal{T}'}}^{a,2p-m-a-1}$) is 0 for $a \geq i$ (resp. $i-1$); thus $\ker(\text{Ext}^1(\theta^{2p-m-1})) \cong \mathcal{E}_{2,\eta_{\mathcal{T}}}^{i-1,2p-m-i}$. This proves (a) of

Proposition 4.5. Assume $[\mathfrak{Z}]_0 = \cdots = [\mathfrak{Z}]_{i-1} = 0$.

(a) If $\nu_3^{i-1} = 0$ and $i \leq m+2$, then ν_3^i factors through

$$\coprod_{\mathcal{T} \in \mathcal{S}[i]} \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), H^{i-1}(\eta_{\mathcal{T}}, R^{2p-m-1} \pi_* \mathbb{Q}(p)));$$

(b) If $\nu_3^i = 0$, then $[\mathfrak{Z}]_i = 0$. In fact, $[\mathfrak{Z}]_i$ is the “topological invariant” of ν_3^i (to be defined below).

Assuming (b), we can use the normal functions to define a filtration $\Lambda^\bullet$ without reference to vanishing of the $\{[\mathfrak{Z}]_i\}$.

Definition 4.6. Set $\Lambda^0 CH^p(X, m) = CH^p(X, m)$, $\Lambda^1 CH^p(X, m) = CH_{hom}^p(X, m)$ ($\Lambda^0 = \Lambda^1$ if $m > 0$), and

$$\Lambda^i CH^p(X, m) := \left\{ \mathcal{Z} \in CH_{hom}^p(X, m) \mid \nu_{\mathfrak{s}(\mathcal{Z})}^1 = \cdots = \nu_{\mathfrak{s}(\mathcal{Z})}^{i-1} = 0 \right\}.$$

To facilitate the proof of (b), we restrict the domains of the $\{\nu_3^i\}$. Choose nested very general hyperplane sections $(\mathcal{S}_{\mathbb{C}} \supsetneq) \mathcal{T}_{\mathbb{C}}^1 \supsetneq \mathcal{T}_{\mathbb{C}}^2 \supsetneq \cdots \supsetneq \mathcal{T}_{\mathbb{C}}^{t-1}$, with $\mathcal{T}^j$ of codimension j and minimal field of definition L_j . These satisfy $k \subsetneq L_1 \subsetneq L_2 \subsetneq \cdots \subsetneq L_{t-1} \subsetneq K$ with $\text{trdeg}(L_j/k) = j$. One can show that there exist smooth affine varieties and morphisms

$$\begin{array}{ccccccc} & \mathcal{M}_0 & & & & & \mathcal{M}_t \\ & \parallel & & \xrightarrow{\rho_2} & & & \parallel \\ (\mathcal{S} \longleftarrow) U & \xrightarrow{\rho_1} & \mathcal{M}_1 & \xrightarrow{\rho_{12}} & \mathcal{M}_2 & \longrightarrow \cdots \longrightarrow & \mathcal{M}_{t-1} \longrightarrow \text{Spec}(k) \\ & & & \xrightarrow{\rho_{t-1}} & & & \end{array}$$

defined $/k$ (with $\dim(\mathcal{M}_j) = t - j$, $k(\mathcal{M}_j) \cong L_j$), such that fibers of each ρ_j are affine j -dimensional (multiple-)hyperplane sections of U . (In particular, $(\mathcal{T}^j \cap U)_{\mathbb{C}} = \rho_j^{-1}(q_j)$ for some very general $q_j \in \mathcal{M}_j(\mathbb{C})$.) Denote the restriction $\eta_{\mathcal{S}} \rightarrow \eta_{\mathcal{M}_j}$ of ρ_j by ρ_j^{η} . Given $\mu \in \mathcal{M}_j$, write $\mathcal{T}_{\mu} := \rho_j^{-1}(\mu)$, and set $H_{\mu} := H^{2p-m-1}(\mathcal{X}_{\eta_{\mathcal{T}_{\mu}}}, \mathbb{Q}(p))$.

The restricted normal function

$$\bar{\nu}_3^i : \eta_{\mathcal{M}_{i-1}}(\mathbb{C}) \rightarrow \coprod_{\mu \in \eta_{\mathcal{M}_{i-1}}(\mathbb{C})} \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), H_{\mu})$$

still clearly satisfies Prop. 4.5(a) (this will be important later). Define a local system on $\eta_{\mathcal{M}_{i-1}}$ with stalks $W_0 H_{\mu}$ by $\mathbb{H}_i := W_0 R^{2p-m-1}(\rho_{i-1}^{\eta} \circ \pi)_* \mathbb{Q}(p)$, and set $\mathcal{H}_i := \mathbb{H}_i \otimes \mathcal{O}_{\eta_{\mathcal{M}_{i-1}}}$; take $\mathcal{F}_i \subset \mathcal{H}_i$ to be the subsheaf with stalks $F^p W_{2p} H^{2p-m-1}(\mathcal{X}_{\eta_{\mathcal{T}_{\mu}}}, \mathbb{C})$, and

$$\mathbb{F}_i := \ker \left\{ \mathbb{H}_i \rightarrow \frac{\mathcal{H}_i}{\mathcal{F}_i} \right\}, \quad \mathcal{J}_i := \text{coker} \left\{ \frac{\mathbb{H}_i}{\mathbb{F}_i} \hookrightarrow \frac{\mathcal{H}_i}{\mathcal{F}_i} \right\}.$$

Here $\mathcal{J}_i$ has stalks $\mathcal{J}^{p,m}(\mathcal{X}_{\eta_{\mathcal{T}_{\mu}}})$, $\mathbb{F}$ is a VMHS of pure type $(0, 0)$ (since its stalks are $\text{Hom}_{\text{MHS}}(\mathbb{Q}(0), H_{\mu})$), and $\bar{\nu}_3^i$ may be viewed as a holomorphic section of $\mathcal{J}_i$. Applying the connecting homomorphism

$$H^0(\eta_{\mathcal{M}_{i-1}}, \mathcal{J}_i) \xrightarrow{\delta} H^1 \left(\eta_{\mathcal{M}_{i-1}}, \frac{\mathbb{H}_i}{\mathbb{F}_i} \right)$$

gives the *topological invariant* $\delta \bar{\nu}_3^i$.

Thinking of $\eta_{\mathcal{M}}$ as some sufficiently small affine $\mathcal{V} \subset \mathcal{M}_{i-1}$ (in the limit), cover it with analytic balls V_{α} . Notice that $\frac{\mathcal{H}_i}{\mathcal{F}_i} \cong \mathbb{R}^{-m}(\rho_{i-1}^{\eta} \circ \pi)_* \mathbf{s}^{\bullet} \tilde{\mathcal{I}}_{(\mathcal{X}_{\eta_{\mathcal{S}}/\eta_{\mathcal{M}_{i-1}}})}(p)$, so that we can apply the version of AJ via (3.2)

continuously fiberwise over each V_α to get a section (omitting the “ $\bullet$ ”) $(2\pi i)^{p-m} \{R_{\alpha,\mu} + (2\pi i)^m \delta_{\Gamma_{\alpha,\mu}}\}$, where $R_{\alpha,\mu} = R_3|_{\mathcal{X}_{\eta_{T_\mu}}}$, $\partial \Gamma_{\alpha,\mu} = T_3|_{\mathcal{X}_{\eta_{T_\mu}}}$. Over $V_\alpha \cap V_\beta$, these differ by only $(2\pi i)^p \delta_{\Gamma_{\alpha,\mu} - \Gamma_{\beta,\mu}}$ (which are rational cycles); hence we get a Čech cocycle in $H^1(\eta_{\mathcal{M}}, \mathbb{H}/\mathbb{F})$, and this is $\delta \bar{\nu}_3^i$.

Now the fundamental class $[\mathfrak{Z}]$ is given by $(2\pi i)^p T_3$, which (fibered by $\rho_{i-1}^\eta \circ \pi$) gives a trivial section⁶ of $\mathbb{H}_i \cong \mathbb{R}^{-m}(\rho_{i-1}^\eta \circ \pi)_* \mathbf{s}^\bullet \mathcal{W}_{\mathbb{Q}, (\mathcal{X}_{\eta_S}/\eta_{\mathcal{M}_{i-1}})}(p)$ and thus maps to some $[\overline{\mathfrak{Z}}] \in W_0 H^1(\eta_{\mathcal{M}_{i-1}}, \mathbb{H}_i)$. (This is by Leray for $\rho_{i-1}^\eta \circ \pi$; while the relevant s.s. doesn't degenerate at E_2 , the $E_\infty^{1,*}$ terms inject into $E_2^{1,*}$.) This $[\overline{\mathfrak{Z}}]$ is calculated by bounding the fibers $T_3|_{\mathcal{X}_{\eta_{T_\mu}}}$ over the V_α and taking differences over $V_\alpha \cap V_\beta$; reusing the $\{\Gamma_{\alpha,\mu}\}$ shows immediately that the image of $[\overline{\mathfrak{Z}}]$ under $j : W_0 H^1(\eta_{\mathcal{M}_{i-1}}, \mathbb{H}_i) \rightarrow H^1(\eta_{\mathcal{M}_{i-1}}, \mathbb{H}_i/\mathbb{F}_i)$ is $\delta \bar{\nu}_3^i$. Moreover, since $\mathbb{F}_i$ is a locally constant system of (limits of) pure HS of weight 0, $W_0 H^1(\eta_{\mathcal{M}}, \mathbb{F}) = 0$ and j is injective. So from $\delta \bar{\nu}_3^i$ we recover $[\overline{\mathfrak{Z}}]$, and map this to $H^1(\eta_{\mathcal{M}_{i-1}}, R^{2p-m-1}(\rho_{i-1}^\eta \circ \pi)_* \mathbb{Q}(p))$, which has a filtration $\mathbb{L}$ (using Leray for π proper) with graded pieces $Gr_{\mathbb{L}}^a \cong H^1(\eta_{\mathcal{M}_{i-1}}, R^{a-1}(\rho_{i-1}^\eta)_* R^{2p-m-a} \pi_* \mathbb{Q}(p))$. By Leray for ρ_{i-1}^η and vanishing of $[\mathfrak{Z}]_0$ thru $[\mathfrak{Z}]_{i-1}$, $[\overline{\mathfrak{Z}}]$ lives in $\mathbb{L}^i$; its image in $Gr_{\mathbb{L}}^i$ identifies with that of $[\mathfrak{Z}]_i$ under

$$\kappa_i : H^i(\eta_S, R^{2p-m-i} \pi_* \mathbb{Q}(p)) \rightarrow H^1(\eta_{\mathcal{M}_{i-1}}, R^{i-1}(\rho_{i-1}^\eta)_* R^{2p-m-i} \pi_* \mathbb{Q}(p)).$$

(This exists by Leray for ρ_{i-1}^η and the vanishing of $R^i(\rho_{i-1}^\eta)_*(\dots)$ by fiber dimension.) Fix $q_i \in \eta_{\mathcal{M}_{i-1}}(\mathbb{C})$, and let $\mathcal{C}_i := \rho_{i-1,i}(q_i) \subset \mathcal{M}_{i-1}$, $W_i := \rho_i^{-1}(q_i)$ (multi-hyperplane section of $U \subset \mathcal{S}$). By affine weak Lefschetz and dimension, we have

$$\begin{array}{ccc} H^i(\eta_S, R^{2p-m-i} \pi_* \mathbb{Q}(p)) & \xrightarrow{\gamma_i} & H^i(\eta_{W_i}, R^{2p-m-i} \pi_* \mathbb{Q}(p)) \\ \downarrow \kappa_i & & \downarrow \cong \\ H^1(\eta_{\mathcal{M}_{i-1}}, R^{i-1}(\rho_{i-1}^\eta)_* R^{2p-m-i} \pi_* \mathbb{Q}(p)) & \longrightarrow & H^1(\eta_{\mathcal{C}_i}, R^{i-1}(\rho_{i-1}^\eta)_* R^{2p-m-i} \pi_* \mathbb{Q}(p)) \end{array}$$

which shows κ_i injective. This completes the proof of Prop. 4.5(b).

4.3. Comparing filtrations. In fact, we have shown $\bar{\nu}_3^i = 0 \implies [\mathfrak{Z}]_i = 0$, and so we can define a third filtration on $CH^p(X, m)$ by $'\Lambda^0 := \Lambda^0$, $'\Lambda^1 := \Lambda^1$, and

$$' \Lambda^i CH^p(X, m) := \left\{ \mathcal{Z} \in CH_{hom}^p(X, m) \mid \bar{\nu}_{\mathfrak{s}(\mathcal{Z})}^1 = \dots = \bar{\nu}_{\mathfrak{s}(\mathcal{Z})}^{i-1} = 0 \right\}.$$

Evidently $\Lambda^i \subset ' \Lambda^i$; obviously $' \Lambda^\bullet$ depends *a priori* on our choices of hyperplane sections.

Theorem 4.7. $\mathcal{L}^r \subseteq \Lambda^r \subseteq ' \Lambda^r$ ($\forall r$), with equality for $r = 0, \dots, m+2$.

⁶we are still assuming $[\mathfrak{Z}]_0 = \dots = [\mathfrak{Z}]_{i-1} = 0$.

Proof. First some notation. For $\mu \in \eta_{\mathcal{M}_{i-1}}(\mathbb{C})$, let

$$\tilde{H}_\mu := H^{i-1}(\eta_{\mathcal{T}_\mu}, R^{2p-m-i}\pi_*\mathbb{Q}(p));$$

define a local system $\tilde{\mathbb{H}}_i := W_0 R^{i-1}(\rho_{i-1}^\eta)_* R^{2p-m-i}\pi_*\mathbb{Q}(p)$ on $\eta_{\mathcal{M}_{i-1}}$ (with stalks $W_0 \tilde{H}_\mu$), and $\tilde{\mathcal{H}}_i, \tilde{\mathcal{F}}_i, \tilde{\mathbb{F}}_i, \tilde{\mathcal{J}}_i$ by analogy to $\mathcal{H}_i, \mathcal{F}_i, \mathbb{F}_i, \mathcal{J}_i$ above. Set $\tilde{\mathbb{H}}_i^{pol} := W_{-1}\tilde{\mathbb{H}}_i$ and $\tilde{\mathcal{J}}_i^{pol} := \text{im}\{\tilde{\mathcal{H}}_i^{pol} \rightarrow \tilde{\mathcal{J}}_i\}$; the Jacobian sheaves have stalks $(\tilde{\mathcal{J}}_i)_\mu = \text{Ext}_{\text{MHS}}^1(\mathbb{Q}(0), \tilde{H}_\mu)$ and $(\tilde{\mathcal{J}}_i^{pol})_\mu = \text{Ext}_{\text{PMHS}}^1(\mathbb{Q}(0), \tilde{H}_\mu)$. Using the analysis preceding Prop. 4.5, $\tilde{\mathcal{J}}_i^{pol} \hookrightarrow \tilde{\mathcal{J}}_i \hookrightarrow \mathcal{J}_i$.

Now we explain some details in the big commutative diagram (4.4) on the next page. First, $\text{Ext}(\gamma_i)$ is injective for $i \leq m+1$, because γ_i is injective ($\forall i$) and $i \leq m+1 \implies$ weights of $H^{i-1}(\eta_{W_{i-1}}, R^{2p-m-i}\pi_*\mathbb{Q}(p))$ are $\leq 2(i-1) + (2p-m-i) - 2p = i - (m+2) < 0$. Also, $\text{Hom}(\kappa_i)$ is injective because κ_i is (see above). Next, the local system underlying $H^{i-1}(\rho_{i-1}^\eta)_* H^{2p-m-i}\pi_*\mathbb{Q}_{\mathcal{X}_{\eta_S}}(p)$ is smooth (as the limit over $\mathcal{V} \subset \mathcal{M}_{i-1}$ allows for removal of jumps), so the $\text{Ext}_{\text{MHM}(\eta_{\mathcal{M}_{i-1}})}^1$ can be replaced by extensions $\text{Ext}_{\text{VMHS}(\eta_{\mathcal{M}})_{ad}}^1$ of admissible variations of MHS. By M. Saito's description [SaM1, Rmk. 3.7(ii)] of Δ , it is clear that $j' \circ \Delta = \tilde{\delta} \circ j \circ \varepsilon_i$ (the “pentagon” in (4.4) commutes).

More precisely, consider an admissible VMHS on an affine (Zar. op.) $\mathcal{V} \subset \mathcal{M}_{i-1}$ with (smooth) underlying local system $\tilde{\mathbb{H}}$, and an extension $\tilde{\mathbb{H}} \rightarrow \mathbb{E} \rightarrow \mathbb{Q}_{\mathcal{V}}(0)$ with class $e \in \text{Ext}_{\text{VMHS}(\mathcal{V})_{ad}}^1(\mathbb{Q}_{\mathcal{V}}(0), \tilde{\mathbb{H}})$. Associate to this a “normal function” $\epsilon(e) \in H^0(\mathcal{V}, \tilde{\mathcal{J}})$ whose value at $v \in \mathcal{V}$ is the class of $\tilde{\mathbb{H}}_v \rightarrow \mathbb{E}_v \rightarrow \mathbb{Q}(0)$ in $\tilde{\mathcal{J}}_v^{pol} = \text{Ext}_{\text{PMHS}}^1(\mathbb{Q}(0), \tilde{\mathbb{H}}_v)$. To produce $\Delta(e) \in \text{Hom}_{\text{MHS}}(\mathbb{Q}(0), H^1(\mathcal{V}, \tilde{\mathbb{F}}))$, consider a cover $\{V_\alpha\}$ of $\mathcal{V}$ by acyclic analytic balls, and local “liftings” (morphisms of local systems) $\sigma_\alpha : \mathbb{Q}_{\mathcal{V}}(0) \rightarrow \mathbb{E}$; and take differences over the intersections. If $\epsilon(e)$ is zero, i.e. if the pointwise extensions of MHS split, then e splits locally and we may take the σ_α to be morphisms of VMHS; it follows that $\Delta(e)$ factors through $\text{Hom}_{\text{MHS}}(\mathbb{Q}(0), H^1(\mathcal{V}, \tilde{\mathbb{F}}))$, which is zero since $\tilde{\mathbb{F}} (= “Hg(\mathcal{H})”) has pure weight 0 ($\implies H^1(\mathcal{V}, \tilde{\mathbb{F}})$ has weights > 1). In the limit over $\mathcal{V}$ and U (with $\tilde{\mathbb{H}}$ essentially $\tilde{\mathbb{H}}_i$), ϵ gives ε_i ; so we have established ε_i 's injectivity.$

We easily show $\mathcal{L}^r \subseteq {}'\Lambda$ ($\forall r$); since the proof doesn't depend on fibers of ρ_i being hyperplane sections, $\mathcal{L}^r \subseteq \Lambda^r$ follows as well. Assume $\mathcal{L}^i \subseteq {}'\Lambda^i$, and let $\mathcal{Z} \in \mathcal{L}^i CH^p(X, m)$ (and $\mathfrak{Z} := \mathfrak{s}(\mathcal{Z})$). It is easy to see that $(j \circ \varepsilon_i \circ \alpha_i)(c_{\text{MHM}}^{X,i}(\mathcal{Z})) =: \tilde{\nu}_3^i$ maps to $\bar{\nu}_3^i$ under $H^0(\tilde{\mathcal{J}}_i) \hookrightarrow H^0(\mathcal{J}_i)$. If $\mathcal{Z} \in \mathcal{L}^{i+1}$, then $\tilde{\nu}_3^i = 0 \implies \bar{\nu}_3^i = 0 \implies \mathcal{Z} \in {}'\Lambda^{i+1}$, done.

(4.4)

$$\begin{array}{ccccc}
& & \mathcal{L}^i CH^p(X, m) & & \\
& & \downarrow c_{\text{MHM}}^{X, i} & & \\
& & \text{Ext}_{\text{MHM}(\eta_S)}^i(\mathbb{Q}_{\eta_S}(0), H^{2p-m-i}\pi_*\mathbb{Q}_{\mathcal{X}_{\eta_S}}(p)) & \xrightarrow{pr} & \text{Hom}_{(\text{P})\text{MHS}}(\mathbb{Q}(0), H^i(\eta_S, R^{2p-m-i}\pi_*\mathbb{Q}(p))) \\
& \nearrow & \downarrow \alpha_i & & \downarrow \text{Hom}(\kappa_i) \\
& \text{Ext}_{\text{PMHS}}^1(\mathbb{Q}(0), H^{i-1}(\eta_S, R^{2p-m-i}\pi_*\mathbb{Q}(p))) & \hookrightarrow & \text{Ext}_{\text{MHM}(\eta_S)}^i(\mathbb{Q}_{\eta_S}(0), H^{2p-m-i}\pi_*\mathbb{Q}_{\mathcal{X}_{\eta_S}}(p)) & \xrightarrow{\Delta} \text{Hom}_{(\text{P})\text{MHS}}(\mathbb{Q}(0), H^1(\eta_{\mathcal{M}_{i-1}}, R^{i-1}(\rho_{i-1}^\eta)_* R^{2p-m-i}\pi_*\mathbb{Q}(p))) \\
& \downarrow & & \downarrow \varepsilon_i & \downarrow \tilde{j} \\
& \text{Ext}_{\text{PMHS}}^1(\mathbb{Q}(0), H^0(\eta_{\mathcal{M}_{i-1}}, R^{i-1}(\rho_{i-1}^\eta)_* R^{2p-m-i}\pi_*\mathbb{Q}(p))) & \hookrightarrow & \text{Ext}_{\text{MHM}(\eta_{\mathcal{M}_{i-1}})}^1(\mathbb{Q}_{\eta_{\mathcal{M}}}(0), H^{i-1}(\rho_{i-1}^\eta)_* H^{2p-m-i}\pi_*\mathbb{Q}_{\mathcal{X}_{\eta_S}}(p)) & \xrightarrow{\Delta} \text{Hom}_{(\text{P})\text{MHS}}(\mathbb{Q}(0), H^1(\eta_{\mathcal{M}_{i-1}}, R^{i-1}(\rho_{i-1}^\eta)_* R^{2p-m-i}\pi_*\mathbb{Q}(p))) \\
& \downarrow & & \downarrow \text{ev}_{q_{i-1}} & \downarrow \tilde{\delta} \\
& \text{Ext}_{\text{PMHS}}^1(\mathbb{Q}(0), H^{i-1}(\eta_{\mathcal{W}_{i-1}}, R^{2p-m-i}\pi_*\mathbb{Q}(p))) & \xlongequal{\quad} & \text{Ext}_{\text{MHM}(q_{i-1})}^1(\mathbb{Q}(0), H^{i-1}(\rho_{i-1}^\eta)_* H^{2p-m-i}\pi_*\mathbb{Q}_{\mathcal{X}_\eta}(p)) & \xrightarrow{j} H^0(\eta_{\mathcal{M}_{i-1}}, \tilde{\mathcal{J}}_i) \\
& \nearrow \text{Ext}(\gamma_i) & & \nearrow \iota_{q_{i-1}}^* & \uparrow \tilde{\delta} \\
& & & H^0(\eta_{\mathcal{M}_{i-1}}, \tilde{\mathcal{J}}_i^{pol}) & \uparrow \tilde{\delta} \\
& & & \downarrow \varepsilon_i & \uparrow \tilde{\delta} \\
& & & H^1(\eta_{\mathcal{M}_{i-1}}, \tilde{\mathbb{H}}_i/\tilde{\mathbb{F}}_i) & \uparrow \tilde{\delta}
\end{array}$$

We remark that whenever $\bar{\nu}_3^i$ lifts to (or “factors through”) some $\tilde{\nu}_3^i \in H^0(\eta_{\mathcal{M}_{i-1}}, \tilde{\mathcal{J}}_i)$, the topological-invariant map δ lifts to $\tilde{\delta}$ in (4.4). Such a lifting occurs (for any i) when $\mathcal{Z} \in \mathcal{L}^i$ as above, but also (for $i \leq m+2$) if $\mathcal{Z} \in {}'\Lambda^i$ (since Prop. 4.5 is valid for *restricted* higher normal functions as well).

Assume inductively $\mathcal{L}^r = {}'\Lambda^r \ \forall r \leq i$, where $i \leq m+1$. To prove $\mathcal{L}^{i+1} = {}'\Lambda^{i+1}$ it suffices to show ${}'\Lambda^{i+1} \cap \mathcal{L}^i \subseteq \mathcal{L}^{i+1}$. Given $\mathcal{Z} \in {}'\Lambda^{i+1} \cap \mathcal{L}^i$, $(j \circ \varepsilon_i \circ \alpha_i)(c_{\text{MHM}}^{X,i}(\mathcal{Z}))$ is zero since it goes to $\bar{\nu}_3^i$ under an injective map. Hence $(\Delta \circ \alpha_i)(c_{\text{MHM}}^{X,i}(\mathcal{Z})) = 0 \implies pr(c_{\text{MHM}}^{X,i}(\mathcal{Z})) = 0$. Injectivity of $Ext(\gamma_i)$ and an obvious diagram chase show that in fact $c_{\text{MHM}}^{X,i}(\mathcal{Z}) = 0$ and so $\mathcal{Z} \in \mathcal{L}^{i+1}$. $\square$

Remark 4.8. (i) The (weak) analogue here of Zucker’s theorem on normal functions [Z, sec. 9] is the following trivial consequence of the diagram (4.4): the image of $\tilde{\delta} : \{\text{im}(j \circ \varepsilon_i) \subseteq H^0(\eta_{\mathcal{M}_{i-1}}, \tilde{\mathcal{J}}_i)\} \rightarrow H^1(\eta_{\mathcal{M}_{i-1}}, \tilde{\mathbb{H}}_i/\tilde{\mathbb{F}}_i)$ is precisely the [rational] (p, p) classes in the target. (We stress that this analogy is imperfect, as we do not have a concept of horizontality for sections of $\tilde{\mathcal{J}}_i$.)

(ii) For $\mathcal{Z} \in \mathcal{L}^i$ ($\forall i$) or $\mathcal{Z} \in {}'\Lambda^i$ ($i \leq m+2$), the diagram actually gives another proof⁷ that $\bar{\nu}_3^i$ has $[\mathfrak{Z}]_i$ as its topological invariant, since $\tilde{\delta}\tilde{\nu}_3^i = (\tilde{\delta} \circ j \circ \varepsilon_i \circ \alpha_i)(c_{\text{MHM}}^{X,i}(\mathcal{Z})) = (\tilde{j} \circ Hom(\kappa_i) \circ pr)(c_{\text{MHM}}^{X,i}(\mathcal{Z})) = (\tilde{j} \circ Hom(\kappa_i))[\mathfrak{Z}]_i$. This proves the conjecture (“Expectation”) of [Ke2, sec. 7.1].

We finish with some remarks on infinitesimal invariants. Assume our affine $U \subset \mathcal{S}$ is sufficiently small that $\pi : \mathcal{X}_U \rightarrow U$ is smooth. Define $\nabla J^{p,i,m}(\mathcal{X}_U/U)$ to be the cohomology of

$$\begin{aligned} \Gamma(\Omega_U^{i-1} \otimes \mathbb{R}^{2p-m-i} \pi_* \Omega_{\mathcal{X}_U/U}^{\bullet \geq p-i+1}) &\xrightarrow{\nabla} \Gamma(\Omega_U^i \otimes \mathbb{R}^{2p-m-i} \pi_* \Omega_{\mathcal{X}_U/U}^{\bullet \geq p-i}) \\ &\xrightarrow{\nabla} \Gamma(\Omega_U^{i+1} \otimes \mathbb{R}^{2p-m-i} \pi_* \Omega_{\mathcal{X}_U/U}^{\bullet \geq p-i-1}) \end{aligned}$$

at the middle term; M. Saito [SaM3] constructed a map

$$\Phi_{p,i,m} : Hom_{\text{MHS}}(\mathbb{Q}(0), H^i(U, R^{2p-m-i} \pi_* \mathbb{Q}(p))) \rightarrow \nabla J^{p,i,m}(\mathcal{X}_U/U).$$

Passing to the $\varinjlim_U$, the *infinitesimal invariant* of ν_3^i (or $\bar{\nu}_3^i$) is given by

$$d\nu_3^i := \Phi_{p,i,m}(\delta\bar{\nu}_3^i) = \Phi_{p,i,m}([\mathfrak{Z}]_i) \in \nabla J^{p,i,m}(\mathcal{X}_\eta/\eta_{\mathcal{S}}).$$

Remark 4.9. For $m = 0$, “higher normal functions” $\rho_{\mathcal{X}_\eta/\eta_{\mathcal{S}}}^{p,i}(\mathfrak{Z})$ were defined by S. Saito [SaS]. These, too, formally take $\Phi_{p,i,m}([\mathfrak{Z}]_i)$ as their infinitesimal invariants. However, for $i \geq 2$ they are missing the “higher

⁷we have proved this for $\mathcal{Z} \in {}'\Lambda^i$ ($\forall i$).

AJ'' data contained in ν_3^i (and indeed contain no more information than $\Phi_{p,i,m}([\mathfrak{Z}]_i)$ itself).

5. LERAY FILTRATION IN THE PRODUCT CASE

Imagine that the smooth projective “ X ” treated in the discussion of spreads in §4, was actually $X_K = X \otimes_k K$ for some X defined $/k$. The k -spread geometry is then $\pi : \mathcal{X} = X \times \mathcal{S} \rightarrow \mathcal{S}$, where $K \cong k(\mathcal{S})$; if also $\mathcal{Z} \in Z^p(X_K, m)$, then $\mathfrak{Z} \in Z^p((X \times \eta_{\mathcal{S}})_k, m)$. This is the setup for §§5, 6, 7.

5.1. Higher Abel-Jacobi maps. For W smooth projective we let (as in [RS]) $\mathcal{K}_{\mathcal{H}}^{\bullet}(W)$ denote a (canonically chosen) complex of MHS with cohomology computing $H^*(W, \mathbb{Q})$. We have a noncanonical isomorphism

$$\mathcal{K}_{\mathcal{H}}^{\bullet}(W) \simeq \bigoplus_j H^j(W, \mathbb{Q})[-j]$$

in the bounded derived category $D^b MHS$. In particular,

$$\mathcal{K}_{\mathcal{H}}^{\bullet}(X \times \eta_{\mathcal{S}}) \simeq \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_{\mathcal{S}}) \otimes \mathcal{K}_{\mathcal{H}}^{\bullet}(X) \simeq_{n.c.}$$

$$(5.1) \quad \bigoplus_j \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_{\mathcal{S}}) \otimes H^{2p-m-j}(X, \mathbb{Q})[-2p+m+j] \\ \implies H_{\mathcal{H}}^{2p-m}(X \times \eta_{\mathcal{S}}, \mathbb{Q}(p)) \cong \text{Ext}_{D^b MHS}^{2p-m}(\mathbb{Q}(-p), \mathcal{K}_{\mathcal{H}}^{\bullet}(X \times \eta_{\mathcal{S}})) \simeq$$

$$(5.2) \quad \bigoplus_j \text{Ext}_{D^b MHS}^j(\mathbb{Q}(-p), \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_{\mathcal{S}}) \otimes H^{2p-m-j}(X))$$

yields an obvious Leray filtration on $H_{\mathcal{H}}^{2p-m}(X \times \eta_{\mathcal{S}}, \mathbb{Q}(p))$, via $\mathcal{L}^i = \bigoplus_{j \geq i}$. This is canonical, and is easily shown to be strictly compatible with $\mathcal{L}^{\bullet}$ of §4, under the map

$$\begin{array}{ccc} \text{Ext}_{D^b PMHS}^{2p-m}(\mathbb{Q}(0), (a_U)_* \pi_* \mathbb{Q}_{X \times U}(p)) & \hookrightarrow & \text{Ext}_{D^b MHS}^{2p-m}(\mathbb{Q}(0), (a_U)_* \pi_* \mathbb{Q}_{X \times U}(p)) \\ \parallel & & \parallel \\ \text{Ext}_{MHM(X \times U)}^{2p-m}(\mathbb{Q}_{X \times U}(0), \mathbb{Q}_{X \times U}(p)) & & H_{\mathcal{H}}^{2p-m}(X \times U, \mathbb{Q}(p)) \end{array}$$

(after taking $\varinjlim_U$). Simply identify $\mathcal{K}_{\mathcal{H}}^{\bullet}(X \times U)(p)$ with $(a_{X \times U})_* \mathbb{Q}_{X \times U}(p) \simeq (a_U)_* \pi_* \mathbb{Q}_{X \times U}(p)$, and use the filtration of (5.1) by $\bigoplus_{\geq i}$ to produce (compatibly) both (5.2) and (4.2).

Consequently we may reproduce the $\mathcal{L}^{\bullet}$ of §4 on $CH^p(X_K, m)$ (in our present special case) by pulling $\mathcal{L}^{\bullet}$ back along the composite

$$CH^p(X_K, m) \xrightarrow[\mathfrak{s} := k\text{-spread}]{\cong} CH^p((X \times \eta_{\mathcal{S}})_k, m) \xrightarrow[c_{\mathcal{H}}]{} H_{\mathcal{H}}^{2p-m}(X \times \eta_{\mathcal{S}}, \mathbb{Q}(p)).$$

We always denote $\mathfrak{s}\langle \mathcal{Z} \rangle$ by $\langle \mathfrak{Z} \rangle$ (or just $\mathfrak{Z}$). We will be particularly interested in the restriction of $\mathcal{L}^\bullet$ (by intersection) to

$$\underline{CH}^p(X_K, m) := \mathfrak{s}^{-1}(\text{im}\{CH^p((X \times S)_k, m) \rightarrow CH^p((X \times \eta_S)_k, m)\}),$$

the “higher Chow cycles with complete k -spread” $\langle \bar{\mathfrak{Z}} \rangle \in CH^p(X \times S, m)$.

Recall that for smooth quasi-projective Y , $\underline{H}^n(Y)$ for Y denotes $\text{im}\{H^n(\bar{Y}) \rightarrow H^n(Y)\}$ for any good compactification $\bar{Y}$, or equivalently the lowest weight filtrand $W_n H^n(Y)$. (Dually, $H_n(Y) := \text{coim}\{H_n(Y) \rightarrow H_n(\bar{Y})\}$.) Calculating for $m > 0$

$$\begin{aligned} & \text{im} \left\{ \begin{array}{l} \text{Ext}_{D^b MHS}^i(\mathbb{Q}(-p), \mathcal{K}_{\mathcal{H}}^\bullet(\mathcal{S}) \otimes H^{2p-m-i}(X)) \\ \rightarrow \text{Ext}_{D^b MHS}^i(\mathbb{Q}(-p), \mathcal{K}_{\mathcal{H}}^\bullet(\eta_S) \otimes H^{2p-m-i}(X)) \end{array} \right\} \\ & \cong \text{im} \left\{ \begin{array}{l} \text{Ext}_{MHS}^1(\mathbb{Q}(-p), H^{i-1}(\mathcal{S}) \otimes H^{2p-m-i}(X)) \\ \rightarrow \text{Ext}_{MHS}^1(\mathbb{Q}(-p), H^{i-1}(\eta_S) \otimes H^{2p-m-i}(X)) \end{array} \right\} \\ & \cong \text{im} \left\{ \begin{array}{l} \text{Ext}_{MHS}^1(\mathbb{Q}(-p), \underline{H}^{i-1}(\eta_S) \otimes H^{2p-m-i}(X)) \\ \rightarrow \text{Ext}_{MHS}^1(\mathbb{Q}(-p), H^{i-1}(\eta_S) \otimes H^{2p-m-i}(X)) \end{array} \right\} \\ (5.3) \quad & \cong \frac{\text{Ext}_{MHS}^1(\mathbb{Q}(-p), \underline{H}^{i-1}(\eta_S) \otimes H^{2p-m-i}(X))}{\text{im} \left\{ \text{Hom}_{MHS} \left(\mathbb{Q}(-p), \left\{ \frac{W_{m+i}}{W_{i-1}} H^{i-1}(\eta_S) \right\} \otimes H^{2p-m-i}(X) \right) \right\}} \end{aligned}$$

we have our basic invariants (for $m > 0$)

$$\begin{array}{ccc} Gr_{\mathcal{L}}^i \underline{CH}^p(X_K, m) & \xhookrightarrow{\quad \quad \quad} & Gr_{\mathcal{L}}^i CH^p(X_K, m) \ni \mathcal{Z} \\ \downarrow \wr & \nearrow & \\ Gr_{\mathcal{L}}^i c_{\mathcal{H}} & & Gr_{\mathcal{L}}^i c_{\mathcal{H}} \\ \downarrow & \nearrow & \uparrow \beta \\ Gr_{\mathcal{L}}^i H_{\mathcal{H}}^{2p-m}(X \times \eta_S, \mathbb{Q}(p)) & \xhookrightarrow{\quad \quad \quad} & Gr_{\mathcal{L}}^i H_{\mathcal{H}}^{2p-m}(X \times \eta_S, \mathbb{Q}(p)) \\ \uparrow \cong & & \uparrow \alpha \\ \frac{\text{Ext}_{MHS}^1(\mathbb{Q}(-p), \underline{H}^{i-1}(\eta_S) \otimes H^{2p-m-i}(X))}{\text{im} \left[\text{Hom}_{MHS} \left(\mathbb{Q}(-p), \frac{W_{m+i}}{W_{i-1}} H^{i-1}(\eta_S) \otimes H^{2p-m-i}(X) \right) \right]} & \xhookrightarrow{\quad \quad \quad} & \text{Ext}_{MHS}^1(\mathbb{Q}(-p), H^{i-1}(\eta_S) \otimes H^{2p-m-i}(X)) \end{array}$$

sending $\mathcal{Z} \mapsto [c_{\mathcal{H}}(\mathfrak{Z})]_i \xrightarrow{\beta} [\mathfrak{Z}]_i = cl_i(\mathcal{Z})$, and if this is 0 (pulling back

along α) to $[AJ(\mathfrak{Z})]_{i-1} = AJ_{i-1}(\mathcal{Z})$. As in §4, these invariants vanish

for $i > \min\{t+1, p\}$. They are called (resp.) i^{th} higher cycle-class and $(i-1)^{\text{st}}$ higher Abel-Jacobi class (of $\mathcal{Z}$).

We want to say something about computing these; a different approach will be taken in §7. The trouble is that the only datum we have from spreading out is $\mathfrak{Z}_{U_0} \in Z^p(Z \times U_0, m)_{\partial_{\mathcal{B}-\text{cl.}}}$. To produce a double-complex cocycle $\mathfrak{Z}^\bullet \in \mathfrak{s}^{-m}\mathcal{M}_{X \times U_0}(p)$ from this might require an explicit (possibly difficult) application of Bloch's moving lemma. (The one exception to this is the case $m = 0$ of algebraic cycles.) So we will make do without the double-complex *AJ*-formulas of §3.

To compute $[\mathfrak{Z}]_i$, we use the C^∞ chain $\overline{T_{\mathfrak{Z}_{U_0}}}$ (viewed as a topological correspondence on $X \times (\mathcal{S}, D_0)$, where $D_0 := \mathcal{S} \setminus U_0$) to induce a map $H_i(\eta_{\mathcal{S}}, \mathbb{Q}) \rightarrow H_{2d-2p+m+i}(X, \mathbb{Q})$. This map is nonzero if and only if $[\mathfrak{Z}]_i \in \text{Hom}_{MHS}(\mathbb{Q}(-p), H^i(\eta_{\mathcal{S}}) \otimes H^{2p-m-i}(X))$ is.

When $[\mathfrak{Z}]_i$ vanishes, one would like an explicit description of $[AJ(\mathfrak{Z})]_{i-1}$ as a collection of “generalized membrane integrals” — i.e., as functionals on test C^∞ forms. Even if $[\mathfrak{Z}] \in \text{Hom}_{MHS}(\mathbb{Q}(-p), H^{2p-m}(X \times \eta_{\mathcal{S}}, \mathbb{Q}))$ vanishes, this can be computationally horrendous. The simplest description (in general) of $F^{d+t-p+1}\{H^{2t-i+1}((\mathcal{S}, D_0), \mathbb{C}) \otimes H^{2d-2p+m+i}(X, \mathbb{C})\}$ in terms of C^∞ forms uses Hodge-filtered Künneth components of $\mathfrak{s}^\bullet \Omega(p)^{\bullet, \bullet}$ from §3 (with $\mathcal{X} = X \times \mathcal{S}$, $\mathcal{Y} = X \times D_0$). (It is well-known that C^∞ forms with compact support on U_0 are not amenable to filtration by [Hodge] type.)

The one really nice case is when $\mathcal{Z} \in \mathcal{L}^i \underline{CH}^p(X_K, m)$, so that there exists $\bar{\mathfrak{Z}} \in CH^p(X \times \mathcal{S}, m)$ — automatically $\stackrel{\text{hom}}{=} 0$ for $m > 0$ (and we assume for the moment $0 = [\bar{\mathfrak{Z}}] \in \text{Hom}_{MHS}(\mathbb{Q}(-p), H^{2p}(X \times \mathcal{S}))$ if $m = 0$). The [KLM] current $(2\pi i)^{p-m} R_{\bar{\mathfrak{Z}}} \pm (2\pi i)^p \delta_{\partial^{-1}T_{\bar{\mathfrak{Z}}}} =: (2\pi i)^{p-m} R_{\bar{\mathfrak{Z}}}^\pm$ yields a functional on (C^∞ forms representing) $F^{d+t-p+1}\{H^{2t-i+1}(\mathcal{S}, \mathbb{C}) \otimes H^{2d-2p+m+i}(X, \mathbb{C})\}$, modulo periods. Now since

$$\begin{aligned} & \text{Hom}_{MHS} \left(\mathbb{Q}(-p), \frac{W_{m+i}}{W_{i-1}} H^{i-1}(\eta_{\mathcal{S}}) \otimes H^{2p-m-i}(X) \right) \\ & \xleftarrow{\cong} \text{Hom}_{MHS} \left(\mathbb{Q}(-p), \frac{W_{m+i}}{W_{i-1}} H^{i-1}(\eta_{\mathcal{S}}) \otimes F_h^{p-i+1} H^{2p-m-i}(X) \right), \end{aligned}$$

we can kill the “ $\text{im}\{Hom\}$ ” in (5.3) by projecting

$$Gr_{\mathcal{L}}^i \underline{H_{\mathcal{H}}^{2p-m}}(X \times \eta_{\mathcal{S}}, \mathbb{Q}(p)) \twoheadrightarrow Ext_{MHS}^1 \left(\mathbb{Q}(-p), \underline{H^{i-1}}(\eta_{\mathcal{S}}) \otimes \frac{H^{2p-m-i}(X)}{F_h^{p-i+1} H^{2p-m-i}(X)} \right);$$

the image of $[AJ(\mathfrak{Z})]_{i-1}$ by this is denoted $[AJ(\mathfrak{Z})]_{i-1}^{tr} = AJ_{i-1}^{tr}(\mathcal{Z})$. The functional described above (an element of $Ext_{MHS}^1(\mathbb{Q}(-p), H^{i-1}(\mathcal{S}) \otimes H^{2p-m-i}(X))$) projects to compute this $[AJ(\mathfrak{Z})]_{i-1}^{tr}$.

In short, we have constructed a slight “quotient” of AJ_{i-1} ,

$$AJ_{i-1}^{tr} : \mathcal{L}^i \underline{CH}^p(X_K, m) \rightarrow Ext_{\text{MHS}}^1 \left(\mathbb{Q}(-p), \underline{H}^{i-1}(\eta_S) \otimes \frac{H^{2p-m-i}(X)}{F_h^{p-i+1}} \right),$$

that can be described explicitly in the spirit of the “Griffiths prescription”; it will be used in §6.

5.2. Weight filtration redux. Our next aim is to Leray-filter the $\mathcal{H}(p)$ -spectral sequence and the residue maps from §3, when $\mathcal{X} = X \times \mathcal{S}$. Recall from (5.2) that for W smooth quasiprojective

$$\begin{aligned} H_{\mathcal{H}}^M(X \times W, \mathbb{Q}(q)) &\cong \bigoplus_{\nu} Ext_{D^b \text{MHS}}^{M-\nu}(\mathbb{Q}(-\nu), \mathcal{K}_{\mathcal{H}}^{\bullet}(W) \otimes H^{\nu}(X)) \\ &=: \bigoplus_{\nu} H_{\mathcal{H}}^{M-\nu}(W, H^{\nu}(X, \mathbb{Q}(q))) =: \bigoplus_{\nu} Gr_{\nu}^{\mathcal{L}} H_{\mathcal{H}}^M(X \times W, \mathbb{Q}(q)) \\ &\cong \bigoplus_{\nu} Gr_{\mathcal{L}}^{M-\nu} H_{\mathcal{H}}^M(X \times W, \mathbb{Q}(q)). \end{aligned}$$

If W is projective, there are obvious cochain complexes

$$Gr_{\nu}^{\mathcal{L}} C_{\mathcal{H}}^{\bullet}(X \times W, \mathbb{Q}(q))$$

for computing the graded pieces, with $\bigoplus_{\nu} \simeq$ to the usual $C_{\mathcal{H}}^{\bullet}$. For $\bullet < 2q$ the $Gr_{\nu}^{\mathcal{L}}$ complex is given by

$$\begin{aligned} C_{top}^{\bullet-\nu}(W, H^{\nu}(X, \mathbb{Q}(q))) \oplus F^q(D^{\bullet-\nu}(W) \otimes H^{\nu}(X, \mathbb{C})) \oplus \\ (D^{\bullet-\nu-1}(W) \otimes H^{\nu}(X, \mathbb{C})), \end{aligned}$$

for $\bullet > 2q + 1$ it is 0, and the reader can fill in the rest.

Passing to the spread situation $\mathcal{X} = X \times \mathcal{S}$, $\mathcal{Y}^i = X \times D^i$ for $D \subset \mathcal{S}$ NCD, we form double complexes

$$Gr_{\nu}^{\mathcal{L}} \mathcal{H}_{\mathcal{X} \setminus \mathcal{Y}}(p)_0^{a,b} := Gr_{\nu}^{\mathcal{L}} C_{\mathcal{H}}^{2p+2a+b}(X \times \widetilde{D^{-a}}, \mathbb{Q}(p+a))$$

with $s^{\bullet} \mathcal{H}(p)_0 \simeq \bigoplus_{\nu} s^{\bullet} Gr_{\nu}^{\mathcal{L}} \mathcal{H}(p)_0$; in fact

$$\mathcal{H}(p)_1^{a,b} \cong \bigoplus_{\nu} H_{\mathcal{H}}^{2p+2a+b-\nu}(\widetilde{D^{-a}}, H^{\nu}(X, \mathbb{Q}(p+a))) \cong \bigoplus_{\nu} Gr_{\nu}^{\mathcal{L}} \mathcal{H}(p)_1^{a,b},$$

and the $\cong$ persists for all pages $r \geq 1$. Hence the $Gr_{\nu}^{\mathcal{L}}$ spectral sequences converge to $Gr_{\nu}^{\mathcal{L}} H_{\mathcal{H}}^{2p+*}(X \times (\mathcal{S} \setminus D), \mathbb{Q}(p))$, and as before we get weight filtrations with maps

$$\begin{aligned} Gr_b^W Gr_{\nu}^{\mathcal{L}} H_{\mathcal{H}}^{2p+a+b}(X \times (\mathcal{S} \setminus D), \mathbb{Q}(p)) &\cong Gr_{\nu}^{\mathcal{L}} \mathcal{H}(p)_{\infty}^{a,b} \\ &\xrightarrow{Res^{-a}} Gr_{\nu}^{\mathcal{L}} \underline{H_{\mathcal{H}}^{2p+2a+b}}(X \times (D^{-a} \setminus D^{-a+1}), \mathbb{Q}(p+a)). \end{aligned}$$

We proceed to the main application.

With the understanding that the limit (over U affine Zariski open in $\mathcal{S}$) defining $\eta_{\mathcal{S}}$ is taken together with a system of good compactifications (or of $(\tilde{\mathcal{S}}, \tilde{D})$ with $\tilde{D}$ a NCD), we can define⁸

$$W_{\ell}Gr_{\mathcal{L}}^iCH^p(X_K, m) := \mathfrak{s}^{-1} \left(W_{\ell}Gr_{\mathcal{L}}^iCH^p((X \times \eta_{\mathcal{S}})_L, m) \right).$$

The following is a simple means for constructing nontrivial m -decomposable higher cycles in the weight-graded pieces.

Theorem 5.1. *Let $\tau_1, \dots, \tau_m \in \mathbb{C}$ be algebraically independent $/\bar{K}$, and set $L := K(\tau_1, \dots, \tau_m)$. The exterior product with $\{\tau_1, \dots, \tau_m\} \in CH^m(\text{Spec}L, m)$ gives a map*

$$Gr_{\mathcal{L}}^i\overline{CH}^p(X_K, n) \hookrightarrow Gr_{-n}^W Gr_{\mathcal{L}}^{i+m}CH^{p+m}(X_L, n+m).$$

Proof. L is the compositum of $L_0 := k(\tau_1, \dots, \tau_m)$ and $K \cong k(\mathcal{S})$; writing $\mathcal{T} := \mathbb{P}_{t_1}^1 \times \dots \times \mathbb{P}_{t_m}^1$ we have $L_0 \cong k(\mathcal{T})$, $L \cong k(\mathcal{S} \times \mathcal{T})$. Taking D, E NCD's in $\mathcal{S}, \mathcal{T}$ (resp.), with $E \supseteq \cup_j |(t_j)|$, set $\mathfrak{D} := (\mathcal{S} \times E) \cup (D \times \mathcal{T})$. The k -spread of the exterior product $\mathcal{Z} \times \{\underline{\tau}\}$ is the exterior product of

$$\mathfrak{Z} \in \mathcal{L}^i\overline{CH}^p(X \times (\mathcal{S} \setminus D), n) \quad \text{and} \quad \{\underline{\tau}\} \in [\mathcal{L}^m]CH^m([\text{pt.} \times] \mathcal{T} \setminus E, m).$$

Assuming $\mathcal{Z}$ is nonzero in $Gr_{\mathcal{L}}^i\overline{CH}^p(X_K, n)$, the class of $\mathfrak{Z}$ is nontrivial (in the limit over all D) in

$$(5.4) \quad Gr_{\mathcal{L}}^i\overline{CH}^p(X \times (\mathcal{S} \setminus D), n) \xrightarrow{Gr_{\mathcal{L}}^i c_{\mathcal{H}}} \underline{H}_{\mathcal{H}}^i(\mathcal{S} \setminus D, H^{2p-n-i}(X, \mathbb{Q}(p)))$$

Evidently the restriction of $Res^m(c_{\mathcal{H}}\{\underline{\tau}\}) \in H_{\mathcal{H}}^0(E^m = \Pi\text{pts.}, \mathbb{Q}(0)) \cong \oplus \mathbb{Q}$ to $p \in \cap_j |(t_j)|$ is ± 1 ($\neq 0$, which puts $\{\underline{\tau}\}$ in the highest weight W_0). Since $\mathfrak{Z}$ is Res-free (hence in W_{-n}), the product gives a class in

$$Gr_{-n}^W Gr_{\mathcal{L}}^{i+m}CH^{p+m}(X \times \{(\mathcal{S} \times \mathcal{T}) \setminus \mathfrak{D}\}, n+m)$$

$$\hookrightarrow Gr_{-n}^W Gr_{\mathcal{L}}^{i+m}H_{\mathcal{H}}^{2p+m-n}(X \times \{(\mathcal{S} \times \mathcal{T}) \setminus \mathfrak{D}\}, \mathbb{Q}(p+m))$$

$$\xrightarrow{Res^m} Gr_{\mathcal{L}}^i \underline{H}_{\mathcal{H}}^{2p-n}(X \times \mathfrak{D}^m \setminus \mathfrak{D}^{m+1}, \mathbb{Q}(p)) \xrightarrow{|(\mathcal{S} \setminus D) \times \{p\}|} \{\text{r.h.s. of (5.4)}\}.$$

Clearly $Res_{(\mathcal{S} \setminus D) \times \{p\}}^m \{Gr_{\mathcal{L}}^{i+m}c_{\mathcal{H}}(\mathfrak{Z} \times \{\underline{\tau}\})\} = Gr_{\mathcal{L}}^i c_{\mathcal{H}}(\mathfrak{Z}) \otimes Res_{\{p\}}^m(c_{\mathcal{H}}\{\underline{\tau}\}) \neq 0$; and it only remains to be said that this does not vanish upon enlarging $\mathfrak{D}$ provided we enlarge D so that $D \times \{p\} \supseteq (\mathcal{S} \times \{p\}) \cap \mathfrak{D}$. $\square$

⁸Warning: one can also define W_* directly on $CH^p(X_K, m)$, and $Gr_{\mathcal{L}}$ and Gr^W do not “commute”. Saying the $Gr_{\mathcal{L}}$ of a class is in W_{ℓ} does not mean the whole class is in W_{ℓ} .

6. DETECTING INDECOMPOSABLES IN THE REGULATOR KERNEL

The j -decomposable cycles are those in the image of the exterior product

$$\theta^j : CH^{p-j}(X_K, m-j) \otimes CH^j(\text{Spec} K, j) \rightarrow \mathcal{L}^j CH^p(X_K, m).$$

For any subgroup $\Xi \subseteq CH^p(X_K, m)$, define

$$\Xi_{j\text{-ind}} := \frac{\Xi}{im(\theta^j) \cap \Xi}.$$

The usual definition of indecomposable is $\Xi_{m\text{-ind}}$, while the “strong indecomposable” notion of [CF] corresponds to $\Xi_{1\text{-ind}}$. We study here the case $\Xi = \mathcal{L}^i \underline{CH}^p(X_K, m)$ — these are cycles with complete spread; that is, their spreads have no residues along substrata of $\mathcal{S}$.

6.1. An invariant for cycles with complete spread. Our aim is to produce, starting from the higher $\{AJ_{i-1}^{tr}\}$ maps of §5.1, further quotients which detect indecomposables. The principal difficulty is that an element of $im(\theta^j) \cap \Xi$ (decomposables with complete spread) may involve terms whose spreads have residues (although the sum of these residues must vanish).

Theorem 6.1. (i) For $1 \leq j \leq m \leq i \leq p$, $Gr_{\mathcal{L}^i \underline{CH}}^i$ (composed with a projection) descends to a map

$$\mathcal{L}^i \underline{CH}_{j\text{-ind}}^p(X_K, m) \rightarrow Ext_{MHS}^1 \left(\mathbb{Q}(-p), \frac{H^{2p-m-i}(X)}{F_h^{p-i}} \otimes \underline{H}^{i-1}(\eta_S) \right).$$

(ii) If $1 \leq j$ and $i \leq p$, but one of the other inequalities fails, then F_h^{p-i} may be replaced by F_h^{p-i+1} (i.e., detecting indecomposables is a trivial matter). If $j = 0$ or $i > p$, there is nothing to detect.

Proof. From §1 we know the map from $\mathcal{L}^i \underline{CH}^p(X_K, m)$ to $Ext_{MHS}^1 \left(\mathbb{Q}(-p), \frac{H^{2p-m-i}(X)}{F_h^{p-i+1}} \otimes \underline{H}^{i-1}(\eta_S) \right)$ is well-defined. So given a class ξ in $\mathcal{L}^i \underline{H}_{\mathcal{H}}^{2p-m}(X \times \eta_S, \mathbb{Q}(p))$ and in the image of

$$(6.1) \quad \begin{aligned} & H_{\mathcal{H}}^{2p-m-j}(X \times \eta_S, \mathbb{Q}(p-j)) \otimes H_{\mathcal{H}}^j(\eta_S, \mathbb{Q}(j)) \longrightarrow \\ & H_{\mathcal{H}}^{2p-m}(X \times \eta_S, \mathbb{Q}(p)), \end{aligned}$$

we need to show its projection $\bar{\xi}$ to

$$Gr_{\mathcal{L}^i \underline{H}_{\mathcal{H}}}^i H_{\mathcal{H}}^{2p-m} \cong Ext_{MHS}^1 \left(\mathbb{Q}(-p), H^{2p-m-i}(X) \otimes H^{i-1}(\eta_S) \right)$$

is contained in the image of

$$(6.2) \quad \begin{aligned} & Ext_{MHS}^1(\mathbb{Q}(-p), F_h^{p-i} H^{2p-m-i}(X) \otimes \underline{H}^{i-1}(\eta_S) \longrightarrow \\ & Ext_{MHS}^1(\mathbb{Q}(-p), H^{2p-m-i}(X) \otimes H^{i-1}(\eta_S)). \end{aligned}$$

Since (6.1) *splits* into a $\oplus$ of Leray-graded pieces, we have immediately that $\bar{\xi} \in$

$$(6.3) \quad \begin{aligned} & Gr_{\mathcal{L}}^i \underline{H}_{\mathcal{H}}^{2p-m}(X \times \eta_S, \mathbb{Q}(p)) \cap \\ & im \left(\begin{array}{c} Gr_{\mathcal{L}}^{i-j} H_{\mathcal{H}}^{2p-m-j}(X \times \eta_S, \mathbb{Q}(p)) \otimes H_{\mathcal{H}}^j(\eta_S, \mathbb{Q}(j)) \\ \rightarrow Gr_{\mathcal{L}}^i H_{\mathcal{H}}^{2p-m}(X \times \eta_S, \mathbb{Q}(p)) \end{array} \right) = \\ & im \left(\begin{array}{c} Ext_{D^b MHS}^i(\mathbb{Q}(-p), H^{2p-m-i}(X) \otimes \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)) \xrightarrow{e_{\mathcal{H}}^j} \\ Ext_{D^b MHS}^i(\mathbb{Q}(-p), H^{2p-m-i}(X) \otimes \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)) \end{array} \right) \cap \\ & im \left(\begin{array}{c} Ext_{D^b MHS}^{i-j}(\mathbb{Q}(-p+j), H^{2p-m-i}(X) \otimes \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)) \otimes Ext_{D^b MHS}^j(\mathbb{Q}(-j), \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)) \\ \xrightarrow{\theta_{\mathcal{H}}^j} Ext_{D^b MHS}^i(\mathbb{Q}(-p), H^{2p-m-i}(X) \otimes \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)) \end{array} \right). \end{aligned}$$

Here we simply take

$$\mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S) := \bigoplus_{\ell} \underline{H}^{\ell}(\eta_S, \mathbb{Q})[-\ell], \quad \underline{\mathcal{K}}_{\mathcal{H}}^{\bullet}(\eta_S) := \bigoplus_{\ell} \underline{H}^{\ell}(\eta_S, \mathbb{Q})[-\ell].$$

Moreover, using an idea of [RS], $Hom_{MHS}(\mathbb{Q}(-j), H^j(\eta_S))$ gives a sub-MHS

$$H^j(\eta_S)'' \subseteq H^j(\eta_S)$$

with $H^j(\eta_S)'' \cap W_{2j-1} H^j(\eta_S) = \{0\}$. By semisimplicity, writing

$\eta : H^j(\eta_S)'' \rightarrow Gr_{2j}^W H^j(\eta_S)$, we have $Gr_{2j}^W H^j(\eta_S) \stackrel{HS}{=} \mathcal{H}' \oplus im(\eta)$. Pulling back the extension $0 \rightarrow W_{2j-1} H^j(\eta_S) \rightarrow H^j(\eta_S) \rightarrow Gr_{2j}^W \rightarrow 0$

along $\mathcal{H}' \hookrightarrow Gr_{2j}^W$ yields $H^j(\eta_S)' \hookrightarrow H^j(\eta_S)$ with $H^j(\eta_S) \stackrel{MHS}{=} H^j(\eta_S)' \oplus H^j(\eta_S)''$. Define more generally

$$H^{\ell}(\eta_S)'' := im \left\{ H^j(\eta_S)'' \otimes H^{\ell-j}(\eta_S) \rightarrow H^{\ell}(\eta_S) \right\},$$

$\mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)'' := \oplus_{\ell} H^{\ell}(\eta_S)''[-\ell]$. A generalization of the above argument (involving both projections and pullbacks of extensions) produces $(\forall \ell)$ $\oplus$ -decompositions of $H^{\ell}(\eta_S)$, hence $\mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S) = \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)' \oplus \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)''$. This induces a $\oplus$ -decomposition on relevant Ext groups (hence also projections and inclusions of Ext 's). Note that $H^{\ell}(\eta_S)' \supseteq W_{\ell+j-1}H^{\ell}(\eta_S) \supseteq W_{\ell}H^{\ell}(\eta_S) = \underline{H}^{\ell}(\eta_S)$.

Since $\mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)' \supseteq \underline{\mathcal{K}}_{\mathcal{H}}^{\bullet}(\eta_S)$, and the projection $\mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S) \twoheadrightarrow \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)'$ is compatible with $\cup$ -products, (6.3) reduces to

$$(6.4) \quad \begin{aligned} & \text{im} \left(\frac{Ext_{D^b MHS}^i(\mathbb{Q}(-p), H^{2p-m-i}(X) \otimes \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)) \xrightarrow{(\epsilon_{\mathcal{H}}^j)'}}{Ext_{D^b MHS}^i(\mathbb{Q}(-p), H^{2p-m-i}(X) \otimes \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)')} \right) \cap \\ & \text{im} \left(\frac{Ext_{D^b MHS}^{i-j}(\mathbb{Q}(-p+j), H^{2p-m-i}(X) \otimes \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)') \otimes Ext_{D^b MHS}^j(\mathbb{Q}(-j), \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)')}{\xrightarrow{(\theta_{\mathcal{H}}^j)'}} Ext_{D^b MHS}^i(\mathbb{Q}(-p), H^{2p-m-i}(X) \otimes \mathcal{K}_{\mathcal{H}}^{\bullet}(\eta_S)') \right). \end{aligned}$$

By construction, $Hom_{MHS}(\mathbb{Q}(-j), H^j(\eta_S)') = 0$. Hence $(\theta_{\mathcal{H}}^j)'$ reduces to

$$(6.5) \quad \begin{aligned} & Hom_{MHS}(\mathbb{Q}(-p+j), H^{2p-m-i}(X) \otimes H^{i-j}(\eta_S)') \otimes Ext_{MHS}^1(\mathbb{Q}(-j), H^{j-1}(\eta_S)) \\ & \longrightarrow Ext_{MHS}^1(\mathbb{Q}(-p), H^{2p-m-i}(X) \otimes H^{i-1}(\eta_S)'). \end{aligned}$$

One easily shows that (in (6.5)) the $Hom = Hom_{MHS}(\mathbb{Q}(-p+j), F_h^{p-i}H^{2p-m-i}(X) \otimes H^{i-j}(\eta_S)').$ (This is straightforward and has nothing to do with $H^{i-j}(\eta_S)$ vs. $H^{i-j}(\eta_S)'$.) Hence (6.5) factors through

$$\begin{aligned} & Ext_{MHS}^1(\mathbb{Q}(-p), F_h^{p-i}H^{2p-m-i}(X) \otimes H^{i-1}(\eta_S)') \\ & \hookrightarrow Ext_{MHS}^1(\mathbb{Q}(-p), H^{2p-m-i}(X) \otimes H^{i-1}(\eta_S)') \end{aligned}$$

and so (6.4) is contained in

$$\begin{aligned} & \text{im} \left(\frac{Ext_{MHS}^1(\mathbb{Q}(-p), H^{2p-m-i}(X) \otimes \underline{H}^{i-1}(\eta_S))}{\rightarrow Ext_{MHS}^1(\mathbb{Q}(-p), H^{2p-m-i}(X) \otimes H^{i-1}(\eta_S)')} \right) \\ & \cap Ext_{MHS}^1(\mathbb{Q}(-p), F_h^{p-i}H^{2p-m-i}(X) \otimes H^{i-1}(\eta_S)') \end{aligned}$$

Using semisimplicity to write $H^{2p-m-i}(X) \cong F_h^{p-i} \oplus \frac{H^{2p-m-i}(X)}{F_h^{p-i}}$ as HS, we see that the above lies in

$$im \left\{ \begin{array}{l} Ext_{MHS}^1(\mathbb{Q}(-p), F_h^{p-i} H^{2p-m-i}(X) \otimes \underline{H^{i-1}}(\eta_S)) \\ \rightarrow Ext_{MHS}^1(\mathbb{Q}(-p), F_h^{p-i} H^{2p-m-i}(X) \otimes H^{i-1}(\eta_S)') \end{array} \right\},$$

which obviously $\subseteq im$ (6.2).

(ii) If $j > i$, the j -decomposables lie in $\mathcal{L}^{i+1}$; if $j > m$, there aren't any; and if $m > i$, the Hom in (6.5) vanishes. $\square$

Setting $i = j := 1$ and $p := q$ in Theorem 6.1, we have for $m \neq 1$ (using (ii))

$$AJ_{\text{ind}} : \mathcal{L}^1 \underline{CH}_{1\text{-ind}}^q(X_k, m) \rightarrow Ext_{MHS}^1(\mathbb{Q}(-q), H^{2p-m-1}(X)) = J^{p,m}(X)$$

(since $F_h^p H^{2p-m-1}(X) = \{0\}$) and for $m = 1$ (using (i))

$$AJ_{\text{ind}} : \mathcal{L}^1 \underline{CH}_{1\text{-ind}}^q(X_k, 1) \rightarrow Ext_{MHS}^1\left(\mathbb{Q}(-q), \frac{H^{2p-2}(X)}{Hg^{p-1}(X)}\right)$$

(since $F_h^{p-1} H^{2p-2}(X) \cong Hg^{p-1}(X)$). It seems natural to call a cycle $\mathcal{W} \in CH_{\text{hom}}^q(X_k, m)$ with $AJ_{\text{ind}}(\mathcal{W}) \neq 0$ “strongly AJ -indecomposable”, but evidently (from the form of AJ_{ind} in the 2 cases) this is the same thing as “regulator-indecomposable” (as in [RS], [CF], [Le2], etc.), so we will use that terminology.

6.2. First application: exterior products. The real point of Theorem 6.1, however, is to detect indecomposable higher cycles in the kernel of AJ . Here is a simple application to classes in the image of the exterior product

$$\mathcal{L}^i CH^{d_1}((Y_1)_K) \otimes \mathcal{L}^1 CH^{p-d_1}((Y_2)_k, m) \rightarrow \mathcal{L}^{i+1} \underline{CH}^p((Y_1 \times Y_2)_K, m),$$

where K/k is finitely generated of transcendence degree i , and Y_1, Y_2 are smooth projective $/k$ of resp. dimensions d_1, d_2 .

Theorem 6.2. *Let $\mathcal{W} \in CH_{\text{hom}}^{p-d_1}((Y_2)_k, m)$ be regulator-indecomposable, and $\mathcal{V} \in \mathcal{L}^i CH^{d_1}((Y_1)_K)$ a 0-cycle whose complete k -spread $\bar{\mathfrak{V}} \in Z^{d_1}((\mathcal{S} \times Y_1)_k)$ induces a nontrivial map $\Omega^i(Y_1) \rightarrow \Omega^i(\mathcal{S})$ of holomorphic i -forms. (See [Ke3] for how to construct such 0-cycles.)*

Then $\mathcal{Z} := \mathcal{V} \times \mathcal{W} \in CH^p((Y_1 \times Y_2)_K, m)$ is strongly indecomposable.

Proof. This is so similar to the proof of Thm. 1 in [Ke3] that we merely indicate briefly how $\mathcal{Z}$ is detected and the technical lemma on HS's needed.

Applying the map of Thm. 6.1 in the appropriate case (substituting the present $i+1$ for i and 1 for j) sends $\mathcal{Z}$ to a quotient (in general)

of $[AJ(\mathfrak{Z})]_i^{tr}$. Since $\mathfrak{Z}$ has complete spread $\bar{\mathfrak{Z}} := \mathcal{W} \times \bar{\mathfrak{W}}$, this can be computed (as in §1) using $(2\pi i)^{p-m} R'_3$ (as a functional on forms). Now $T_{\bar{\mathfrak{Z}}} = T_{\mathcal{W}} \times \bar{\mathfrak{W}} \implies \partial^{-1} T_{\bar{\mathfrak{Z}}} := (\partial^{-1} T_{\mathcal{W}}) \times \bar{\mathfrak{W}}$, and $R_{\bar{\mathfrak{Z}}} = R_{\mathcal{W}} \cdot (2\pi i)^{d_1} \delta_{\bar{\mathfrak{W}}}$; so $\frac{1}{(2\pi i)^{d_1}} R'_3 = R'_{\mathcal{W}} \cdot \delta_{\bar{\mathfrak{W}}}$, and (the quotient of) $[AJ(\mathfrak{Z})]_i^{tr}$ may be computed by the image of $[\bar{\mathfrak{W}}]_i \otimes AJ_{(\text{ind})}(\mathcal{W})$. (We are assuming $AJ_{\text{ind}}(\mathcal{W}) \neq 0$.)

Define sub-Hodge structures

$$\mathcal{H}_1 := H^{2d_1-i}(Y_1) \otimes H^{2p-2d_1-m-1}(Y_2) \otimes H^i(\mathcal{S})$$

$$\mathcal{G}_0 := H^{2p-i-m-1}(Y_1 \times Y_2) \otimes N^1 H^i(\mathcal{S}) + F_h^{p-i-1} H^{2p-i-m-1}(Y_1 \times Y_2) \otimes H^i(\mathcal{S})$$

of $\mathcal{H}_0 := H^{2p-i-m-1}(Y_1 \times Y_2) \otimes H^i(\mathcal{S})$, and

$$\mathcal{H}_{\bar{\mathfrak{W}}} := \mathbb{Q} \langle [\bar{\mathfrak{W}}]_i \rangle \otimes H^{2p-2d_1-m-1}(Y_2)$$

$$\mathcal{G}_1 := H^{2d_1-i}(Y_1) \otimes F_h^{p-d_1-1} H^{2p-2d_1-m-1}(Y_2) \otimes H^i(\mathcal{S})$$

sub-HS of $\mathcal{H}_1$. From the last paragraph,

(6.6)

$$[\bar{\mathfrak{W}}]_i \otimes AJ_{\text{ind}}(\mathcal{W}) \in \mathbb{Q} [\bar{\mathfrak{W}}]_i \otimes J_{\text{ind}}^{p,m}(Y_2) \cong Ext_{MHS}^1 \left(\mathbb{Q}(-p), \frac{\mathcal{H}_{\bar{\mathfrak{W}}}}{\mathcal{G}_1 \cap \mathcal{H}_{\bar{\mathfrak{W}}}} \right)$$

and the Thm. 6.1 quotient of $[AJ(\mathfrak{Z})]_i^{tr}$ in $Ext_{MHS}^1(\mathbb{Q}(-p), \mathcal{H}_0/\mathcal{G}_0)$, map to the same element of

$$(6.7) \quad \frac{Ext_{MHS}^1(\mathbb{Q}(-p), \mathcal{H}_1/\mathcal{G}_1)}{Ext_{MHS}^1(\mathbb{Q}(-p), (\mathcal{H}_1 \cap \mathcal{G}_0)/(\mathcal{G}_1 \cap \mathcal{G}_0))}.$$

One shows that

$$\mathcal{H}_{\bar{\mathfrak{W}}} \cap (\mathcal{H}_1 \cap \mathcal{G}_0) \subseteq \mathcal{G}_1,$$

so that the map from (6.6) to (6.7) is an injection and we are done. $\square$

6.3. Relation to a criterion of Lewis. Given $\mathcal{S}'$ smooth projective $/k$ and $\alpha \in H^{\dim(\mathcal{S}')-i, \dim(\mathcal{S}')}(\mathcal{S}', \mathbb{C})$, let $cl_{\mathcal{S}', \alpha}^{\{p-m, i-m, 0\}}$ denote the composition

$$\begin{array}{ccc} CH^{p-i}(X \times \mathcal{S}') & \xrightarrow{cl} & H^{2p-2i}(X \times \mathcal{S}', \mathbb{C}) \xrightarrow[\text{K\"unneth}]{\text{Hodge}} H^{p-i, p-m}(X, \mathbb{C}) \otimes H^{i, 0}(\mathcal{S}', \mathbb{C}) \\ & & \downarrow \cup \alpha \\ & & H^{p-i, p-m}(X, \mathbb{C}). \end{array}$$

The next application (Theorem 6.3 below) strengthens and generalizes a result of Collino and Fakhruddin [CF]. To put this in context, we first explain how Theorem 6.1 essentially subsumes the invariant

they use for indecomposables in $\ker(AJ)$, due to [Le2]. This may be expressed as a map

$$(6.8) \quad \mathcal{L}^i \underline{CH}_{m\text{-ind}}^p(X_K, m) \rightarrow \frac{H^{p-i, p-m}(X, \mathbb{C})}{H^{\{p-m, i-m, 0\}}(X)} \otimes H^{i-1, 0}(\mathcal{S}, \mathbb{C})$$

where as usual $k(\mathcal{S}) \cong K$, $m \geq 1$, and

$$H^{\{p-m, i-m, 0\}}(X) := \sum_{S', \alpha} \text{im} \left(cl_{S', \alpha}^{\{p-m, i-m, 0\}} \right).$$

To describe (6.8), let H and G be quotient Hodge structures (resp.) of $H^{2p-m-i}(X, \mathbb{Q})$ and $H^{i-1}(\mathcal{S}, \mathbb{Q})$, such that $G_{\mathbb{C}}^{i-1, 0} \cong H^{i-1, 0}(\mathcal{S}, \mathbb{C})$; let $\mathcal{H} := H \otimes G$ and write $pr_{\mathcal{H}} : H^{2p-m-1}(X \times \mathcal{S}, \mathbb{Q}) \twoheadrightarrow \mathcal{H}$. Consider the composition $\mathcal{R}_{\mathcal{H}}^{\{p, i, m\}}$:

$$\begin{array}{ccc} Ext_{\text{MHS}}^1(\mathbb{Q}(-p), \mathcal{H}) & \xleftarrow{pr_{\mathcal{H}} \circ AJ} & CH^p(X \times \mathcal{S}, m) \\ \downarrow & & \\ Ext_{\mathbb{R}\text{-MHS}}^1(\mathbb{R}(-p), \mathcal{H}_{\mathbb{R}}) & \xlongequal{\quad} & \mathcal{H}_{\mathbb{R}(p-1)} \cap \{\mathcal{H}_{\mathbb{C}}^{p-m, p-1} \oplus \dots \oplus \mathcal{H}_{\mathbb{C}}^{p-1, p-m}\} \\ & & \downarrow \\ & & H_{\mathbb{C}}^{p-i, p-m} \otimes H^{i-1, 0}(\mathcal{S}, \mathbb{C}) \xleftarrow{\quad} \mathcal{H}_{\mathbb{C}}^{p-1, p-m} \end{array}$$

Let $\mathcal{H}_1 := H^{2p-m-i}(X) \otimes H^{i-1}(\mathcal{S})$, $\mathcal{H}_2 := \frac{H^{2p-m-i}(X)}{F_h^{p-i}} \otimes \underline{H}^{i-1}(\eta_{\mathcal{S}})$ be choices of $\mathcal{H}$ as above, and consider the commutative diagram

$$\begin{array}{ccc} CH^p(X \times \mathcal{S}, m) & \xrightarrow{\quad} & H^{p-i, p-m}(X, \mathbb{C}) \otimes H^{i-1, 0}(\mathcal{S}, \mathbb{C}) \\ \downarrow \overline{pr_{\mathcal{H}_2}} \circ AJ & \searrow \mathcal{R}_{\mathcal{H}_1}^{\{p, i, m\}} & \downarrow \\ & \xrightarrow{\mathcal{R}_{\mathcal{H}_1}^{\{p, i, m\}}} & \frac{H^{p-i, p-m}(X, \mathbb{C})}{H^{\{p-m, i-m, 0\}}(X)} \otimes H^{i-1, 0}(\mathcal{S}, \mathbb{C}) \\ & \searrow \mathcal{R}_{\mathcal{H}_2}^{\{p, i, m\}} & \downarrow (*) \\ Ext_{\text{MHS}}^1(\mathbb{Q}(-p), \mathcal{H}_2) & \xrightarrow{(**)} & \left(\frac{H^{2p-m-i}(X)}{F_h^{p-i}} \right)_{\mathbb{C}}^{(p-i, p-m)} \otimes H^{i-1, 0}(\mathcal{S}, \mathbb{C}). \end{array}$$

The proof in [Le2] implies that $\overline{\mathcal{R}_{\mathcal{H}_1}^{\{p, i, m\}}}$ is well-defined on the subquotient $\mathcal{L}^i \underline{CH}_{m\text{-ind}}^p(X_K, m)$ of $CH^p(X \times \mathcal{S}, m)$, and this gives (6.8). Moreover, $\overline{pr_{\mathcal{H}_2}} \circ AJ$ is just the map of our Theorem 6.1 (so this is also well-defined on $\mathcal{L}^i \underline{CH}_{m\text{-ind}}^p$). Now the GHC says that $F_h^{p-i} H^{2p-m-i}(X) = N^{p-i} H^{2p-m-i}(X)$; and Lewis shows that the Hard Lefschetz Conjecture (implied by GHC) gives $\frac{H^{p-i, p-m}(X, \mathbb{C})}{H^{\{p-m, i-m, 0\}}(X)} \cong \left(\frac{H^{2p-m-i}(X)}{N^{p-i}} \right)_{\mathbb{C}}^{(p-i, p-m)}$. Hence $\text{GHC} \implies (*)$ is an isomorphism, which means that $\overline{pr_{\mathcal{H}_2}} \circ AJ$

factors $\overline{\mathcal{R}_{\mathcal{H}_1}^{\{p,i,m\}}}$. The point is that we conjecturally aren't losing any information via (*), while we definitely are losing data along (**).

6.4. Second application: Pontryagin cycles. We now describe the improvement of [CF, Prop. 1.1]. Let C be a smooth projective curve $/k$ of genus g . Fix $i \in \mathbb{N}$; $q_1, \dots, q_i \in \mathcal{C}(k)$; $o \in J(\mathcal{C})(k)$ [origin of group law]; and $P := (p_1, \dots, p_i) \in \mathcal{C}^{\times i}(\mathbb{C})$ very general, i.e. $K := k(P) \subset \mathbb{C}$ has $\text{trdeg}(K/k) = i$. We define

$$\tilde{B}_i : Z_{\mathbb{R}}^p(J(C), m) \rightarrow Z_{\mathbb{R}}^p(J(C)_K, m)$$

by $\mathcal{Z} \mapsto \mathcal{Z} * (AJ(p_1 - q_1) - o) * \dots * (AJ(p_i - q_i) - o) =: \mathcal{Z}(i)$. This induces

$$B_i : CH_{\text{hom}}^p(J(C), m) \rightarrow \mathcal{L}^{i+1} \underline{CH}^p(J(C)_K, m)$$

since the *Pontryagin cycle* $\mathcal{Z}(i)$ is in $\mathcal{L}^{i+1}$ by the parallelogram law [Bl3], [Ke2, 3.2].

Theorem 6.3. *Assume*

$$(6.9) \quad i \leq 2p - g - m - 1,$$

and let $\langle \mathcal{Z} \rangle \in CH_{\text{hom}}^p(J(C), m)$ be given.

(a) If $AJ(\mathcal{Z}) \neq 0$, then $AJ_i(\mathcal{Z}(i)) \neq 0$, hence

$$0 \neq \langle \mathcal{Z}(i) \rangle \in Gr_{\mathcal{L}}^{i+1} \underline{CH}^p(J(C)_K, m).$$

(b) Let $m \geq 1$. If $\mathcal{Z}$ is regulator-indecomposable, then

$$0 \neq \overline{\langle \mathcal{Z}(i) \rangle} \in Gr_{\mathcal{L}}^{i+1} \underline{CH}_{1\text{-ind}}^p(J(C)_K, m);$$

in particular $\mathcal{Z}(i)$ is (strongly) indecomposable.

Remark 6.4. The case corresponding to [CF] is (b) $m = 1$. No assumption on the sub-Hodge structures of the $H^i(J(C), \mathbb{Q})$ is required.

We establish this with a parade of definitions and lemmas. Define cycles $/k$

$$\mathfrak{A}_i := \sum_{j=0}^i (-1)^j \sum_{|J|=j} \left\{ (u; c_1, \dots, c_i) \mid u = \sum_{\ell \in J} AJ_C(c_\ell - q_\ell) \right\} \in Z^g(J(C) \times C^{\times i}),$$

$$\mathfrak{G} := \{(u_1, u_2, u_3) \mid u_1 + u_2 = u_3\} \in Z^{2g}(J(C)^{\times 3}).$$

Let π_I , $I \subseteq \{1, 2, 3, 4\}$, denote various projections from $J(C)^{\times 3} \times C^{\times i}$ (where factors 1, 2, 3 are copies of $J(C)$, and $C^{\times i}$ is the 4th factor). In the otherwise self-explanatory diagram

$$\begin{array}{ccc}
& H^{2p-m-1}(J(C) \times C^{\times i}) & \\
& \downarrow \text{K\"unneth} & \\
H^{2p-m-1}(J(C)) & \xrightarrow{\beta_i} & H^{2p-m-i-1}(J(C)) \otimes H^i(C^{\times i}) =: \mathcal{H} \\
\downarrow p & \searrow \bar{\beta}'_i & \downarrow p' \\
\frac{H^{2p-m-1}(J(C))}{F_h^{p-1}} & & \frac{H^{2p-m-i-1}(J(C))}{F_h^{p-i}} \otimes \underline{H}^i(\eta_{C^i}) \\
& \searrow \bar{\beta}''_i & \downarrow p'' \\
& & \frac{H^{2p-m-i-1}(J(C))}{F_h^{p-i-1}} \otimes \underline{H}^i(\eta_{C^i})
\end{array}$$

define

$$\beta_i(\xi) := \pi_{34*} \{ \pi_{123}^*[\mathfrak{G}] \cup \pi_{24}^*[\mathfrak{A}_i] \cup \pi_1^* \xi \}.$$

Since β_i is induced by action of a correspondence, all arrows are morphisms of (M)HS. (Note that well-definedness of $\bar{\beta}_i''$ is clear once one knows $\bar{\beta}'_i$ is a morphism of HS.) We also write

$$\begin{array}{ccc}
H^{2p-m-1}(J(C), \mathbb{C}) & \xrightarrow{\bar{\beta}_i^{(i,0)}} & H^{2p-m-i-1}(J(C), \mathbb{C}) \otimes H^{i,0}(C^{\times i}, \mathbb{C}) \\
& \searrow \bar{\beta}_i & \uparrow pr^{(i,0)} \\
& & H^{2p-m-i-1}(J(C), \mathbb{C}) \otimes H^i(C^{\times i}, \mathbb{C}).
\end{array}$$

Lemma 6.5. *If (6.9) holds, $\bar{\beta}_i^{(i,0)}$ and $\bar{\beta}_i$ are injective.*

Proof. It suffices to examine $\bar{\beta}_i^{(i,0)}$. Note that pullback along $\{p \mapsto AJ(p-q)\} : C \rightarrow J(C)$ is independent of $q \in C$, and call this AJ^* . Let $\{\Omega_\ell\}_{\ell=1}^g \subset \Omega^1(J(C))$ be a basis such that $\{\omega_\ell := AJ^* \Omega_\ell\} \subset \Omega_C^1$ satisfies $\int_C \omega_k \wedge \overline{\omega_\ell} = \delta_{k\ell}$. Write $\Omega_k := \bigwedge_{k \in K} \Omega_k$, and let $\tau_{IJ} := \Omega_I \wedge \overline{\Omega_J}$ for any pair of multi-indices $I, J \subseteq \{1, \dots, g\}$ with $|I| + |J| = 2g - 2p + m + 1$.

Given nonzero $\xi \in H^{2p-m-1}(J(C))$, $\exists I, J$ such that $\langle \xi, \tau_{IJ} \rangle \neq 0$. Since (6.9) holds, we may take $K = \{k_1, \dots, k_i\} \subseteq \{1, \dots, g\} \setminus I \cup J$. Write $\pi_0 : J(C) \times C^{\times i} \rightarrow J(C)$, $\pi_j : J(C) \times C^{\times i} \rightarrow C$ ($j = 1, \dots, i$) for the obvious projections. By an easy calculation (essentially the same as that in [CF, Lemma 1.3]),

$$\langle \beta_i(\xi), \pi_0^*(\Omega_K \wedge \tau_{IJ}) \wedge \pi_1^*(\overline{\omega_{k_1}}) \wedge \dots \wedge \pi_i^*(\overline{\omega_{k_i}}) \rangle = \pm \langle \xi, \tau_{IJ} \rangle;$$

hence $\bar{\beta}_i^{(i,0)}(\xi) \neq 0$. □

For a map of (pure $\mathbb{Q}$ -)Hodge structures $\mathcal{H}_1 \xrightarrow{\theta} \mathcal{H}_2$ of weight $< 2p$, write $Ext^1(\theta)$ for the induced map

$$Ext_{\text{MHS}}^1(\mathbb{Q}(-p), \mathcal{H}_1) \cong \frac{\mathcal{H}_{1,\mathbb{C}}}{F^p \mathcal{H}_{1,\mathbb{C}} + \mathcal{H}_1} \xrightarrow{\bar{\theta}} \frac{\mathcal{H}_{2,\mathbb{C}}}{F^p \mathcal{H}_{2,\mathbb{C}} + \mathcal{H}_2} \cong Ext_{\text{MHS}}^1(\mathbb{Q}(-p), \mathcal{H}_2).$$

Lemma 6.6. *The following diagram*

$$\begin{array}{ccc} CH_{\text{hom}}^p(J(C), m) & \xrightarrow{B_i} & \mathcal{L}^{i+1} \underline{CH}^p(J(C)_K, m) \\ \downarrow AJ & & \downarrow AJ_i^{\text{tr}} \\ Ext_{\text{MHS}}^1(\mathbb{Q}(-p), H^{2p-m-1}(J(C))) & \xrightarrow{Ext^1(\beta'_i)} & Ext_{\text{MHS}}^1\left(\mathbb{Q}(-p), \frac{H^{2p-m-i-1}(J(C))}{F_h^{p-i}} \otimes \underline{H}^i(\eta_{C^i})\right) \\ \downarrow Ext^1(\mathfrak{p}) & & \downarrow Ext^1(\mathfrak{p}'') \\ Ext_{\text{MHS}}^1\left(\mathbb{Q}(-p), \frac{H^{2p-m-1}(J(C))}{F_h^{p-1}}\right) & \xrightarrow{Ext^1(\beta'_{i'})} & Ext_{\text{MHS}}^1\left(\mathbb{Q}(-p), \frac{H^{2p-m-i-1}(J(C))}{F_h^{p-i-1}} \otimes \underline{H}^i(\eta_{C^i})\right) \end{array}$$

is commutative.

Proof. Let $\iota_P : J(C)[\times\{P\}] \hookrightarrow J(C) \times C^{\times i}$ be the inclusion (defined $/K$). Define a map $\mathfrak{B}_i$ factoring $\tilde{B}_i$

$$\begin{array}{ccc} Z_{\mathbb{R}}^p(J(C), m) & \xrightarrow{\tilde{B}_i} & Z_{\mathbb{R}}^p(J(C)_K, m) \\ & \searrow \mathfrak{B}_i & \uparrow \iota_P^* \\ & & Z_{\mathbb{R}}^p(J(C) \times C^{\times i}, m) \end{array}$$

by sending $\mathcal{Z} \mapsto \pi_{34}^* \{\pi_{123}^* \mathfrak{G} \cdot \pi_{24}^* \mathfrak{A}_i \cdot \pi_1^* \mathcal{Z}\} =: \overline{\mathfrak{Z}(i)}$. Clearly $\tilde{\mathfrak{B}}_i$ descends modulo $\overset{\text{rat}}{\equiv}$. Since $\overline{\mathfrak{Z}(i)}$ is a complete spread for $\mathcal{Z}(i)$, $AJ_i^{\text{tr}}(\mathcal{Z}(i))$ is a projection of $AJ_{J(C) \times C^i}(\overline{\mathfrak{Z}(i)})$ and it suffices to show

$$\begin{array}{ccc} CH_{\text{hom}}^p(J(C), m) & \xrightarrow{\mathfrak{B}_i} & CH_{\text{hom}}^p(J(C) \times C^{\times i}, m) \\ \downarrow AJ & & \downarrow AJ \\ Ext_{\text{MHS}}^1(\mathbb{Q}(-p), H^{2p-m-1}(J(C))) & \xrightarrow{Ext^1(\beta_i)} & Ext_{\text{MHS}}^1(\mathbb{Q}(-p), H^{2p-m-1}(J(C) \times C^{\times i})) \end{array}$$

commutes. Now $\overline{\mathfrak{Z}(i)}$ is a (signed) sum of families of translates (over $C^{\times i}$) of $\mathcal{Z}$. Translating $\mathcal{Z}$ on $J(C)$ gives a new (higher) cycle whose KLM currents δ_T , Ω , R are simply push-forwards of $\delta_{T_{\mathcal{Z}}}$, $\Omega_{\mathcal{Z}}$, $R_{\mathcal{Z}}$ along the translation. The same thing happens with the closed current $R' := R - (2\pi i)^* \delta_{\partial^{-1}T} + d^{-1}\Omega$, whose class in $H^{2p-m-1}(J(C), \mathbb{C})$ descends to $AJ_{J(C)}(\mathcal{Z})$. More precisely, we can choose

$$R'_{\overline{\mathfrak{Z}(i)}} := \pi_{34*} \{\pi_{123}^* \delta_{\mathfrak{G}} \cdot \pi_{24}^* \delta_{\mathfrak{A}_i} \cdot \pi_1^* R'_{\mathcal{Z}}\},$$

and taking classes gives the result. $\square$

Consider the sub-Hodge structures

$$\mathcal{N} := H^{2p-m-i-1}(J(C)) \otimes N^1 H^i(C^{\times i}),$$

$$\mathcal{F}_0 := F_h^{p-i} H^{2p-m-i-1}(J(C)) \otimes H^i(C^{\times i}),$$

$$\mathcal{F}_1 := F_h^{p-i-1} H^{2p-m-i-1}(J(C)) \otimes H^i(C^{\times i})$$

of $\mathcal{H}$; clearly $\mathfrak{p}'$ and $\mathfrak{p}'' \circ \mathfrak{p}'$ are (resp.) quotients of $\mathcal{H}$ by $\mathcal{N} + \mathcal{F}_0$, $\mathcal{N} + \mathcal{F}_1$.

Lemma 6.7. *Provided (6.9) holds, we have:*

$$(\mathcal{N} + \mathcal{F}_1) \cap \text{im}(\bar{\beta}_i) \subseteq \bar{\beta}_i(F_h^{p-i}),$$

$$(\mathcal{N} + \mathcal{F}_0) \cap \text{im}(\bar{\beta}_i) = \{0\}.$$

Proof. Consider $\xi \in H^{2p-m-1}(J(C), \mathbb{C})$ with $\bar{\beta}_i(\xi) \in \mathcal{N}_{\mathbb{C}} + \mathcal{F}_{\ell, \mathbb{C}}$, $\ell = 0$ or 1 . We have $\text{pr}^{(i,0)}(\mathcal{N}_{\mathbb{C}}) = \{0\}$, $F_h^{p-i-\ell} H^{2p-m-i-1}(J(C), \mathbb{C}) \subset F^{p-i-\ell}$, hence

$$(6.10) \quad \bar{\beta}_i^{(i,0)}(\xi) \in F^{p-i-\ell} H^{2p-m-i-1}(J(C), \mathbb{C}) \otimes H^{i,0}(C^{\times i}, \mathbb{C}).$$

Since $\bar{\beta}_i$ is a morphism of HS, it decomposes $(\otimes \mathbb{C})$ by type; hence $\bar{\beta}_i^{(i,0)}$ must decompose into a direct sum (over $a \in \mathbb{Z}$) of maps

$$H^{a, 2p-m-a-1}(J(C), \mathbb{C}) \rightarrow H^{a-i, 2p-m-a-1}(J(C), \mathbb{C}) \otimes H^{i,0}(C^{\times i}, \mathbb{C}).$$

These are injective (because $\bar{\beta}_i^{(i,0)}$ is), so (6.10) $\implies \xi \in F^{p-\ell} H^{2p-m-1}(J(C), \mathbb{C})$.

Now $\bar{\beta}_i^{-1}(\mathcal{N} + \mathcal{F}_{\ell})$ is necessarily a subHS of $H^{2p-m-1}(J(C))$. The last paragraph shows $\bar{\beta}_i^{-1}(\mathcal{N}_{\mathbb{C}} + \mathcal{F}_{\ell, \mathbb{C}}) \subset F^{p-\ell}$ ($\ell = 0, 1$); hence

$$\bar{\beta}_i^{-1}(\mathcal{N} + \mathcal{F}_{\ell}) \subseteq F_h^{p-\ell} H^{2p-m-1}(J(C)).$$

Applying $\bar{\beta}_i$ to this gives the result, noting (for $\ell = 0$) that $F_h^p H^{2p-m-1}(J(C)) = \{0\}$. $\square$

Proof. (of Theorem 6.3). Since $\bar{\beta}_i$ is injective (Lemma 6.5), Lemma 6.7 implies $\bar{\beta}'_i$, $\bar{\beta}''_i$, hence $\text{Ext}^1(\bar{\beta}'_i)$, $\text{Ext}^1(\bar{\beta}''_i)$, are injective too. By assumption, (a) $AJ(\mathcal{Z})$ resp. (b) $\text{Ext}^1(\mathfrak{p})(AJ(\mathcal{Z}))$ is nonzero. So Lemma 6.6 $\implies$ nonvanishing of (a) $AJ_i^{tr}(\mathcal{Z}(i))$ resp. (b) $\text{Ext}^1(\mathfrak{p}'')(AJ_i^{tr}(\mathcal{Z}(i)))$, as desired. $\square$

7. REDUCED HIGHER ABEL-JACOBI MAPS

We now give a description of part of each higher Abel-Jacobi invariant in the spirit of Griffiths-Green [GG]. As in §5.1 we avoid double complexes, but we don't restrict to the “nice case” examined there.

Roughly speaking: reduced higher AJ maps are, like the AJ_i^{tr} we have been using, Hodge-theoretically defined quotients of the previously defined AJ_i . Only we quotient somewhat farther, in order to get something which is explicitly computable in terms of generalized membrane integrals on X — even when one does not have a homologically trivial complete spread. Our expectation is that they will be useful in special cases of arithmetic interest (as have their analogues for 0-cycles, see [Ke1, sec. 16], [Ke2, sec. 6.3]), rather than for proving results of a general flavor (such as Theorems 6.2 and 6.3 above). Assume $m > 0$ for the discussion.

Set $V = F^{d-p+i} H^{2d-2p+m+i}(X, \mathbb{C})$. The “period lattice”

$$\Lambda := \text{im} \left\{ H^{2p-m-i}(X, \mathbb{Q}(p)) \rightarrow \frac{H^{2p-m-i}(X, \mathbb{C})}{F^{p-i+1} H^{2p-m-i}(X, \mathbb{C})} \cong V^\vee \right\}$$

has in general $\dim_{\mathbb{Q}} \Lambda \neq 2 \dim_{\mathbb{C}} V$. There are (well-defined) maps

$$(7.1) \quad Gr_{\mathcal{L}}^i CH^p(X_K, m) \xrightarrow{\overline{AJ_{i-1}}} Hom_{\mathbb{Q}}(\overline{H_{i-1}}(\eta_S, \mathbb{Q}), V^\vee / \Lambda)$$

where $\overline{H_{i-1}}(\eta_S) := \text{coim}\{H_{i-1}(\eta_S) \rightarrow H_{i-1}(\mathcal{S})\}$, and (for cycles without complete spread)

$$(7.2) \quad Gr_{\mathcal{L}}^i CH^p(X_K, m) \xrightarrow{\overline{cl_i}} Hom_{\mathbb{C}}(V, F^i W_{m+i} H^i(\eta_S, \mathbb{C})),$$

$$(7.3) \quad \ker(\overline{cl_i}) \xrightarrow{\overline{AJ_{i-1}}} Hom_{\mathbb{Q}}(H_{i-1}(\eta_S, \mathbb{Q}), V^\vee / \Lambda).$$

We note that as $\ker(\overline{cl_i}) \supseteq \ker(cl_i)$ is not in general an equality, $\overline{AJ_{i-1}}$ may be defined “before” AJ_{i-1} (of §5.1) is.

Whether or not $\mathcal{Z} \in \mathcal{L}^i$ has complete k -spread $\mathfrak{Z}$ ($\partial_{\mathcal{B}}$ -closed in $Z^p(X \times \mathcal{S}, m)$) does not matter so much for the formulas for $\overline{AJ_{i-1}}$. In either case one has (for $U \subset \mathcal{S}$ “sufficiently small” Zariski open) $\mathfrak{Z}_U \in Z_{\mathbb{R}}^p(X \times U, m)$, $\partial_{\mathcal{B}}$ -closed, with irreducible components equidimensional $/U$, and restricting to the class of $\mathfrak{Z} = \mathfrak{s}(\mathcal{Z}) \in CH^p(X \times \eta_S, m)$. Given $[\gamma] \in H_{i-1}(\eta_S, \mathbb{Q})$ resp. $\overline{H_{i-1}}(\eta_S, \mathbb{Q})$, its image in $H_{i-1}(U)$ resp. $\overline{H_{i-1}}(U)$ may be represented by $\gamma \in Z_{i-1}^{top}(U, \mathbb{Q})$ such that $\gamma \times X \times \square^m$ intersects

properly all $\mathfrak{Z}_U \cap \{\mathcal{S} \times X \times \partial_{f,\mathbb{R}}^{I,\ell} \square^m\}$. (See §8.2 for a definition of $\partial_{f,\mathbb{R}}^{I,\ell} \square^m$.) Finally, we take classes $\in V$ to be represented by test forms $\omega \in Z_d(X, F^{d-p+i} \Omega_{X_\infty}^{2d-2p+m+i})$.

Writing $\pi_X : X \times \mathcal{S} \rightarrow X$, $\pi_U : X \times U \rightarrow U$, define

$$T_3(\gamma) := \pi_X((\gamma \times X) \cap T_3) \in Z_{2d-2p+m+i}^{top}(X, \mathbb{Q}),$$

$$R_3(\gamma) := \pi_{X*}(\delta_{\gamma \times X} \cdot R_3) \in \mathcal{D}^{2p-m-i}(X)$$

(i.e. $(R_3(\gamma))(\omega) := \int_{\gamma \times X} \pi_X^* \omega \wedge R_3$ defined by appropriate $\lim_{\epsilon \rightarrow 0} \int$'s), and

$$\Omega_3(\omega) := \pi_{U*}(\pi_X^* \omega \wedge \Omega_3) \in \Gamma_{d-cl.}(U, \mathcal{D}_U^{i,0}) = \Omega^i(U).$$

Note $T_3(\gamma) \stackrel{\text{hom}}{=} 0$ since $\mathfrak{Z} \in \mathcal{L}^i \implies [\mathfrak{Z}]_{i-1} = 0$. The following result says how to compute (7.1), (7.2), (7.3).

Theorem 7.1. (i) Given $\langle \mathcal{Z} \rangle \in \mathcal{L}^i CH^p(X_K, m)$, $[\omega] \in V$, $\overline{cl}_i \langle \mathcal{Z} \rangle [\omega]$ is computed by $(2\pi i)^{p-m} \Omega_3(\omega) \in \Omega^i(U)$.

(ii) Given $\langle \mathcal{Z} \rangle \in \{\ker(\overline{cl}_i) \subseteq \mathcal{L}^i CH^p(X_K, m)\}$ resp. $\mathcal{L}^i \underline{CH}^p(X_K, m)$, $[\gamma] \in H_{i-1}(\eta_S, \mathbb{Q})$ resp. $\overline{H}_{i-1}(\eta_S, \mathbb{Q})$, $\overline{AJ}_{i-1} \langle \mathcal{Z} \rangle [\gamma]$ is computed by

$$(2\pi i)^{p-m} R_3(\gamma) + (2\pi i)^p \delta_{\partial^{-1} T_3(\gamma)} =: (2\pi i)^{p-m} R_3(\gamma)' \in \mathcal{D}^{2p-m-i}(X),$$

viewed as a functional on V .

Proof. In the complete spread case (7.1), the Theorem essentially follows from writing $\overline{AJ}_{i-1}$ as the composition of AJ_{i-1} with a projection and using the end of §5.1. For the projection, start with

$$\begin{aligned} Ext_{MHS}^1 \left(\mathbb{Q}(-p), \underline{H}^{i-1}(\eta_S) \otimes \frac{H^{2p-m-i}(X)}{F_h^{p-i+1} H^{2p-m-i}(X)} \right) &\cong \\ \frac{\underline{H}^{i-1}(\eta_S, \mathbb{C}) \otimes \frac{H^{2p-m-i}(X, \mathbb{C})}{F_h^{p-i+1} H^{2p-m-i}(X, \mathbb{C})}}{F^p \{\text{num}\} + H^{i-1}(\eta_S, \mathbb{Q}) \otimes \frac{H^{2p-m-i}(X, \mathbb{Q}(p))}{F_h^{p-i+1} H^{2p-m-i}(X, \mathbb{Q}(p))}}. \end{aligned}$$

Since $F_h^{p-i+1} H^{2p-m-i}(X, \mathbb{C}) \subseteq F^{p-i+1} H^{2p-m-i}(X, \mathbb{C})$, and $F^p \left(H^{i-1}(\eta_S, \mathbb{C}) \otimes \frac{H^{2p-m-i}(X, \mathbb{C})}{F_h^{p-i+1} H^{2p-m-i}(X, \mathbb{C})} \right) = 0$, this projects to

$$\frac{\underline{H}^{i-1}(\eta_S, \mathbb{C}) \otimes \frac{H^{2p-m-i}(X, \mathbb{C})}{F_h^{p-i+1} H^{2p-m-i}(X, \mathbb{C})}}{im \left\{ \underline{H}^{i-1}(\eta_S, \mathbb{Q}) \otimes \frac{H^{2p-m-i}(X, \mathbb{Q}(p))}{F_h^{p-i+1}} \right\}} \cong \underline{H}^{i-1}(\eta_S, \mathbb{Q}) \otimes V^\vee / \Lambda$$

and thus to the r.h.s. of (7.1).

In the incomplete spread case (7.3), the result is considerably deeper and involves a study of meromorphic $V^\vee/\Lambda$ -valued differential characters (sufficiently parallel to §§10, 11 of [Ke1] that we won't give details). $\square$

There are three special cases or extensions. First, there is a version of (7.1) for indecomposables, namely (for $1 \leq j \leq m \leq i \leq p$)

$$\mathcal{L}^i \underline{CH}_{j\text{-ind}}^p(X_K, m) \xrightarrow{\overline{AJ}_{i-1, \text{ind}}} \text{Hom}_{\mathbb{Q}} \left(\overline{H}_{i-1}(\eta_S, \mathbb{Q}), \frac{\{F^{d-p+i+1} H^{2d-2p+m+i}(X, \mathbb{C})\}^\vee}{\text{im}\{H_{2d-2p+m+i}(X, \mathbb{Q}(p))\}} \right),$$

again computed by $(2\pi i)^{p-m} R_{\mathfrak{Z}}(\gamma)'$ (only viewed as a functional [mod periods] on a smaller space). For example, for $\mathcal{L}^2 \underline{CH}_{\text{ind}}^3(X_K, 1)$ with X a 3-fold, the target is $\text{Hom} \left(\overline{H}_1(\eta_S), \frac{\{\Omega^3(X)\}^\vee}{\text{im}\{H_3(X, \mathbb{Q}(3))\}} \right)$.

Second, there is the heretofore neglected case $m = 0$ of algebraic cycles. Although these have complete spread $\mathfrak{Z}$, it is not possible in general to arrange $\mathfrak{Z} \stackrel{\text{hom}}{=} 0$; so this is more like the situation of (7.2), (7.3) than (7.1). One has a map

$$\{ker(cl_i) \subseteq \mathcal{L}^i CH^p(X_K)\} \xrightarrow{\overline{AJ}_{i-1}} \text{Hom}_{\mathbb{Q}} \left(\overline{H}_{i-1}(\eta_S, \mathbb{Q}), \frac{\{F^{d-p+i} H^{2d-2p+i}(X, \mathbb{C})\}^\vee}{\Lambda} \right),$$

with $\overline{AJ}_{i-1} \langle \mathcal{Z} \rangle [\gamma]$ computed by $(2\pi i)^p \int_{\partial^{-1}\mathfrak{Z}(\gamma)} (\cdot)$ since $R_{\mathfrak{Z}} = 0$ and $T_{\mathfrak{Z}} = \mathfrak{Z}$. (Here $\mathfrak{Z}(\gamma) := \pi_X \{\mathfrak{Z} \cap (\gamma \times X)\}$.)

Finally, there is the case of $X = \text{Spec} K$, a point. In order for V to be $\mathbb{C}$ and not $\{0\}$, we must have $2d - 2p + m + i = 0$ (with $d = 0$), hence $i = 2p - m$. Moreover, $d - p + i \leq 0$ (again with $d = 0$) $\implies p \leq m$. (7.2) maps $Gr_{\mathcal{L}}^{2p-m} CH^p(\text{Spec} K, m)$ to (dropping the weight) $F^{2p-m} H^{2p-m}(\eta_S, \mathbb{C})$, and is given by $\mathcal{Z}(\mapsto \mathfrak{Z}) \mapsto \Omega_{\mathfrak{Z}}$. This is zero if $\text{trdeg}(K/k) = 2p - m - 1$, in which case (7.3) reduces to

$$Gr_{\mathcal{L}}^{2p-m} CH^p(\text{Spec} K, m) \rightarrow \text{Hom}_{\mathbb{Q}}(H_{2p-m-1}(\eta_S, \mathbb{Q}), \mathbb{C}/\mathbb{Q}(p)) \cong J^{p,m}(\eta_S),$$

represented by $(2\pi i)^{p-m} R'_{\mathfrak{Z}}$ (a d -closed $(2p - m - 1)$ -current on some $U \subset \mathcal{S}$). This is just the [KLM] formula for AJ on η_S .

8. ABEL-JACOBI FOR MOTIVIC COHOMOLOGY OF RELATIVE AND SINGULAR VARIETIES

In this section we essentially reverse the Gysin spectral sequences of §§3, 5. This allows us to study, for example, AJ maps on $H_{\mathcal{M}}^{2p-n}(\mathcal{Y}, \mathbb{Q}(p))$ rather than $H_{\mathcal{M}, \mathcal{Y}}^{2p-n+2}(\mathcal{X}, \mathbb{Q}(p+1)) \cong CH^p(\mathcal{Y}, n)$ when $\mathcal{Y} \subset \mathcal{X}$ is a complete NCD. This is technically more involved, since pulling back (cycles, currents) is considerably harder than pushing forward. The reward is in the arithmetically fascinating examples of §8.4, as well as [DK] and other forthcoming papers.

8.1. Currents with pullback. We follow [Ki, sec. 4] for the first part of this subsection.

Let X be a smooth projective analytic variety over $\mathbb{C}$. Recall that a current $T \in \mathcal{D}_X^r(U)$ on an open subset of X is a continuous linear functional on C^∞ forms compactly supported on U .

Definition 8.1. *Normal currents* on open $U \subset X$ are defined by

$$\mathcal{N}_X^r(U) := \left\{ T \in \mathcal{D}_X^r(U) \left| \begin{array}{l} T, dT \text{ admit extensions to} \\ \text{continuous linear functionals on} \\ \text{continuous forms compactly supported on } U \end{array} \right. \right\};$$

$\mathcal{N}_X^r$ is their sheafification.

Now let $V \subset X$ be a smooth closed subvariety of codimension q , and set

$$\Psi := \text{characteristic function of } X \setminus V.$$

Call $T \in \mathcal{N}_X^r(U)$ of *V-residue type* if $T = \Psi T$, and *V-transversal* if also $dT = \Psi dT$. We first treat the case of $q = 1$:

Definition 8.2. Assuming $V \cap U \subset U$ defined by $z = 0$,

$$\mathcal{N}_X^r\{V\}(U) := \left\{ T \in \mathcal{N}_X^r(U) \left| \begin{array}{l} T \text{ is } V\text{-transversal, and } \exists \\ S, \mathfrak{S} \in \mathcal{N}_X^{r+1}(U) \text{ of } V\text{-residue type} \\ \text{such that } T \wedge dz = zS, dT \wedge dz = z\mathfrak{S} \end{array} \right. \right\}$$

are the *normal currents of V-intersection type* on U , whose sheafification yields $\mathcal{N}_X^r\{V\}$.

Intuitively, we require the “singularities” of our currents to properly intersect V .

Remark 8.3. Note that $S, \mathfrak{S}$ are unique if they exist; we think of S as “ $T \wedge \frac{dz}{z}$ ”. By definition S operates on continuous forms; hence T operates on continuous $\log(V)$ forms, *a fortiori* on $C^\infty \log(V)$ forms.

The map

$$\iota_V^* : \mathcal{N}_X^r\{V\} \rightarrow \mathcal{N}_V^r$$

is given by sending $T \mapsto (1 - \Psi)dS =: \iota_V^*(T)$. One may think of $\iota_V^*(T)$ as “ $\text{Res}_V(T \wedge \frac{dz}{z})$ ”; showing this is indeed (the push-forward of) a current on V uses normality of T .

Next, for arbitrary codimension q , let $\sigma : \tilde{X} \rightarrow X$ be the blowup along V , with $\sigma^{-1}(V) =: \tilde{V}$ and $\sigma^{-1}(U) =: \tilde{U}$. (Note that the pullback of currents is defined along the projective-bundle projection $\sigma|_{\tilde{V}}$.)

Definition 8.4. More generally, $\mathcal{N}_X^r\{V\}(U) :=$

$$\sigma_* \left\{ \tilde{T} \in \mathcal{N}_{\tilde{X}}^r\{\tilde{V}\}(\tilde{U}) \mid \iota_{\tilde{V}}^*(\tilde{T}) = (\sigma|_{\tilde{V}})^* \tau \text{ for some } \tau \in \mathcal{N}_V^r(U \cap V) \right\}.$$

Finally, we consider the case where $V = \bigcup_{i=1}^N V_i$ is a normal crossing divisor (NCD) on X . (Here we depart from [Ki].) Writing $I, J, K \subseteq \{1, \dots, N\}$ for multi-indices, each $V_I := \bigcup_{i \in I} V_i$ must be smooth of codimension $|I|$. Note that if T is V_i -transversal ($\forall i$), then T is V_I -transversal ($\forall I$).

Take $U \subset X$ sufficiently small that

- (a) $\exists K = \{k_1, \dots, k_m\}$, $m \leq d := \dim(X)$, such that only $V_{k_1}, \dots, V_{k_m}$ intersect U ; and
- (b) $\exists$ analytic chart $\{z_{k_1}, \dots, z_{k_m}; \underline{w}\}$ on U such that $U \ni \{\underline{0}\}$, $U \cap V_k = \{z_k = 0\}$ and $U \cap V = \{z_{k_1} \cdots z_{k_m} = 0\}$.

We write locally $dz_I := \bigwedge_{i \in I \cap K} dz_i$, $z_I := \prod_{i \in I \cap K} z_i$.

Definition 8.5. For $V = \text{NCD}$, $T \in \mathcal{N}_X^r\{V\}(U) \iff$ the following hold:

- (a) $T \in \mathcal{N}_X^r(U)$ is V_j -transversal ($\forall j$)
- (b) $\forall I$, $\exists S_I, \mathfrak{S}_I \in \mathcal{N}_X^{r+|I|}(U)$
 - (i) satisfying $T \wedge dz_I = z_I S_I$, $dT \wedge dz_I = z_I \mathfrak{S}_I$; and
 - (ii) of V_j -residue type $\forall j \in I$,
 - (iii) V_j -transversal $\forall j \notin I$,
 - (iv) V_J -transversal $\forall J$ with $|J| > 1$.

Remark 8.6. Again, if such $S_I, \mathfrak{S}_I$ exist then they are unique; the sheaves $\mathcal{N}_X^r\{V\}$ are called (once more) normal currents of V -intersection type.

To describe the pullbacks, write $X = V_\emptyset$, and put

$$V^I := V_I \bigcap \bigcup_{j \notin I} V_j \subseteq V_I$$

(e.g., $V^\emptyset = V$ itself). Then $\mathcal{N}_{V_I}^r\{V^I\}$ has pullback maps to any $\mathcal{N}_{V_J}^r\{V^J\}$ for $J \supseteq I$, and pullbacks commute. More precisely, the assignment $T \mapsto (1 - \Psi_j)dS_j =: \iota_{V_j}^*(T)$ generalizes for $J = \{j_1, \dots, j_\ell\}$ to $\iota_{V_J}^* : \mathcal{N}_X^r\{V\} \rightarrow \mathcal{N}_{V_J}^r\{V^J\}$ by taking $\iota_{V_J}^* := \iota_{V_J}^* \circ \iota_{V_{J \setminus \{j_\ell\}}}^* \circ \dots \circ \iota_{V_{\{j_1, j_2\}}}^* \circ \iota_{V_{j_1}}^*$. That this is independent of the ordering of the elements of J follows from the existence of the $S_{J'}$ for all $J' \subseteq J$.

Here is the Poincaré lemma for intersection currents.

Lemma 8.7. *The inclusions*

$$\Omega_X^\bullet \subset \Omega_{X^\infty}^\bullet \subset \mathcal{N}_X^\bullet\{V\} \subset \mathcal{N}_X^\bullet \subset \mathcal{D}_X^\bullet$$

are $F^\bullet$ -filtered quasi-isomorphisms in the cases considered above (Def. 8.2, 8.4, 8.5).

The main idea here is to show that King's Cauchy slice-kernels K_i (which furnish the coboundary between $\cap$ -currents and holomorphic forms) preserve $F^p\mathcal{N}_X^\bullet\{V\}$ at the stalk level. Roughly speaking, given $S, \mathfrak{S}$ for T , one must produce normal currents $S', \mathfrak{S}'$ for $K_i(\text{cutoff function} \cdot T)$. For the non-NCD cases, this is done in [Ki]; for $V = \text{NCD}$ the formulas for $S', \mathfrak{S}'$ are more complicated (we don't give them here), but the other details of the proof are the same.

Now assume $V \subset X$ is a NCD (with both X and V complete), and write $C_{top}^r\{V\}(X)$ for real-codimension- r C^∞ -chains properly intersecting all the V_I . Set

$$C_{\mathcal{D}}^\bullet(X, \mathbb{Q}(p))_V := \text{Cone} \left\{ \begin{array}{c} C_{top}^\bullet\{V\}(X, \mathbb{Q}(p)) \\ \oplus F^p\mathcal{N}_X^\bullet\{V\}(X) \end{array} \longrightarrow \mathcal{N}_X^\bullet\{V\}(X) \right\} [-1].$$

It is an obvious corollary of Lemma 8.7 that this computes Deligne cohomology of X ; that is,

$$C_{\mathcal{D}}^\bullet(X, \mathbb{Q}(p))_V \xrightarrow{\simeq} C_{\mathcal{D}}^\bullet(X, \mathbb{Q}(p)).$$

Now define a double complex $\mathfrak{K}_V^{\bullet, \bullet}(p)$ by

$$\mathfrak{K}_V^{\ell, m}(p) := \oplus_{|I|=\ell} C_{\mathcal{D}}^{2p+m}(V_I, \mathbb{Q}(p))_{V^I}$$

with differentials $D : \mathfrak{K}_V^{\ell, m}(p) \rightarrow \mathfrak{K}_V^{\ell, m+1}(p)$ (Deligne homology differential) and

$$(8.1) \quad \sum_{|I|=\ell} \sum_{i \notin I} (-1)^{<i>_I} (\iota_{V_{I \cup i} \subset V_I})^* = \mathfrak{J} : \mathfrak{K}_V^{\ell, m}(p) \rightarrow \mathfrak{K}_V^{\ell+1, m}(p),$$

where $<i>_I$ means the position in which i appears in $\{1, \dots, N\} \setminus I$. Write $s^\bullet$ for associated simple (total) complex, and $\mathbb{D}$ for total differential.

Proposition 8.8. *Cohomology of*

$$C_{\mathcal{D}}^{2p+\bullet}(V, \mathbb{Q}(p)) := s^\bullet(\mathfrak{K}_V^{\bullet, \geq 1, \bullet}(p))[1]$$

$$\text{resp. } C_{\mathcal{D}}^{2p+\bullet}((X, V), \mathbb{Q}(p)) := s^\bullet(\mathfrak{K}_V^{\bullet, \geq 0, \bullet}(p))$$

computes Deligne cohomology of V resp. (X, V) .

Remark 8.9. We emphasize the distinction (quite significant for singular V) between Deligne cohomology $H_{\mathcal{D}}^*(V, \mathbb{Q}(p))$ of V , and $H_{\mathcal{D}, V}^*(X, \mathbb{Q}(p))$ — which is Deligne homology of V .

Noting that $\mathfrak{K}_V^{0, \bullet}(p) = C_{\mathcal{D}}^{2p+\bullet}(X, \mathbb{Q}(p))_V$, we see that

$$C_{\mathcal{D}}^{2p+\bullet}((X, V), \mathbb{Q}(p)) \xrightarrow{\cong} \text{Cone} \left\{ C_{\mathcal{D}}^{2p+\bullet}(X, \mathbb{Q}(p))_V \rightarrow C_{\mathcal{D}}^{2p+\bullet}(V, \mathbb{Q}(p)) \right\} [-1];$$

this gives rise to a long-exact sequence

(8.2)

$$\rightarrow H_{\mathcal{D}}^{q-1}(V, \mathbb{Q}(p)) \rightarrow H_{\mathcal{D}}^q((X, V), \mathbb{Q}(p)) \rightarrow H_{\mathcal{D}}^q(X, \mathbb{Q}(p)) \rightarrow H_{\mathcal{D}}^q(V, \mathbb{Q}(p)) \rightarrow .$$

For $Y := X, V$, or (X, V) one has as usual $H_{\mathcal{D}}^q(Y, \mathbb{Q}(p)) \cong H_{\mathcal{H}}^q(Y, \mathbb{Q}(p))$ whenever $W_{2p}H^q(Y, \mathbb{Q}) = H^q(Y, \mathbb{Q})$, and this is the case here when $q \leq 2p$.

8.2. Higher Chow chains with pullback. Let X be a nonsingular projective algebraic variety over a field K , with $\mathbb{Q} \subseteq K \subseteq \mathbb{C}$. For technical reasons, we assume K has a finitely generated field extension $L \subseteq \mathbb{C}$ invariant under complex conjugation.

Given $I \subseteq \{1, \dots, n\}$, $f : I \rightarrow \{0, \infty\}$, $j \in \hat{I} \cup \{0\}$ (where $\hat{I} := \{1, \dots, n\} \setminus I$), define subsets

$$\begin{aligned} \partial_f^I \square^n &:= \cap_{\ell \in I} \{z_\ell = f(\ell)\}, \\ \partial_{f, \mathbb{R}}^{I, j} \square^n &:= \begin{cases} [\cap_{\ell \in \{1, \dots, j\}} T_{z_\ell}] \cap \partial_f^I \square^n & \text{if } j \geq 1 \\ \partial_f^I \square^n & \text{if } j = 0 \end{cases} \end{aligned}$$

of $\square^n$. Write $\iota_f^I : \square^{n-|I|} \hookrightarrow \square^n$, $\pi_I : \square^n \twoheadrightarrow \square^{n-|I|}$ for the obvious inclusions and projections.

Definition 8.10. $Z_{\mathbb{R}}^p(X, n)$ is the free abelian group $(\otimes \mathbb{Q})$ generated by irreducible $\mathfrak{Z} \in Z^p((X \times \square^n)_K)$ with $\mathfrak{Z}_{\mathbb{C}}$ meeting all $X_{\mathbb{C}} \times \partial_{f, \mathbb{R}}^{I, j} \square^n$ properly as real analytic varieties, modulo degenerate cycles. Write $\partial_f^I \mathfrak{Z} := (\iota_f^I)^* \mathfrak{Z} \in Z_{\mathbb{R}}^p(X, n - |I|)$, $\partial_{\mathcal{B}} \mathfrak{Z} := \sum_{j=1}^n (-1)^j (\partial_o^j - \partial_\infty^j) \mathfrak{Z} \in Z_{\mathbb{R}}^p(X, n - 1)$.

Remark 8.11. Recall $\mathfrak{Z} \in Z^p(X, n)$ if the intersections $\mathfrak{Z} \cap (X \times \partial_f^I \square^n)$ are proper. The following description is useful for checking whether a higher Chow chain is in good real position: given irreducible $\mathfrak{Z} \in Z^p(X, n)$, $\mathfrak{Z}$ belongs to $Z_{\mathbb{R}}^p(X, n)$ iff (i) $\mathfrak{Z}_{\mathbb{C}}$ intersects all $X_{\mathbb{C}} \times (T_{z_1} \cap \dots \cap T_{z_j})$ properly $/\mathbb{R}$ and (ii) each $\partial_f^I \mathfrak{Z}_{\mathbb{C}}$ has the same property (in $X \times \square^{n-|I|}$).

We describe V -intersection chains first in the case where $V \subset X$ is a nonsingular K -subvariety of codimension q .

Definition 8.12. $Z_{\mathbb{R}}^p(X, n)_V$ has generators as in Def. 8.10, with the additional requirement that $\mathfrak{Z}_{\mathbb{C}}$ properly intersect all $V \times \partial_{f, \mathbb{R}}^{I, j} \square^n$ properly $/\mathbb{R}$. (For $Z^p(X, n)_V$ we just demand proper intersections with $V \times \partial_f^I \square^n$ in addition to $X \times \partial_f^I \square^n$.)

Remark 8.13. One can also define the “normalized” subcomplex $C^p(X, \bullet)_V \subset Z^p(X, \bullet)_V$ consisting of cycles with all facet intersections *zero* except

for that with $X \times \partial_0^n \square^n$; this subcomplex is well-known to be quasi-isomorphic. It is convenient to write $\partial^- \square^n := (\cup_{i=1}^{n-1} \partial_0^i \square^n) \cup (\cup_{i=1}^n \partial_\infty^i \square^n)$. Put $C_{\mathbb{R}}^p(X, n)_V := C^p(X, n) \cap Z_{\mathbb{R}}^p(X, n)_V$.

Lemma 8.14. (MOVING LEMMA I) *The inclusions*

$$\begin{array}{ccc} Z_{\mathbb{R}}^p(X, \bullet)_V & \hookrightarrow & Z^p(X, \bullet)_V \\ \downarrow & & \downarrow \\ Z_{\mathbb{R}}^p(X, \bullet) & \hookrightarrow & Z^p(X, \bullet) \end{array}$$

are all quasi-isomorphisms.

For the right vertical arrow, this is due to [Lv1, Cor 3.2].⁹ It will therefore suffice to examine the top horizontal inclusion, which we do in an appendix at the end of the subsection. Easy homological algebra shows one can tack a subscript “ $\mathbb{R}$ ” onto the moving lemmas of Bloch [Bl1] and Levine:

Corollary 8.15. *We have $\frac{Z_{\mathbb{R}}^p(X, \bullet)}{Z_{\mathbb{R}}^{p-q}(X, \bullet)} \simeq 0$, $Z_{\mathbb{R}}^p(X, \bullet)_V \xrightarrow{\simeq} Z_{\mathbb{R}}^p(X, \bullet)$ and $\frac{Z_{\mathbb{R}}^p(X, \bullet)}{Z_{\mathbb{R}}^{p-q}(D, \bullet)} \xrightarrow{\simeq} Z_{\mathbb{R}}^p(X \setminus D, \bullet)$ (for $D \subset X$ a closed subvariety of pure codimension q).*

The old moving lemmas can also be (with some more algebra) combined in the following way, which we will require for an application:

Lemma 8.16. (MOVING LEMMA II) *Given $D \subset X$ a NCD,*

$$\frac{Z^p(X, \bullet)_V}{Z^{p-1}(D, \bullet)_{V \cap D}} \xrightarrow{\simeq} Z^p(X \setminus D, \bullet)_{V \setminus V \cap D}$$

provided $V \cap D$ is proper and avoids the codimension-2 skeleton of D . (One may also replace Z by $Z_{\mathbb{R}}$ here.)

Next we want to augment V to a collection $\mathcal{S} := \{S_1, \dots, S_n\}$ of closed (possibly singular) subvarieties of X . By imposing the requirements of Def. 8.12 for each S_i one obtains $Z_{(\mathbb{R})}^p(X, \bullet)_{\mathcal{S}}$. All the moving lemmas still hold with $\mathcal{S}$ replacing V .

Given $V \subset X$ a NCD, write $\mathcal{V}^I$ for the collection of all V_J with $J \supsetneq I$ (see §8.1). We take $Z_V^{\ell, m}(p) := \oplus_{|I|=\ell} Z^p(V_I, -m)_{\mathcal{V}^I}$ with differentials $\partial_{\mathcal{B}} : Z_V^{\ell, m}(p) \rightarrow Z_V^{\ell, m+1}(p)$ and $\mathfrak{I} : Z_V^{\ell, m}(p) \rightarrow Z_V^{\ell+1, m}(p)$ (essentially identical to (8.1) above), and total differential $\mathbb{D}$. Since there is already a notion of $Z^p(V, -\bullet)$ (computing K' -theory), we set

$$\tilde{Z}^p((X, V), -\bullet) := s^\bullet(Z_V^{\bullet \geq 0, \bullet}(p)), \quad \tilde{Z}^p(V, -\bullet) := s^\bullet(Z_V^{\bullet \geq 1, \bullet}(p))[1].$$

⁹[Lv1] proves this for the normalized complexes; use Rem. 8.13 to transfer the result.

These have respective $(-n)^{\text{th}}$ cohomologies

$$H_{\mathcal{M}}^{2p-n}((X, V), \mathbb{Q}(p)) =: \widetilde{CH}^p((X, V), n)$$

and

$$H_{\mathcal{M}}^{2p-n}(V, \mathbb{Q}(p)) =: \widetilde{CH}^p(V, n)$$

($\neq CH^p(V, n)$ in general). One also defines $Z_{V, \mathbb{R}}^{\ell, m}(p)$ and $\widetilde{Z}_{\mathbb{R}}^p$'s in the obvious way, with the same cohomologies (by Moving Lemma I). Comments about cones as in §8.1 yield a version of (8.2) with $\mathcal{M}$ replacing $\mathcal{D}$. Note that negative n produces nontrivial $\widetilde{CH}$ groups in both cases: e.g. for $N > m$, $\widetilde{CH}^p((X \times \square^N, X \times \partial \square^N), -m) \cong CH^p(X, N - m) \cong \widetilde{CH}^p(X \times \partial \square^{N+1}, -m)$.

Finally, the following notion of relative cycles is convenient:

$$RZ^p(X, V) := \ker(\mathfrak{J}) \subseteq \{Z_V^{0,0}(p) = Z^p(X)_{\mathcal{V}^0}\}.$$

We also define

$$RZ^p(V) := \ker(\mathfrak{J}) \subseteq \{Z_V^{1,0}(p) = \oplus_{i=1}^N Z^p(V_i)_{\mathcal{V}^i}\}.$$

Obviously X, V need not be complete, and V need not have normal crossings (or live in some X) for these to be defined.

Appendix: proof of Moving Lemma I. In the following proof, we will abuse notation by writing $Z^p(X)_V$ for $Z^p(X)_{\mathcal{V}^0}$, i.e. using a NCD to denote the collection of its own “faces”; the same goes for “real analytic NCD’s”. So as a subscript, $\square_{\mathbb{R}}^n$ will mean the set of all $\partial_{f, \mathbb{R}}^{I, j} \square^n$ (including $|I|, j = 0$), $\partial \square^n$ the collection of $\partial_f^I \square^n$ ($|I| \geq 1$), and so on. We write $(\cdot)_X$ for $X \times (\cdot)$.

In the square of inclusions

$$\begin{array}{ccc} C_{\mathbb{R}}^p(X, \bullet)_V & \xrightarrow{\mathcal{I}} & Z_{\mathbb{R}}^p(X, \bullet)_V \\ \mathcal{J} \downarrow & & \mathcal{J}' \downarrow \\ C^p(X, \bullet) & \xrightarrow{\mathcal{I}'} & Z^p(X, \bullet), \end{array}$$

it is well-known that $\mathcal{I}'$ is a quasi-isomorphism. We will show here that **(a)** $\mathcal{J}$ is a quasi-isomorphism and **(b)** $\mathcal{I}$ is a quasi-surjection¹⁰ (in fact, it is a $\simeq$). It follows that $\mathcal{J}'$ is a quasi-isomorphism; applying this to

$$\begin{array}{ccc} Z_{\mathbb{R}}^p(X, \bullet)_V & \xrightarrow{\quad} & Z_{\mathbb{R}}^p(X, \bullet) \\ & \searrow \quad \swarrow & \\ & Z^p(X, \bullet) & \end{array}$$

¹⁰For $\alpha \in Z_{\mathbb{R}}^p(X, n)_{V, \partial_{\mathcal{B}}\text{-cl.}}$, $\exists \beta \in Z_{\mathbb{R}}^p(X, n+1)_V$ and $\gamma \in C_{\mathbb{R}}^p(X, n)_{V, \partial_{\mathcal{B}}\text{-cl.}}$ s.t. $\alpha = \partial_{\mathcal{B}}\beta + \mathcal{I}(\gamma)$.

for $V = V, \emptyset$ shows $Z_{\mathbb{R}}^p(X, \bullet)_V \xrightarrow{\simeq} Z_{\mathbb{R}}^p(X, \bullet)$ as required.

(a) We must show

$$\frac{RZ^p(\square_X^n, \partial \square_X^n)_{\square_V^n \cup \square_{\mathbb{R}, X}^n}}{\partial_0^{n+1} \left\{ RZ^p(\square_X^{n+1}, \partial^- \square_X^{n+1})_{\square_V^n \cup \square_{\mathbb{R}, X}^n} \right\}} \xrightarrow{\mathcal{J}} \frac{RZ^p(\square_X^n, \partial \square_X^n)}{\partial_0^{n+1} \left\{ RZ^p(\square_X^{n+1}, \partial^- \square_X^{n+1})_{\partial_0^{n+1} \square_X^{n+1}} \right\}}$$

is an isomorphism. This makes heavy use of [Lv1]. Set

$$\mathbb{I} := \{(I, f) \mid I \subseteq \{1, \dots, n\}, f : I \rightarrow \{0, \infty\}\},$$

and note $|\mathbb{I}| = 3^n$; [Lv1] constructs a scheme via iterated doubles, which as a topological space is

$$\mathcal{T}^n := \mathbb{I} \times \square_X^n \left/ \left\langle \begin{array}{l} (I, f, x) \sim (I', f', x') \text{ if} \\ I \subset I', f = f'|_I, x = x' \in \partial_{f'}^{I' \setminus I} \square_X^n \end{array} \right\rangle \right.$$

The preimage of a subvariety (or real analytic subspace) $W \subset \square_X^n$ under the obvious projection $\mathcal{T}^n \longrightarrow \square_X^n$, is denoted $\mathcal{T}^n(W)$. We may combine the inclusions $j_f^I : \square_X^n \hookrightarrow \mathcal{T}^n$ into

$$j^* := \sum_{I, f} (-1)^{|I|} (j_f^I)^* : Z^p(\mathcal{T}^n) \rightarrow Z^p(\square_X^n).$$

Given $\xi \in RZ^p(\square_X^n, \partial \square_X^n)$, Levine's argument in [Lv1, Thm. 2.3-4] (viewed through the lens of cycles, and his 2.7, 3.1) implies $\exists$ of $\alpha \in RZ^p(\square_X^{n+1}, \partial^- \square_X^{n+1})_{\partial_0^{n+1} \square_X^{n+1}}$, $f : \mathcal{T}^n \hookrightarrow H = \text{homogeneous space for } GL_n(K)$, and $\eta \in Z^p(H)_{f(\mathcal{T}^n)}$ such that the following holds. Writing $f^\sharp$ for the composition

$$Z^p(H)_{f(\mathcal{T}^n)} \xrightarrow{f^*} RZ^p(\mathcal{T}^n) \xrightarrow{j^*} RZ^p(\square_X^n, \partial \square_X^n),$$

$f^\sharp \eta = \xi + \partial_0^{n+1} \alpha$. Moreover, if $g \in GL_n(K)$ has $g^* \eta \in Z^p(H)_{f(\mathcal{T}^n)}$, then $f^\sharp g^* \eta = \xi + \partial_0^{n+1} \alpha'$.

Our point is that by extending scalars to some $L \supset K$, we can choose $g \in GL_m(L)$ so that $g^* \eta \in Z^p(H_L)_{f(\mathcal{T}^n)_L \cup f(\mathcal{T}^n(\square_V^n))_L \cup f(\mathcal{T}^n(\square_{\mathbb{R}, X}^n))}$, which implies $f^\sharp g^* \eta \in RZ^p(\square_{X_L}^n, \partial \square_{X_L}^n)_{\square_V^n \cup \square_{\mathbb{R}, X}^n}$. Namely, choose L such that $[L : L \cap \mathbb{R}] = 2$, and “realify” the entire situation, replacing all L -varieties by underlying $L \cap \mathbb{R}$ -varieties of twice the dimension and $GL_m(L)$ by its isomorphic copy in $GL_{2m}(L \cap \mathbb{R})$. (The realification of H_L is a homogeneous space for this isomorphic copy, not for $GL_{2m}(L \cap \mathbb{R})$.) Then the $f(\mathcal{T}^n(\square_{\mathbb{R}, X}^n))$ are now algebraic (our $L \cap \mathbb{R}$) and we may apply Kleiman transversality (as quoted in [Lv1]) to obtain g as described.

In fact, if $[L : K] < \infty$ then one can choose g such that

$$\sigma^*(g^*\eta) = (\sigma^*g)^*\eta \in Z^p(H_L)_{f(\mathcal{T}^n)_L \cup f(\mathcal{T}^n(\square_V^n))_L \cup f(\mathcal{T}^n(\square_{\mathbb{R},X}^n))}$$

$\forall \sigma \in \text{Gal}(L/K)$, so that

$$\text{Norm}(f^\sharp g^*\eta) := \frac{1}{[L : K]} \sum_{\sigma} f^\sharp(\sigma^*(g^*\eta)) \in RZ^p(\square_X^n, \partial \square_X^n)_{\square_V^n \cup \square_{\mathbb{R},X}^n},$$

completing the argument for surjectivity of $\mathcal{J}$ in that case. If L/K is merely finitely generated, then we can combine the norm argument with specialization of transcendentals.

For injectivity, given $\Xi \in RZ^p(\square_X^n, \partial \square_X^n)_{\square_V^n \cup \square_{\mathbb{R},X}^n}$ such that $\exists \alpha \in RZ^p(\square_X^{n+1}, \partial^- \square_X^{n+1})_{\partial_0^{n+1} \square_X^{n+1}}$ with $\Xi = \partial_0^{n+1} \alpha$, we construct $\beta \in RZ^p(\square_X^{n+1}, \partial^- \square_X^{n+1})_{\square_V^n \cup \square_{\mathbb{R},X}^n}$ with $\partial_0^{n+1} \beta = \Xi$. Write

$$\begin{aligned} \square_X^{n+1} \setminus |\Xi| \times \square &=: {}^\circ \square_X^{n+1}, & {}^\circ \partial \square_X^{n+1} &:= \partial \square_X^{n+1} \cap {}^\circ \square_X^{n+1}, \\ {}^\circ \mathcal{T}^{n+1} &:= \mathcal{T}^{n+1} \setminus \mathcal{T}^{n+1}(|\Xi| \times \square), \end{aligned}$$

etc.

The argument in the proof of [Lv1, Thm. 2.7] implies $\exists$ of $f : {}^\circ \mathcal{T}^{n+1} \hookrightarrow H[=\text{homogeneous space for } GL_m(K)]$ and $\eta \in Z^p(H)_{f({}^\circ \mathcal{T}^{n+1})}$, such that if $f^\sharp$ denotes the composition

$$\begin{aligned} Z^p(H)_{f({}^\circ \mathcal{T}^{n+1})} &\xrightarrow{f^*} RZ^p({}^\circ \mathcal{T}^{n+1}) \xrightarrow{j^*} RZ^p({}^\circ \square_X^{n+1}, {}^\circ \partial \square_X^{n+1}) \\ &\quad \begin{array}{c} \text{Zariski closure} \\ + \text{add degenerate} \\ \text{cycle} \end{array} \downarrow \\ &\quad RZ^p(\square_X^{n+1}, \partial^- \square_X^{n+1}), \end{aligned}$$

then $\partial_0^{n+1}(f^\sharp \eta) = \Xi$. Moreover, $\partial_0^{n+1} \circ f^\sharp$ of any g^* -translate of η also maps to Ξ .

Make an extension to L as before, choose a translate of η [with all $\text{Gal}(L/K)$ -conjugates] properly intersecting $f({}^\circ \mathcal{T}^{n+1}(\square_V^{n+1})) / L$ and (in the realification) $f(\mathcal{T}^{n+1}(\square_{\mathbb{R},X}^{n+1}))$, apply $f^\sharp$ and take norm; this yields β as desired.

(b) We first recall some definitions:

$$\begin{aligned} d^p(X, n) &:= \sum_{i=1}^n \text{im} \left\{ \pi_i^* : Z^p(\square_X^{n-1})_{\partial \square_X^{n-1}} \rightarrow Z^p(\square_X^n)_{\partial \square_X^n} \right\} \\ d_{\mathbb{R}}^p(X, n) &:= d^p(X, n) \cap Z^p(\square_X^n)_{\square_{\mathbb{R},X}^n} = \sum_{i=1}^n \pi_i^* \left(Z^p(\square_X^{n-1})_{\square_{\mathbb{R},X}^{n-1}} \right) \\ Z^p(X, n) &= \frac{Z^p(\square_X^n)_{\partial \square_X^n}}{d^p(X, n)}, \quad Z_{\mathbb{R}}^p(X, n) = \frac{Z^p(\square_X^n)_{\square_{\mathbb{R},X}^n}}{d_{\mathbb{R}}^p(X, n)}. \end{aligned}$$

Now given¹¹ $\mathcal{Z} \in Z_{\mathbb{R}}^p(X, n)$ with $\partial_{\mathcal{B}}\mathcal{Z} = 0$, let $\tilde{\mathcal{Z}}_1 \in Z^p(\square_X^n)_{\square_{\mathbb{R}, X}^n}$ be a representative, and define inductively

$$\tilde{\mathcal{Z}}_{i+1} := \tilde{\mathcal{Z}}_i - \pi_i^* \partial_{\infty}^i \tilde{\mathcal{Z}}_i \quad \text{for } i = 1, \dots, n.$$

This produces a representative $\mathfrak{Z} := \tilde{\mathcal{Z}}_{n+1} \in RZ^p(\square_X^n, \partial_{\infty}\square_X^n)_{\square_{\mathbb{R}, X}^n} \subset Z^p(\square_X^n)_{\square_{\mathbb{R}, X}^n}$, where $\partial_{\infty}\square_X^n := \cup_{j=1}^n \partial_{\infty}^j \square_X^n$.

Consider the projections

$$\begin{array}{ccc} \square_X^n & (x; z_1, \dots, z_{i-1}, z_i, z_{i+1}, z_{i+2}, \dots, z_n) & \\ \pi_{i+2} \uparrow & \uparrow & \\ \square_X^{n+1} & (x; z_1, \dots, z_{i-1}, z_i, z_{i+1}, z'_{i+1}, z_{i+2}, \dots, z_n) & \\ \pi_{\{i, i+1\}} \downarrow & \downarrow & \\ \square_X^{n-1} & (x; z_1, \dots, z_{i-1}, z'_{i+1}, z_{i+2}, \dots, z_n) & \end{array}$$

and set $Y_i := \{(1 - z_i)(1 - z_{i+1}) = (1 - z'_{i+1})\} \in Z^1(\square_X^{n+1})$. Sending $\mathcal{W} \mapsto (\pi_{i+2})_* \left(Y_i \cdot \pi_{\{i, i+1\}}^* (\partial_0^i \mathcal{W}) \right)$ yields a map

$$M_i \in \text{End} \left(RZ^p \left(\square_X^n, \cup_{j=1}^{i-1} \partial_0^j \square_X^n \bigcup \partial_{\infty} \square_X^n \right)_{\square_{\mathbb{R}, X}^n} \right).$$

(It is easy to check that real good position is preserved. For example, to show $\dim_{\mathbb{R}}(T_{z_1} \cap \dots \cap T_{z_j} \cap M_i(\mathcal{W})) \leq 2(\dim(X) + n - p) - j$, one uses the facts that $z_i, z_{i+1} \in \mathbb{R}^- \implies z'_{i+1} = 1 - (1 - z_i)(1 - z_{i+1}) \in \mathbb{R}^-$, and that $\partial_0^i \mathcal{W}$ is in good real position if $\mathcal{W}$ is.) Moreover,

$$\partial_0^i M_i(\mathcal{W}) = \partial_0^i \mathcal{W} = \partial_0^{i+1} M_i(\mathcal{W})$$

and so

$$\mathcal{W} - M_i(\mathcal{W}) \in RZ^p \left(\square_X^n, \cup_{j=1}^i \partial_0^j \square_X^n \bigcup \partial_{\infty} \square_X^n \right)_{\square_{\mathbb{R}, X}^n}.$$

We claim there is a cycle

$$\mathcal{H}_i(\mathcal{W}) \in RZ^p \left(\square_X^{n+1}, \cup_{j=1}^{i-1} \partial_0^j \square_X^{n+1} \bigcup \partial_{\infty} \square_X^{n+1} \right)_{\square_{\mathbb{R}, X}^{n+1}}$$

with

$$(8.3) \quad \partial_{\mathcal{B}} \mathcal{H}_i(\mathcal{W}) = (-1)^{i+1} M_i(\mathcal{W}) - \mathcal{H}_i(\partial_{\mathcal{B}} \mathcal{W}).$$

¹¹We omit the subscript V throughout, but all steps are still valid with this reinstated.

Consider more projections:

$$\begin{array}{ccc}
\Box_X^{n+1} & (x; z_1, \dots, z_{i-1}, z_i, z_{i+1}, z'_{i+1}, z_{i+2}, \dots, z_n) & \\
\pi_{i+3} \uparrow & & \uparrow \\
\Box_X^{n+2} & (x; z_1, \dots, z_{i-1}, z_i, z_{i+1}, z'_{i+1}, z''_{i+1}, z_{i+2}, \dots, z_n) & \\
\pi_{\{i, i+1, i+2\}} \downarrow & & \downarrow \\
\Box_X^{n-1} & (x; z_1, \dots, z_{i-1}, z''_{i+1}, z_{i+2}, \dots, z_n) &
\end{array}$$

and set $\mathcal{Y}_i := \{(1 - z_i)(1 - z_{i+1})(1 - z'_{i+1}) = (1 - z''_{i+1})\} \in Z^1(\Box_X^{n+2})$; then $\mathcal{H}_i(\mathcal{W}) := (\pi_{i+3})_* \left(\mathcal{Y}_i \cdot \pi_{\{i, i+1, i+2\}}^* (\partial_0^i \mathcal{W}) \right)$ does the job.

Now using (8.3) with $\partial_{\mathcal{B}} \mathcal{W} = 0$, inductively define $\tilde{\mathfrak{Z}}_1 := \mathfrak{Z}$ and

$$\tilde{\mathfrak{Z}}_{i+1} := \tilde{\mathfrak{Z}}_i - M_i(\tilde{\mathfrak{Z}}_i) = \tilde{\mathfrak{Z}}_i \pm \partial_{\mathcal{B}} \mathcal{H}_i(\tilde{\mathfrak{Z}}_i) \quad \text{for } i = 1, \dots, n-1.$$

Clearly $\tilde{\mathfrak{Z}}_n$ differs from $\mathfrak{Z}$ by $\partial_{\mathcal{B}}$ of a cycle in $Z^p(\Box_X^{n+1})_{\Box_{\mathbb{R}, X}^{n+1}}$, and belongs to $RZ^p(\Box_X^n, \partial^- \Box_X^n)_{\Box_{\mathbb{R}, X}^n}$; since $\tilde{\mathfrak{Z}}_n$ is $\partial_{\mathcal{B}}$ -closed, it actually lives in $RZ^p(\Box_X^n, \partial \Box_X^n)_{\Box_{\mathbb{R}, X}^n} = \ker(\partial_{\mathcal{B}}) \subseteq C_{\mathbb{R}}^p(X, n)$.

8.3. Abel-Jacobi maps. An irreducible $\mathfrak{Z} \in Z_{\mathbb{R}}^p(X, n)$ produces “KLM currents” $R_{\mathfrak{Z}}, \Omega_{\mathfrak{Z}}, \delta_{T_{\mathfrak{Z}}}$ [KLM, sec. 5.4]. For these to operate on continuous forms, the integrals described there must be bounded, and this in fact follows from the boundedness against C^∞ forms. Since d of these currents is given in terms of $R_{\partial_{\mathcal{B}} \mathfrak{Z}}, \Omega_{\partial_{\mathcal{B}} \mathfrak{Z}}, \delta_{T_{\partial_{\mathcal{B}} \mathfrak{Z}}}$, it follows that they are normal.

Let

$$\mathfrak{R}_{\mathfrak{Z}}^{i,j,I,f} := \left\{ \begin{array}{c} \text{subset of } X_{\mathbb{C}}^{an} \text{ over which } \mathfrak{Z}_{\mathbb{C}} \cap (X_{\mathbb{C}} \times \partial_{f, \mathbb{R}}^{I,j} \Box^n) \\ \text{is equidimensional of real dimension } i \end{array} \right\},$$

and consider $V \subset X$ smooth of codimension 1. To say $\mathfrak{Z} \in Z_{\mathbb{R}}^p(X, n)_V$, is equivalent to stipulating that all $\mathfrak{R}_{\mathfrak{Z}}^{i,j,I,f}$ properly intersect V . Roughly speaking, this ensures that $R_{\mathfrak{Z}}, \Omega_{\mathfrak{Z}}, \delta_{T_{\mathfrak{Z}}}$ have their most “serious” kinds of singularities or discontinuities (deltas, log-branch jumps) transversely to V . More precisely, the restriction of the KLM formula $\mathfrak{Z} \mapsto (-2\pi i)^{p+\bullet} ((2\pi i)^{-\bullet} T_{\mathfrak{Z}}, \Omega_{\mathfrak{Z}}, R_{\mathfrak{Z}})$ [KLM, sec. 5.5] to V -intersection higher Chow chains, produces a map of complexes

$$Z_{\mathbb{R}}^p(X, -\bullet)_V \xrightarrow{\widetilde{AJ}} C_{\mathcal{D}}^{2p+\bullet}(X, \mathbb{Q}(p))_V$$

commuting with the pullback. This extends automatically to $\text{codim}_X(V) > 1$ since the pullback of $\mathfrak{Z} \in Z_{\mathbb{R}}^p(X, n)_V$ under V -blowup $\sigma : \tilde{X} \rightarrow X$ belongs to $Z_{\mathbb{R}}^p(\tilde{X}, n)_{\tilde{V}}$.

The upshot of all this is the following

Proposition 8.17. *Let $V \subset X$ be a NCD with smooth components; assume X, V complete. The KLM formula induces a map of double complexes*

$$Z_{V, \mathbb{R}}^{\ell, m}(p) \xrightarrow{\widetilde{AJ}} \mathfrak{K}_V^{\ell, m}(p),$$

hence maps of total complexes

$$\tilde{Z}_{\mathbb{R}}^p((X, V), -\bullet) \rightarrow C_{\mathcal{D}}^{2p+\bullet}((X, V), \mathbb{Q}(p)), \quad \tilde{Z}_{\mathbb{R}}^p(V, -\bullet) \rightarrow C_{\mathcal{D}}^{2p+\bullet}(V, \mathbb{Q}(p))$$

computing AJ on motivic cohomology.

This yields a map of long-exact sequences

$$\begin{array}{ccccccc} \longrightarrow & \widetilde{CH}^p((X, V), n) & \longrightarrow & CH^p(X, n) & \longrightarrow & \widetilde{CH}^p(V, n) & \xrightarrow{n \rightarrow n-1} \\ & \downarrow AJ & & \downarrow AJ & & \downarrow AJ & \\ \longrightarrow & H_{\mathcal{D}}^{2p-n}((X, V), \mathbb{Q}(p)) & \longrightarrow & H_{\mathcal{D}}^{2p-n}(X, \mathbb{Q}(p)) & \longrightarrow & H_{\mathcal{D}}^{2p-n}(V, \mathbb{Q}(p)) & \longrightarrow \end{array}$$

Much of the foregoing construction extends to more general singular or relative varieties (over our field $K \subseteq \mathbb{C}$). Levine [Lv2, sec. IV.3] describes a procedure (partly due to Hanamura) for associating to any reduced finite-type K -scheme V a cohomological motive $\mathbb{Z}_V$ in $D_{mot}^b(Sm_K)$.¹² This is a diagram of motives of smooth quasi-projective varieties (arrows opposite to the actual morphisms), obtained via cubical hyperresolutions; it can be used to produce $\tilde{Z}_{(\mathbb{R})}^p(V, \bullet)$ computing $H_{\mathcal{M}}^*(V, \mathbb{Q}(p))$ ($V = \text{NCD}$ as done above is essentially a special case). What we have done for $H_{\mathcal{D}}$ (and $\widetilde{AJ}$) will also extend in a straightforward way under the following conditions. For each V_{α} occurring in the hyperresolution, (a) V_{α} is complete and (b) the irreducible components of images of morphisms (in the hyperresolution) into V_{α} are *locally* components of [coskeletons of] a NCD. Here are the two simplest examples that go beyond $V = \text{NCD}$; another is given in §8.4.

Example 8.18. V a degenerate elliptic curve of Kodaira type I_1 , i.e. a rational curve with ordinary double point p . In $D_{mot}^b(Sm_K)$,

$$\mathbb{Z}_V \cong \text{Cone}\{\mathbb{Z}_{\mathbb{P}^1} \xrightarrow{\iota_0^* - \iota_{\infty}^*} \mathbb{Z}_{\{p\}}\}[-1].$$

For us,

$$\tilde{Z}_{\mathbb{R}}^p(V, -\bullet) := \text{Cone}\{Z_{\mathbb{R}}^p(\mathbb{P}^1, -\bullet)_{\{0, \infty\}} \xrightarrow{\iota_0^* - \iota_{\infty}^*} Z_{\mathbb{R}}^p(\{p\}, -\bullet)\}[-1]$$

¹²triangulated motivic category, defined in [Lv2, I.2.1.4].

computes $H_{\mathcal{M}}^{2p+*}(V, \mathbb{Q}(p))$. If V is embedded in some smooth projective X , then motivic cohomology of (X, V) is computed by

$$\tilde{Z}_{\mathbb{R}}^p((X, V), -\bullet) := \text{Cone}\{Z_{\mathbb{R}}^p(X, -\bullet)_{\{V, p\}} \rightarrow \tilde{Z}_{\mathbb{R}}^p(V, -\bullet)\}[-1].$$

Example 8.19. V a $K3$ surface with rational singularity at p . There exists a minimal (crepant) resolution $b: \tilde{V} \rightarrow V$, with $b^{-1}(p) =: D$ an A - D - E curve [BHPV, sec. III.2], hence a NCD. The weak 2-resolution

$$\begin{array}{ccc} D & \xrightarrow{\iota_D} & \tilde{V} \\ \downarrow b|_D & & \downarrow b \\ p & \xrightarrow{\iota_p} & V \end{array}$$

produces an isomorphism $\mathbb{Z}_V \cong \text{Cone}\{\mathbb{Z}_{\tilde{V}} \oplus \mathbb{Z}_{\{p\}} \rightarrow \mathbb{Z}_D\}[-1]$ in $D_{mot}^b(Sm_K)$, and

$$\tilde{Z}_{\mathbb{R}}^p(V, -\bullet) := \text{Cone}\left\{Z_{\mathbb{R}}^p(\tilde{V}, -\bullet)_{\mathcal{D}^0} \oplus Z_{\mathbb{R}}^p(\{p\}, -\bullet) \xrightarrow{\iota_D^* - (b|_D)^*} \tilde{Z}_{\mathbb{R}}^p(D, -\bullet)\right\}[-1]$$

computes $H_{\mathcal{M}}$ of V .

In both examples, cochains for $H_{\mathcal{D}}$ are constructed similarly, and $\widetilde{AJ}$ goes through.

8.4. Regulator period computations on singular varieties. We start by setting up a pairing between homology and cohomology of (X, V) (a similar procedure works for V). Consider the double complexes $\mathcal{D}_V^{\ell, m}(p) := \oplus_{|I|=\ell} \mathcal{N}_{V_I}^{2p+m}\{V^I\}(V_I)$ with differentials $d, \mathfrak{I}$ and $\mathcal{D}_{\ell, m}^V(-p) := \oplus_{|I|=\ell} \mathcal{D}_{V_I}^{2d-2p-2\ell-m}(V_I)$ with differentials $d, Gy := 2\pi i \sum_{|I|=\ell} \sum_{i \in I} (-1)^{<i>_{I \setminus i}} (\iota_{V_I \subset V_{I \setminus i}})_*$. The associated simple complexes (with resp. total differentials $d \pm \mathfrak{I}, d \pm Gy$ both written $\mathbb{D}$)

$$C_{DR}^{2p+\bullet}((X, V), \mathbb{C}) := s^{\bullet}(\mathcal{D}_V^{\bullet, \bullet}(p)), \quad C_{2p+\bullet}^{DR}((X, V), \mathbb{C}) := s^{\bullet}(\mathcal{D}_{\bullet, \bullet}^V(-p))$$

will be our (co)chains; write Z_{DR}, Z^{DR} for the $\mathbb{D}$ -(co)cycles. In particular, put $Z_{2p+M}^{DR, \infty}((X, V), \mathbb{C}) :=$

$$\ker(\mathbb{D}) \cap \left\{ \oplus_{\ell} \left(\oplus_{|I|=\ell} \Omega_{V_I^{\infty}}^{2d-2p-\ell-M} \langle \log V^I \rangle (V_I) \right) \right\} \subset s^M(\mathcal{D}_{\bullet, \bullet}^V(-p))$$

for $\mathbb{D}$ -cycles given by C^{∞} log forms (modulo $\mathbb{D}$ -boundary, they all are). The entrywise pairings

$$\mathcal{N}_{V_I}^{2p+\ell+M}\{V^I\}(V_I) \otimes \Omega_{V_I^{\infty}}^{2(d-\ell)-(2p+\ell+M)} \langle \log V^I \rangle (V_I) \longrightarrow \mathbb{C}$$

(which exist by Rem. 8.3), *summed over the diagonal* (in the double complex), yield

$$Z_{DR}^{2p+M}((X, V), \mathbb{C}) \otimes Z_{2p+M}^{DR, \infty}((X, V), \mathbb{C}) \longrightarrow \mathbb{C},$$

which descends to

$$(8.4) \quad H_{DR}^{2p+M}((X, V), \mathbb{C}) \otimes H_{2p+M}^{DR}((X, V), \mathbb{C}) \longrightarrow \mathbb{C}.$$

Assume henceforth that $n \geq 1$, and also that $n \geq p$ or $p > d (= \dim(X))$. We describe in this case a (bilinear) “regulator period” map

$$(8.5) \quad \widetilde{CH}^p((X, V), n) \otimes H_{2p-n-1}^{DR}((X, V), \mathbb{Q}) \longrightarrow \mathbb{C}/\mathbb{Q}(p)$$

in terms of $\xi = \{\xi_I \in Z_{\mathbb{R}}^p(V_I, n + \ell)\}_{\substack{\ell, I \\ |I| = \ell}}$ and $\alpha = \{\alpha_I\}_{\ell, I} \in$

$Z_{2p-n-1}^{DR, \infty}((X, V), \mathbb{C})$ representing classes in the respective $\otimes$ -factors. Since $n \geq 1$, $F^p H^{2p-n}((X, V), \mathbb{C}) \cap H^{2p-n}((X, V), \mathbb{Q}) = 0$; since also $n \geq p$ or $p > d$, $H_D^{2p-n}((X, V), \mathbb{Q}(p)) \cong \frac{H^{2p-n-1}((X, V), \mathbb{C})}{H^{2p-n-1}((X, V), \mathbb{Q}(p))}$. So

$$\widetilde{AJ}(\xi) = \{(-2\pi i)^{p-n-\ell} ((2\pi i)^{n+\ell} T_{\xi_I}, \Omega_{\xi_I}, R_{\xi_I})\}_{\ell, I}$$

can be moved (with only rational ambiguity) by $\mathbb{D}$ -coboundary to

$$\mathcal{R} := \{(-2\pi i)^{p-n-\ell} (0, 0, R'_{\xi_I})\}_{\ell, I} \in \text{im} \{Z_{DR}^{2p-n-1}((X, V), \mathbb{C})\}.$$

Now (8.5) of $\xi \otimes \alpha$ is given by (8.4) of $\mathcal{R} \otimes \alpha$, which is

$$(-2\pi i)^{p-n} \sum_{\ell, I} \int_{V_I} R'_{\xi_I} \wedge \alpha_I.$$

By taking limits we can replace the $\log(V^I)$ C^∞ -forms α_I by relative topological cycles $\gamma_I \in Z_{2p-n-\ell-1}^{top}((V_I, V^I), \mathbb{Q})$. (Here $\gamma = \{\gamma_I\}$ gives a $\mathbb{D}$ -cycle in $C_{2p-n-1}^{DR}((X, V), \mathbb{C})$ by taking the associated δ -currents; it is these δ -currents which are limits of $\{\alpha_I^\epsilon\}$'s, and by definition $\int_\gamma R := \lim_{\epsilon \rightarrow 0} \int R \wedge \alpha^\epsilon$.) In fact, unless $p = n$ and $p \leq d$, the Ω_{ξ_I} are all 0. So the $\mathbb{D}$ -coboundary involves only topological chains, and

$$(8.6) \quad (-2\pi i)^{p-n} \int_\gamma R_\xi := (-2\pi i)^{p-n} \sum_\ell \sum_{|I|=\ell} \int_{\gamma_I} R_{\xi_I} \in \mathbb{C}/\mathbb{Q}(p)$$

computes (8.5). We find these periods in two examples, in which V replaces (X, V) and $p > \dim(V)$ suffices (in lieu of $p > \dim(X)$).


Example 8.20. $(\widetilde{CH}^2(—, 2) [\cong H_{\mathcal{M}}^2(—, \mathbb{Q}(2))]$ **of a singular curve.**)

Let ζ_1, ζ_2 be roots of unity (other than -1) with $\zeta_1 + \zeta_2 \neq 1$. In $\mathbb{A}^2$, let $V_1^\circ = \{x = \zeta_1\}$, $V_2^\circ = \{y = \zeta_2\}$, $V_3^\circ = \{x + y = 1\}$; their closures in $\mathbb{P}^2$ are denoted V_i . Set $V = \cup V_i$ and $V^\circ = V_1 \cup V_2 \cup V_3^\circ$; V is defined

(and we work) over $K = \mathbb{Q}(\zeta_1, \zeta_2)$. We first want to construct a higher Chow cocycle

$$\xi \in \ker(\mathbb{D}) \subset \tilde{Z}_{\mathbb{R}}^2(V, 2) = \oplus_i Z_{\mathbb{R}}^2(V_i, 2)_{\mathcal{V}^i} \bigoplus \oplus_{i < j} Z_{\mathbb{R}}^2(V_{ij}, 3),$$

and then use (8.6) to pair $\widetilde{AJ}(\xi)$ with the $\gamma = \{\gamma_I\}$ shown:



Denote the graph of (x, y) in $Z_{\mathbb{R}}^2(\mathbb{A}_K^2, 2)_{(\mathcal{V}^\circ)^\emptyset}$ by $\{x, y\}$; the *Tame* symbols of its restrictions to the η_{V_i} are torsion ($= 0$ for us), and its pullback ξ_3° to V_3° is $\partial_{\mathcal{B}}$ -closed. The Zariski closures of $\{x, y\}|_{V_3^\circ} =: \xi_i^\circ$ ($i = 1, 2$) may be “completed” to $\partial_{\mathcal{B}}$ -closed $\xi_i \in Z_{\mathbb{R}}^2(V_i, 2)_{\mathcal{V}^i}$ by adding appropriate α_1, α_2 supported resp. on $\{y = 0, \infty\} \subset V_1$, $\{x = 0, \infty\} \subset V_2$. (If $\zeta_i = 1$, there is nothing to do.) Hence we have

$$\xi^\circ := (\xi_1, \xi_2, \xi_3^\circ; 0, 0, 0) \in \tilde{Z}_{\mathbb{R}}^2(V^\circ, 2).$$

Applying Moving Lemma II to ξ_3° yields a cochain in $Z_{\mathbb{R}}^2(V_3, 2)_{\mathcal{V}^3}$ with restriction $\xi_3^\circ + \partial_{\mathcal{B}}\beta_3$ to V_3° ; we may complete it into a $\partial_{\mathcal{B}}$ -cocycle ξ_3 by adding α_3 supported on $\{(\infty, \infty)\} \in V_3$. Keeping track of $\mathbb{D}\beta_3$ (not just $\partial_{\mathcal{B}}\beta_3$) in this “move”, yields the desired

$$\xi := (\xi_1, \xi_2, \xi_3, \pm\beta_3|_{V_{13}}, \pm\beta_3|_{V_{23}}, 0),$$

with regulator period $\int_{\gamma} R_{\xi} = \{\sum_{i=1}^3 \int_{\gamma_i} R_{\xi_i}\} \pm \frac{1}{2\pi i} \{R_{(\beta|_{V_{13}})} + R_{(\beta|_{V_{23}})}\}$. But general principles (Stokes’s theorem for currents) ensure that R_{ξ° must have (mod $\mathbb{Q}(2)$) the same period, which is easy to compute:

$$\begin{aligned} \int_{\gamma} R_{\xi^\circ} &= \sum_{i=1}^3 \int_{\gamma_i} R\{x, y\} \\ &= \int_{y=1-\zeta_1}^{y=\zeta_2} \log \zeta_1 d\log y + \int_{x=\zeta_1}^{x=1-\zeta_2} (-2\pi i) \log \zeta_2 \delta_{T_x} \\ &\quad + \int_{y=\zeta_2}^{y=1-\zeta_1} \{\log(1-y) d\log y - 2\pi i \log y \delta_{T_{1-y}}\} \\ &= -\log \zeta_1 \log(1-\zeta_1) + 0 + \{\text{Li}_2(\zeta_2) - \text{Li}_2(1-\zeta_1)\} \\ &\equiv \text{Li}_2(\zeta_1) + \text{Li}_2(\zeta_2) \in \mathbb{C}/\mathbb{Q}(2). \end{aligned}$$

This really boils down to a (lifted) Borel regulator map computation for an element of $K_3^{\text{ind}}(K)$, in the sense that one may explicitly construct

ρ so that

$$\begin{array}{ccc} H_{\mathcal{M}}^2(V, \mathbb{Q}(2)) & \xrightarrow[\cong]{\rho} & CH^2(K, 3) \\ \downarrow AJ & & \downarrow AJ \\ H_D^2(V, \mathbb{Q}(2)) & \xleftarrow[\cong]{} H^1(V, \mathbb{C}/\mathbb{Q}(2)) \xrightarrow[\cong]{f_\gamma} & \mathbb{C}/\mathbb{Q}(2) \end{array}$$

commutes.


Example 8.21. $(\widetilde{CH}^3(\text{---}, 3)$ of a singular surface.)

In $(\mathbb{P}^1)^3$, let $V_1 = \{x = 1\}$, $V_2 = \{y = 1\}$, $V_3 = \{z = 1\}$, $V_4 = \{z = \frac{(x-1)(y-1)}{xy}\}$, and $V := \cup V_i$. This is not quite a NCD as V_{14} and V_{24} are reducible. For this reason we write $V^4 (= V_{14} \cup V_{24} \cup V_{34}) =: W = \cup W_j$, with

$$W_1 = \{x = 1, z = 0\}, \quad W_2 = \{y = 1, z = 0\}, \quad W_3 = \{x = 1, y = 0\}, \\ W_4 = \{x = 0, y = 1\}, \quad \text{and } W_5 = \{z = 1\} \cap V_4.$$

Let $D = D_1 \cup D_2$ where $D_1 := \{x = 0, z = \infty\} \subset V_4$, $D_2 := \{y = 0, z = \infty\} \subset V_4$, and set $V_4^\circ := V_4 \setminus D$, $W_j^\circ := W_j \cap V_4^\circ$, $W^\circ = \cup W_j^\circ$. Finally let $C_1 = \{x = \infty\} \cap V_4$, $C_2 = \{y = \infty\} \cap V_4$. We work over $K = \mathbb{Q}$.

We will compute the regulator period of $\xi \in \ker(\mathbb{D}) \subset \widetilde{Z}_{\mathbb{R}}^3(V, 3)$ over the topological 2-cycle $\gamma = \{\gamma_I\}$ with $\gamma_1 = \{1\} \times [0, 1] \times [0, 1]$, $\gamma_2 = [0, 1] \times \{1\} \times [0, 1]$, $\gamma_3 = (\{x + y \geq 1\} \subset [0, 1]^2) \times \{1\}$, $\gamma_4 = (\{x + y \geq 1\} \subset [0, 1]^2) \cap V_4$.



To construct ξ , the idea is to move and complete the pullback of $\{x, y, z\} \in Z_{\mathbb{R}}^3((\mathbb{G}_m)^3, 3)$ to $\widetilde{Z}_{\mathbb{R}}^3(V \cap (\mathbb{G}_m)^3, 3)$ (which has trivial *Tame* symbols on components). Since $\{x, y, z\}$ is identically zero on V_1 , V_2 , and V_3 , we do this construction in the subcomplex $\widetilde{Z}_{\mathbb{R}}^3((V_4, W), -\bullet)$ of $\widetilde{Z}_{\mathbb{R}}^3(V, -\bullet)$.

The images of the maps

$$\mathbb{P}^1 \setminus \{0\} \times \mathbb{P}^1 \setminus \{0, 1, \infty\} \hookrightarrow V_4^\circ \times \square^3$$

$$\begin{aligned}
(t, u) &\mapsto (\infty, t, 1 - t^{-1}; u^{-1}, 1 - \frac{1}{tu}, 1 - u) \\
&\mapsto (\infty, t, 1 - t^{-1}; u^{-1}, 1 - \frac{1}{u}, 1 - u) \\
&\mapsto (t, \infty, 1 - t^{-1}; u^{-1}, 1 - \frac{1}{tu}, 1 - u) \\
&\mapsto (t, \infty, 1 - t^{-1}; u^{-1}, 1 - \frac{1}{u}, 1 - u)
\end{aligned}$$

yield higher Chow cochains $\mathcal{A}_1, \mathcal{A}_2$ resp. $\mathcal{A}_3, \mathcal{A}_4$ supported on C_1 resp. C_2 , such that $\xi_4^\circ := \{x, y, z\}|_{V_4^\circ} - \mathcal{A}_1 + \mathcal{A}_2 + \mathcal{A}_3 - \mathcal{A}_4$ is $\partial_{\mathcal{B}}$ -closed on V_4° , and

$$\xi_{rel}^\circ := (\xi_4^\circ; 0, \dots, 0) \in \ker(\mathbb{D}) \subset \tilde{Z}_{\mathbb{R}}^3((V_4^\circ, W^\circ), 3).$$

That one (if so inclined) can then *repeatedly* apply the Moving Lemma II to produce $\xi_{rel} \in \ker(\mathbb{D}) \subset \tilde{Z}_{\mathbb{R}}^3((V_4, W), 3)$, is a consequence of exactness of

(8.7)

$$H_{\mathcal{M}}^3((V_4, W), \mathbb{Q}(3)) \longrightarrow H_{\mathcal{M}}^3((V_4^\circ, W^\circ), \mathbb{Q}(3)) \xrightarrow{Res}$$

$$\{H_{\mathcal{M}}^2((D_2 \setminus D_{12}, D_2 \cap W_3), \mathbb{Q}(3)) \oplus H_{\mathcal{M}}^2((D_1 \setminus D_{12}, D_1 \cap W_4), \mathbb{Q}(3))\}$$

at the middle term. We could compute $Res(\xi_{rel}^\circ)$, but this is unnecessary: middle-exactness of

$$H_{\mathcal{M}}^1(\{1\}, \mathbb{Q}(2)) \rightarrow H_{\mathcal{M}}^2((\mathbb{A}_{\mathbb{Q}}^1, \{1\}), \mathbb{Q}(2)) \rightarrow H_{\mathcal{M}}^2(\mathbb{A}_{\mathbb{Q}}^1, \mathbb{Q}(2))$$

and vanishing of $CH^2(\mathbb{A}_{\mathbb{Q}}^1, 2) \cong K_2^M(\mathbb{Q})_{\mathbb{Q}}$ and $CH^2(Spec(\mathbb{Q}), 3)$, imply the second line of (8.7) is zero.

To compute $\int_{\gamma} R_{\xi}$, note that by construction it is just $\int_{\gamma_4} R_{\xi_{rel}}$, which must be $\int_{\gamma_4} R_{\xi_{rel}^\circ} = \int_{\gamma_4} R\{x, y, z\}$ by (repeated) application of Stokes. That is, $\int_{\gamma} R_{\xi} =$

$$\int_{\gamma_4} \left\{ \log(x) d\log(y) \wedge d\log(z) + 2\pi i \log(y) d\log(z) \delta_{T_x} + (2\pi i)^2 \log(z) \delta_{T_x \cap T_y} \right\}.$$

Writing $\Delta := \{x + y \geq 1\} \subset [0, 1]^2$, the latter is simply

$$\int_{\Delta} \log(x) d\log(y) \wedge d\log\left\{\frac{(x-1)(y-1)}{xy}\right\}.$$

since $\gamma_4 \cap T_x \subset W_4$ (and $\log(y) d\log(z)$, $\log(z) \delta_{T_y}$ are zero on W_4). So we get

$$\int_{\Delta} \log(x) d\log(y) \wedge d\log\left\{\frac{(x-1)}{x}\right\}$$

$$\begin{aligned}
&= - \int_{x=0}^1 \log(x) \left(\int_{y=1-x}^1 d\log y \right) [\mathrm{dlog}(x-1) - \mathrm{dlog}(x)] \\
&= \int_{x=0}^1 \log(x) \log(1-x) [\mathrm{dlog}(1-x) - \mathrm{dlog}(x)] \\
&= -2 \int_{u=0}^1 \log(u) \log(1-u) \mathrm{dlog}(u) \\
&= 2 \sum_{n>0} \frac{1}{n} \int_0^1 u^{n-1} \log(u) du = 2 \sum_{n>0} \frac{1}{n} \left(-\frac{1}{n^2} \right) = -2\zeta(3)
\end{aligned}$$

in $\mathbb{C}/\mathbb{Q}(3)$.

This can be interpreted as a Borel regulator computation (for $K_5(\mathbb{Q})_{\mathbb{Q}}^{(3)}$) as follows. The element $\Xi := \left\{ \frac{z}{z-1}, \frac{(x-1)(y-1)}{(x-1)(y-1)-xyz}, \frac{x}{x-1} \right\} \in CH^3((\mathbb{P}^1)^3 \setminus V, 3)$ has $Res_V(\Xi) \in CH^2(V, 2)$ with cycle-class in $Hom_{\mathrm{MHS}}(\mathbb{Q}(-3), H_V^4((\mathbb{P}^1)^3, \mathbb{Q}))$ given by ∂T_{Ξ} — which is precisely γ . Hence integration against γ splits $H^2(V)$, that is, gives a morphism of MHS $H^2(V, \mathbb{Q}) \rightarrow \mathbb{Q}(0)$. One may construct a compatible “splitting” of motivic cohomology, in the sense that

$$\begin{array}{ccc}
H_{\mathcal{M}}^3(V, \mathbb{Q}(3)) & \xrightarrow{\rho} & CH^3(\mathrm{Spec}(\mathbb{Q}), 5) \\
\downarrow AJ & & \downarrow AJ \\
H_D^3(V, \mathbb{Q}(3)) & \xleftarrow{\cong} H^2(V, \mathbb{C}/\mathbb{Q}(3)) \xrightarrow{\int_{\gamma}} & \mathbb{C}/\mathbb{Q}(3)
\end{array}$$

commutes. This phenomenon will be treated in a subsequent work.

REFERENCES

- [Ar1] D. Arapura, *The Leray spectral sequence is motivic*, Invent. Math. (2005)
- [Ar2] ———, *Notes on D-modules and mixed Hodge modules*, preprint, 2006.
- [As] M. Asakura, *Motives and algebraic de Rham cohomology*, in “The arithmetic and geometry of algebraic cycles” (Banff), CRM Proc. Lect. Notes 24, AMS, 2000, 133-154.
- [BHPV] W. Barth, K. Hulek, C. Peters, A. van de Ven, “Compact Complex Surfaces” (2nd ed.), Springer-Verlag, Berlin, 2004.
- [Be] A. Beilinson, *Notes on absolute Hodge cohomology*, Contemp. Math. 55 (1986), 35-68.
- [Bl1] S. Bloch, *The moving lemma for higher Chow groups*, J. Alg. Geom. 3 (1994), no. 3, 493-535.
- [Bl2] ———, *Algebraic cycles and the Beilinson conjectures*, Contemp. Math. 58 (1986), 65-79.
- [Bl3] ———, *Some elementary theorems about algebraic cycles on Abelian varieties*, Invent. Math. 37 (1976), 215-228.

- [BZ] J.-L. Brylinski and S. Zucker, *An overview of recent advances in Hodge theory*, Several complex variables VI, Encyclopedia Math. Sci. 69 (Springer, 1990), 39-142.
- [CF] A. Collino and N. Fakhruddin, Indecomposable higher Chow cycles on Jacobians
- [De1] P. Deligne, *Theorie de Hodge II*, Publ. Math. IHES 40 (1971), 5-58.
- [De2] ———, *Theorie de Hodge III*, Publ. Math. IHES 44 (1974), 5-77.
- [DK] C. Doran and M. Kerr, *Algebraic K-theory of toric hypersurfaces*, in preparation.
- [GG] M. Green and P. Griffiths, *Hodge-theoretic invariants for algebraic cycles*, Intl. Math. Res. Not. 2003, no. 9, 477-510.
- [Gr] P. Griffiths, *Some results on algebraic cycles on algebraic manifolds*, Algebraic Geometry (Bombay, 1968), Oxford Univ. Press, London, 1969, 93-191.
- [Ja1] U. Jannsen, *Deligne homology, Hodge-D-conjecture, and motives*, in “Beilinson’s Conjectures on Special Values of L -functions” (Academic Press, Boston, MA, 1988), 305-372.
- [Ja2] ———, “Mixed motives and Algebraic K -theory”, Lect. Notes in Math., vol. 1400, Springer, Berlin, 1990.
- [Ke1] M. Kerr, *Higher Abel-Jacobi maps for 0-cycles*, to appear in K -theory.
- [Ke2] ———, *A survey of transcendental methods in the study of Chow groups of zero-cycles*, to appear in “Mirror Symmetry V”.
- [Ke3] ———, *Exterior products of zero-cycles*, to appear in J. reine angew. Math.
- [Ke4] ———, *A regulator formula for Milnor K -groups*, K -theory 29 (2003), pp. 175-210.
- [Ke5] ———, *Motivic irrationality proofs*, in preparation.
- [KLM] M. Kerr, J. Lewis, S. Müller-Stach, *The Abel-Jacobi map for higher Chow groups*, Compos. Math. 142 (2006), no. 2, 374-396.
- [Ki] J. King, *Log complexes of currents and functorial properties of the Abel-Jacobi map*, Duke Math. J. 50 (1983), no. 1, pp. 1-53.
- [Le1] J. Lewis, *A filtration on the Chow groups of a complex projective variety*, Compos. Math. 128 (2001), no. 3, 299-322.
- [Le2] ———, *A note on indecomposable motivic cohomology classes*, J. reine angew. Math. 485 (1997), 161-172.
- [Le3] ———, “A survey of the Hodge conjecture” (2nd edition), AMS, Providence, 1999.
- [Lv1] M. Levine, *Bloch’s higher Chow groups revisited*, in “ K -theory (Strasbourg, 1992)”, Asterisque No. 226 (1994), 10, pp. 235-320.
- [Lv2] ———, “Mixed Motives”, AMS, Providence, RI, 1998.
- [RS] A. Rosenschon and M. Saito, *Cycle map for strictly decomposable cycles*, Amer. J. Math. 125 (2003), no. 4, 773-790.
- [SaM1] M. Saito, *Extension of mixed Hodge modules*, Compos. Math. 74 (1990), 209-234.
- [SaM2] ———, *Mixed Hodge modules*, Publ. RIMS, Kyoto Univ., 26 (1990), 221-333.
- [SaM3] ———, *Direct image of logarithmic complexes and infinitesimal invariants of cycles*, preprint 2005.

- [SaM4] ———, Hodge-type conjecture for higher Chow groups, arXiv:math.AG/0201113 v4.
- [SaS] S. Saito, *Higher normal functions and Griffiths groups*, J. Alg. Geom. 11 (2002), no. 1, pp. 161-201.
- [Z] S. Zucker, *Hodge theory with degenerating coefficients: L_2 cohomology in the Poincaré metric*, Ann. of Math. 109 (1979), 415-476.

Matt Kerr matkerr@math.uchicago.edu
 Department of Mathematics, University of Chicago, Chicago, IL 60637,
 USA

James D. Lewis lewisjd@gpu.srv.ualberta.ca
 Department of Mathematics, University of Alberta, Edmonton, Al-
 berta, T6G 2G1, Canada